

Lipids, Biochemicals, and Standards for
Life Science Research

2009/2010

About Matreya LLC

- Matreya strives to develop, manufacture and deliver products of the highest value to our customers.
- Quality will always be the best achievable by state-of-the-art techniques, typically greater than 98%
- We strive for rapid delivery. 95+% of our products are shipped within 24 hours of receipt of an order.
- Within the area of sphingolipids and glycolipids, we have earned a reputation as the preferred problem solver and technology leader.
- When you demand quality and consistency, you may rely on Matreya lipids.

Matreya Products for Biochemistry Research.

We offer one of the widest selection of ceramides for intracellular signaling research available. We stock antibodies to glycosphingolipids as well as inhibitors of enzymes involved in glycosphingolipid metabolism.

Our products provide the valuable tools for the study of cell membrane and its structure, growth regulators in the cellular metabolism, and intracellular mediators.

We are able to make our products better and better with the latest technology in Chromatography, Mass Spectrometry, and NMR techniques.

We are proud to offer our products as a valuable tool for your life science research needs.

Matreya Products for Microbiology Research.

Matreya stocks many unusual fatty acid standards produced by bacteria that are useful for culture characterization.

Matreya Products for the Food and Agriculture Industries.

Many of Matreya's fatty acid products have been industry standards for many years. The acids and their methyl esters are used as standards in analysis and quality control.

Custom Preparations.

Our experience in chemical synthesis and the extraction and purification of natural products allows us to produce custom preparations with the same high quality and purity as the products listed in the catalog. Depending on the complexity of the molecule, delivery will be 4 to 12 weeks after receipt of an order, usually less than 6 weeks.

If you can't find a product in the catalog, please check the INDEX, where we also try to list common synonyms for our products.

TABLE OF CONTENTS

TABLE OF CONTENTS	i
Technical Service	iii
Natural Products.....	iii
Storage.....	iii
Sphingolipid Structures and Pathways.....	iii
Package Weight	iii
Sphingoid bases, sphingolipids and glycosphingolipids	1
Sphingosines.....	2
Synthetic sphingosines with C18 sphingoid base.....	2
Synthetic sphingosines with sphingoid bases other than C18.....	2
Synthetic dihydrosphingosines	3
3-keto-Dihydrosphingosines	5
Phytosphingosines.....	5
Other sphingosine derivatives and precursors	6
Ceramides	7
Synthetic ceramides derived from C18-sphingosine	7
2-Hydroxy ceramides	10
Ceramide made from sphingosines with sphingoid bases other than C18	10
Dihydroceramides	11
2-Hydroxy dihydroceramides	11
Ceramides from natural sources.....	12
Phytoceramides	13
Fluorescent ceramides.....	14
Phosphosphingolipids	15
Sphingomyelins.....	15
Sphingosylphosphorylcholines (SPC).....	17
Sphingosine phosphates.....	18
Fluorescent sphingomyelins	18
Glycosphingolipids	19
Galactosylceramides and glucosylceramides.....	19
Lactosylceramides	23
Ceramide trihexosides	24
Globosides	25
Labeled glycolipids.....	25
Gangliosides	28
Glycosphingolipid reference mixes for TLC	31
Antibodies directed against glycolipids	32
Enzyme Inhibitors	34
Glycerolipids	40
Glycerophospholipids.....	40
Natural phospholipids	40
Synthetic phospholipids	42
Phosphatidylinositols	45
Bacterial tetraethers.....	47
Glycosyl glycerides.....	48
Fatty acids	48
Simple fatty acids	48
Saturated fatty acids and methyl esters.....	48
Unsaturated fatty acids and methyl esters.....	52
Trans fatty acids and methyl esters	57
Conjugated linoleic acid isomers (CLA)	59
Other CLA products and derivatives.....	61

Hydroxy fatty acids	61
2-Hydroxy fatty acids and methyl esters	61
3-Hydroxy fatty acids and methyl esters	64
Omega hydroxy fatty acids	66
Other hydroxy fatty acids	68
Branched and cyclic fatty acids	69
iso-Fatty acids and esters	69
anteiso-Fatty acids and esters	70
Methylated fatty acids	70
Cyclopropyl fatty acids and esters	71
Unusual fatty acids and derivatives	71
Other lipids	72
Tocopherols	72
Cholestane derivatives	73
Plant sterols and steryl glucosides	74
Propyleneglycol Monoesters	75
Standards and reference compounds	75
Food industry mixes	75
Each methyl ester mix is carefully prepared by weight.	75
Polyunsaturated fatty acid methyl esters mixes	76
Carbohydrate mixes	76
Other fatty acid methyl ester mixes	77
AOCS animal and vegetable oil reference mixes (RM mixes).....	78
Custom mixes	80
GLC Standard mixes.....	81
Water soluble fatty acid mixes	82
Microbiology standard mixes	83
Biochemical research standard mixes	83
Glycosphingolipid mixtures for TLC	84
Biochemicals and reagents	85
Stable isotope labeled compounds	85
Fluorescent compounds.....	86
Sphingolipid Structures and Pathways Wall Chart.....	95
Cross reference for product numbers and catalog pages.....	96
Product name index	100
International Dealers & Representatives.....	107
Tables	
Table I. AOCS Oil Reference Mixes.....	78
Table II. Standards for GC analysis	81
Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC.	90

All chemicals listed in this catalog are for investigational use only. They are not intended for human consumption or to be used in food or food additives. None are for general drug or medicinal use on humans. We believe the information in this catalog, offered in good faith, is accurate.

Limited Warranty: All Matreya Products, except those specifically exempted, are warranted (for 90 days) to be free of defects in materials and workmanship, if properly stored. Any replacements required as a result of such defects will be made without charge provided that such defective products are returned with a written explanation. Please request a Returned Goods Authorization before returning products under this warranty.

Technical Service

Our technical service department may be contacted by telephone at 800.342.3595, or by e-mail to techservice@matreya.com.

Note that there are excellent resources for general information on lipid nomenclature, distribution and biochemistry available on the internet. One such source is www.lipidlibrary.co.uk maintained by W.W. Christie and colleagues. Another source for detailed discussion on isolation and purification protocols is www.cyberlipid.org. This site also maintains discussion groups for lipid researchers.

Natural Products

Some of our glycolipids are extracted from natural sources. These products have a normal heterogeneity in their lipid components, particularly in the fatty acids. Variations include carbon chain length as well as the presence or absence of 2-hydroxy fatty acids. Products based on sphingosine may contain longer chain sphingoid bases as well as chains with multiple double bonds. This heterogeneity may result in additional spots showing on TLC plates or multiple peaks in LC analyses. We have listed the typical fatty acid compositions of our natural products in the appendix.

Storage

Catalog items in unopened containers are stable for at least one year when stored under the conditions indicated in the catalog listing. Items containing unsaturated fatty acids are subject to oxidation and should be stored in solution in organic solvents or under argon. Glycolipids and phospholipids should not be stored in aqueous solutions due to potential hydrolysis.

Sphingolipid Structures and Pathways

In a clear and straightforward manner, this wall chart indicates the structures and relationships between most commonly discussed sphingolipids. A one-page thumbnail version of the chart is shown on page 95. Full size copies (approximately 35 x 26 inches) are available on request to customer service.

Package Weight

Unless otherwise specified, the package will contain at least the indicated amount and usually slightly more. The user is cautioned to always measure the required amount from the container.


Sphingoid bases, sphingolipids and glycosphingolipids.

Sphingoid bases such as sphingosine are the characteristic structural unit of the sphingolipids. The bases are long chain aliphatic amines, containing two or three hydroxyl groups, and typically a *trans*-double bond at C4. In animal tissues most abundant base is sphingosine with a C18 aliphatic chain containing a double bond in position 4. The saturated analogue is dihydrosphingosine or sphinganine. In plants the common long chain base is the 4 hydroxy saturated base phytosphingosine.

Sphingolipids are widely distributed in animal tissues, particularly cell membranes. Sphingoid bases linked to fatty acids via an amide bond at C2 are ceramides and are present in trace amounts in most tissues. Glycosphingolipids (ceramides having various mono- and oligosaccharides on the OH group at C1) are neutral glycosphingolipids (i.e., cerebrosides and globosides). Those with sialic acid derivatized sugars are acidic glycolipids (i.e., gangliosides). They are amphiphilic and can be solubilized in buffers via sonication and micelle formation.


Gangliosides are present in substantial amounts in nerve cell membranes, and together with globosides are found in the membranes of white and red blood cells. These plus the glycosphingolipids of the lacto- and neolacto-series are involved in cell recognition (e.g. blood group determinants). Glycolipid expression on the surface of cells determines their antigenicity as well as their status, i.e. differentiated vs. undifferentiated (embryonic), normal vs. malignant, etc. (1). The ganglioside GM1 stimulates nerve growth (2,3) and has been reported to have a curative effect on experimental Parkinsonism (4). For an overview see (5). Gangliosides are also being investigated as potential anti-tumor vaccines (57). Glycosphingolipids are also essential for the correct functioning of cell surface receptors (6). Matreya is your best source for many sphingolipids. Most of Matreya's sphingosines and ceramides are fully synthetic and as such 98%+ pure. Others, particularly the glycosphingolipids are highly purified natural products (98%+), and can be used either as standards or biochemical reagents without further purification.

Through total synthesis, all four isomers of sphingosine are available as well as a number of sphingosines with other than 18 carbons and a number of ceramides (for details in using ceramides in cell culture see Hauser et al. [9]). Fluorescent labeled ceramides, glycosphingolipids and sphingomyelins are also available for study. D. N. Brindley and his group have been exploring the interaction of ceramides, sphingosine and sphingosine 1-phosphate in regulating DNA synthesis and phospholipase D activity (10). **See Literature References on page 96.**



Sphingosines

Synthetic sphingosines with C18 sphingoid base



1802 D-erythro-Sphingosine 25 mg

Sphingosine, C18 chain $C_{18}H_{37}NO_2$ CAS#: 123-78-4

Source: synthetic **Mol. Wt.:** 299 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** alcohols, chloroform, DMSO **Storage:** -20°C

Selective inhibitor of phosphokinase C

References:

Y. Hannun et al., *Science*, **235**, 670, 1987

S. Spiegel et al., *Proc. Intern. Conf. Biol. Function Glycosphingolipids*, Santa Barbara, CA 1990

1806 L-threo-Sphingosine 10 mg

L-threo-Sphingosine, C18 chain $C_{18}H_{37}NO_2$

Source: synthetic **Mol. Wt.:** 299 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** -20°C

1826 L-erythro-Sphingosine 5 mg

L-erythro-Sphingosine, C18 chain $C_{18}H_{37}NO_2$ CAS#: 6036-75-5

Source: synthetic **Mol. Wt.:** 299 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** -20°C

1827 D-threo-Sphingosine 5 mg

D-threo-Sphingosine, C18 chain $C_{18}H_{37}NO_2$ CAS#: 6036-85-7

Source: synthetic **Mol. Wt.:** 299 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** -20°C

1304 Sphingosine 10 mg

D-erythro-Sphingosine $C_{18}H_{37}NO_2$ CAS#: 123-78-4

Source: semi-synthetic, bovine **Mol. Wt.:** 299 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** alcohol, chloroform **Storage:** -20°C

Synthetic sphingosines with sphingoid bases other than C18


Varying chain lengths allow the study of translocation effects of sphingosines and ceramides into cells.

1833 D-erythro-C14-Sphingosine 5 mg

Sphingosine with C14 chain $C_{14}H_{29}NO_2$


Source: synthetic **Mol. Wt.:** 243 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** -20°C

- 1835 D-erythro-C16-Sphingosine** **5 mg**
Sphingosine with C16 chain $C_{16}H_{33}NO_2$
- Source:** synthetic **Mol. Wt.:** 271 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$
- 1837 D-erythro-C10-Sphingosine** **5 mg/ml, 1 ml**
Sphingosine with C10 chain $C_{10}H_{21}NO_2$
- Source:** synthetic **Mol. Wt.:** 187 **Purity:** 98+% by TLC, GC **Appearance:** liquid **Solvent:** ethanol **Solubility:** ethanol **Storage:** $-20^{\circ}C$
- 1838 D-erythro-C12-Sphingosine** **5 mg**
Sphingosine with C12 chain $C_{12}H_{25}NO_2$ CAS#: 6918-49-6
- Source:** synthetic **Mol. Wt.:** 215 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$
- 1840 D-erythro-C20-Sphingosine** **5 mg**
Sphingosine with C20 chain $C_{20}H_{41}NO_2$
- Source:** synthetic **Mol. Wt.:** 328 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$



Synthetic dihydrosphingosines

D,L-*threo*-Dihydrosphingosine has also been found to be a significant inhibitor of sphingosine kinase (11). The D,L-*erythro*-isomer has been used as an inactive control. We now offer all four isomers in pure form making detailed studies possible. Safingol, the L-*threo*-isomer is a potent inhibitor of PKC and as such is capable of reversing multi-drug resistance in cancer cells (12). **See Literature References on page 96.**




- 1807 L-threo-Dihydrosphingosine (Safingol)** **5 mg**
1807-025 L-threo-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ CAS#: 15639-50-6 **25 mg**
- Source:** synthetic **Mol. Wt.:** 301 **Melting Point ($^{\circ}C$):** 103-114 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$
- References:**
C.W. Sachs et al., *ibid.*, **270**, 26639, 1995
G.K. Schwartz et al., *J. Natl. Cancer Inst.*, **87**, 1394, 1995

1831 1831-1	D-erythro-Dihydrosphingosine D-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 764-22-7	25 mg 1 g
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C Inhibitor of PLA ₂ and PLD	
	References: D.I. Yule et al., <i>ibid</i> , 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, <i>Adv. in Lipid Res.</i> , 26 , 59, 1993 C. W. Sachs et al., <i>J. Biol. Chem.</i> , 270 , 26639, 1995	
1846	L-erythro-Dihydrosphingosine L-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂	1 mg
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., <i>ibid</i> , 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, <i>Adv. in Lipid Res.</i> , 26 , 59, 1993 C. W. Sachs et al., <i>J. Biol. Chem.</i> , 270 , 26639, 1995	
1851	D-threo-Dihydrosphingosine D-threo-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 6036-86-8	1 mg
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., <i>ibid</i> , 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, <i>Adv. in Lipid Res.</i> , 26 , 59, 1993 C. W. Sachs et al., <i>J. Biol. Chem.</i> , 270 , 26639, 1995	
1324	D,L-erythro-Dihydrosphingosine D,L-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 3102-56-5	25 mg
	Source: synthetic Mol. Wt.: 301 Purity: erythro 77%; threo 23% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C Inhibitor of sphingosine kinase	
	References: D.I. Yule et al., <i>ibid</i> , 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, <i>Adv. in Lipid Res.</i> , 26 , 59, 1993 C. W. Sachs et al., <i>J. Biol. Chem.</i> , 270 , 26639, 1995	
1326	D,L-C16-Dihydrosphingosine (mixed isomers) D,L-Sphinganine with C16 chain C ₁₆ H ₃₅ NO ₂	10 mg
	Source: synthetic Mol. Wt.: 273 Purity: erythro 90%, threo 10% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., <i>ibid</i> , 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, <i>Adv. in Lipid Res.</i> , 26 , 59, 1993 C. W. Sachs et al., <i>J. Biol. Chem.</i> , 270 , 26639, 1995	
1845	D-erythro-C20-Dihydrosphingosine D-erythro-Sphinganine, C20 chain C ₂₀ H ₄₃ NO ₂ CAS#: 24006-62-0	5 mg
	Source: synthetic Mol. Wt.: 330 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1, warm ethanol Storage: -20°C	

- 1839** **D,L-erythro-C20-Dihydrosphingosine** **10 mg**
 D,L-erythro-Sphinganine, C20 chain $C_{20}H_{43}NO_2$
- Source:** synthetic **Mol. Wt.:** 330 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol 5:1, warm ethanol **Storage:** $-20^{\circ}C$

3-keto-Dihydrosphingosines



Catalog number 1876


- 1876** **3-keto-Dihydrosphingosine•HCl** **10 mg**
 3-keto-Sphinganine hydrochloride $C_{18}H_{37}NO_2 \cdot HCl$ **CAS#:** 18944-28-0
- Source:** synthetic **Mol. Wt.:** 336 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

- 1891** **3-keto-C6-Dihydrosphingosine•HCl** **10 mg**
 1-Hydroxy-2-amino-3-keto-hexane • HCl $C_6H_{13}NO_2 \cdot HCl$
- Source:** synthetic **Mol. Wt.:** 168 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** ethanol, methanol, DI water **Storage:** $-20^{\circ}C$

- 1892** **3-keto-C8-Dihydrosphingosine•HCl** **10 mg**
 1-Hydroxy-2-amino-3-keto-octane • HCl $C_8H_{17}NO_2 \cdot HCl$
- Source:** synthetic **Mol. Wt.:** 196 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DI water **Storage:** $-20^{\circ}C$

- 1893** **3-keto-C12-Dihydrosphingosine•HCl** **10 mg**
 1-Hydroxy-2-amino-3-keto-dodecane • HCl $C_{12}H_{25}NO_2 \cdot HCl$
- Source:** synthetic **Mol. Wt.:** 252 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

Phytosphingosines




Catalog number 1330

- 1330** **Phytosphingosine** **50 mg**
1330-1 4-Hydroxysphinganine $C_{18}H_{39}NO_3$ **CAS#** 554-62-1 **1 g**
- Source:** natural, yeast (*Pichia ciferri*) **Mol. Wt.:** 318 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol, chloroform/methanol 2:1 (warm) **Storage:** $-20^{\circ}C$

Reference:

Keliu, Xiping Zhang, Robert L. Lester, and Robert C. Dickson. *J. Biol. Chem.*, **280**, Issue 24, 22679-22687, June 17, 2005

Other sphingosine derivatives and precursors



Catalog number 1896

1320 **N,N-Dimethyl-D-erythro-sphingosine** **5 mg/ml, 1 ml**
C₂₀H₄₁NO₂ CAS#: 119567-63-4

Source: synthetic **Mol. Wt.:** 328 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: isopropanol **Solubility:** chloroform, ethanol, isopropanol, methanol
Storage: -20°C

Inhibitor of phosphokinase C

Reference:

B. Felding-Habermann et al., *Biochemistry*, **29**, 6314, 1990

1896 **N,N-Dihexyl-D-erythro-sphingosine** **5 mg/ml, 1 ml**
Sphingosine with tertiary amine group C₃₀H₆₁NO₂

Source: synthetic **Mol. Wt.:** 468 **Purity:** 95% by TLC **Appearance:** liquid
Solvent: ethanol **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1805 **N-Palmitoyl serinol** **10 mg**
C₁₉H₃₉NO₃ CAS#: 126127-31-9

Source: synthetic **Mol. Wt.:** 329 **Purity:** 98+% by TLC, GC **Appearance:** white solid
Solubility: chloroform, methanol, ethanol **Storage:** -20°C

Sphingosine precursor

Ceramides

Ceramide is a fatty acid amide of sphingosine. It may be formed by dehydrogenation of dihydroceramide; by hydrolysis of sphingomyelin or glycosphingolipids; or by acylation of free sphingosine. Ceramide functions as a precursor in the synthesis of sphingomyelin (by an exchange reaction with phosphatidylcholine and phosphatidylethanolamine); of glycosphingolipids (by glycosylation with UDP-hexose); and of free sphingosine and fatty acid by hydrolysis. The sphingosine can be phosphorylated by a kinase to form sphingosine-1-phosphate, which may undergo further hydrolysis or cleavage.

Control of sphingolipid metabolism maintains vital balance points in cell physiology. Two of ceramide's metabolites, sphingosine-1-phosphate and glucosylceramide, produce cell proliferation. Sphingosine-1-phosphate is also a highly active regulator of angiogenesis, vascular maturation, cardiac development, immunity, and directed cell movement. Sphingosine, the free base, is a potent inhibitor of protein kinase C and is involved in intracellular calcium regulation.

Sphingolipid enzymes seem to be particularly active in cancers, so modifying their activities by exogenous action may provide alternatives to chemical therapies. These enzymes are controlled by many known agents, such as 1,25-dihydroxy-vitamin D₃, tumor necrosis factor- α , nerve growth factor, interleukin 1, endothelial growth factor, glutathione, arachidonic acid, dexamethasone, many anticancer drugs, therapeutic radiation, and activators of the FAS receptor.


Ceramide exerts numerous biological effects, including induction of cell maturation, cell cycle arrest, terminal cell differentiation, cell senescence, and cell death. Other effects include producing reactive oxygen in mitochondria (followed by apoptosis) and stimulating phosphorylation of certain proteins (especially mitogen activated protein). It also stimulates some protein phosphatases (especially protein phosphatase 2A). Thus ceramide is an important controller of protein activity

It is apparent from these relationships that ceramide exists at the crux of several enzyme reaction cycles and that experiments involving sphingolipid function call for control of all of the cycles and their branch-off points. Matreya is the major supplier of these lipids, which can be used as standards for analysis of tissues (a much needed part of modern research) and identification of major sphingolipids.

Ceramides with short side chains have been shown to enter easily into cells where they are biologically active. Ceramides with longer side chains, however, also enter cells if dissolved in dodecane-isopropanol first. Fluorescent labeled ceramides and sphingomyelin made from fluorescent labeled acids instead of plain fatty acids are also available for the study of the localization and metabolism of sphingolipids in the cell. Matreya now offers all four isomers of C₂, C₄, C₆ and C₁₈ ceramides. The corresponding dihydroceramides are being used as inactive controls

In three major reviews, Radin (54-56) has discussed the biochemistry and chemistry of ceramide and outlined many potential approaches to cancer therapy using ceramides and related compounds as generators of apoptosis.

See Literature References on page 96.



Catalog number 1901

Synthetic ceramides derived from C₁₈-sphingosine

1901 **N-Acetyl-D-erythro-sphingosine** **10 mg**
N-C₂:0-D-erythro-Ceramide C₂₀H₃₉NO₃ **CAS#:** 3102-57-6

Source: synthetic **Mol. Wt.:** 342 **Purity:** 98+ by TLC, GC **Appearance:** white solid
Solubility: chloroform, ethanol, methanol, DMSO, (up to 5 mg/ml) **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1829	<p>N-Acetyl-L-threo-sphingosine N-C2:0-L-threo-Ceramide C₂₀H₃₉NO₃</p> <p>Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	1 mg
1847	<p>N-Acetyl-L-erythro-sphingosine N-C2:0-L-erythro-Ceramide C₂₀H₃₉NO₃</p> <p>Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	1 mg
1900	<p>N-Hexanoyl-D-erythro-sphingosine N-C6:0-D-erythro-Ceramide C₂₄H₄₇NO₃ CAS#: 124753-97-5</p> <p>Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO (up to 5 mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	10 mg
1828	<p>N-Hexanoyl-L-threo-sphingosine N-C6:0-L-threo-Ceramide C₂₄H₄₇NO₃</p> <p>Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	1 mg
1848	<p>N-Hexanoyl-L-erythro-sphingosine N-C6:0-L-erythro-Ceramide C₂₄H₄₇NO₃</p> <p>Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC; GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	1 mg
1809	<p>N-Hexanoyl-D-threo-sphingosine N-C6:0-D-threo-Ceramide C₂₄H₄₇NO₃</p> <p>Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol, DMSO (up to 5 mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	1 mg
1903	<p>N-Octanoyl-D-erythro-sphingosine N-C8:0-D-erythro-Ceramide C₂₆H₅₁NO₃ CAS#: 74713-59-0</p> <p>Source: synthetic Mol. Wt.: 426 Purity: 98+ by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml) Storage: -20°C</p> <p>Reference: J. M. L. Hauser et al., J. Biol. Chem. 269, 6803, 1994</p>	10 mg

1830	N-Octanoyl-L-threo-sphingosine N-C8:0-L-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1810	N-Octanoyl-D-threo-sphingosine N-C8:0-D-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1333	N-Decanoyl-D-erythro-sphingosine N-C10:0-D-erythro-Ceramide C ₂₈ H ₅₅ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 454 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, (up to 5mg/ml) Storage: -20°C	
2037	N-Pentadecanoyl-D-erythro-sphingosine N-C15:0-D-erythro-Ceramide C ₃₃ H ₆₅ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 524 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
1915	N-Hexadecanoyl-D-erythro-sphingosine N-C16:0-D-erythro-Ceramide C ₃₄ H ₆₇ NO ₃ CAS#: 24696-26-2	10 mg
	Source: synthetic Mol. Wt.: 538 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
	Reference: Yosuke Osawa, Hiroshi Uchinami, Jacek Bielawski, Robert F. Schwabe, Yusuf A. Hannun, and David A. Brenner. J. Biol. Chem., 280 , Issue 30, 27879-27887, July 29, 2005	
2038	N-Heptadecanoyl-D-erythro-sphingosine N-C17:0-D-erythro-Ceramide C ₃₅ H ₆₉ NO ₃ CAS#: 24696-26-2	10 mg
	Source: synthetic Mol. Wt.: 552 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
1832	N-Octadecanoyl-D-erythro-sphingosine N-C18:0-D-erythro-Ceramide C ₃₆ H ₇₁ NO ₃ CAS#: 2304-81-6	10 mg
	Source: synthetic Mol. Wt.: 566 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, hot ethanol, chloroform/methanol 2:1 (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
2039	N-Nonadecanoyl-D-erythro-sphingosine N-C19:0-D-erythro-Ceramide C ₃₇ H ₇₃ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 580 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	

1843 **N-Octadecanoyl-L-threo-sphingosine** **1 mg**
N-C18:0-L-threo-Ceramide C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1850 **N-Octadecanoyl-L-erythro-sphingosine** **1 mg**
N-C18:0-L-erythro-Ceramide C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1855 **N-Octadecanoyl-D-threo-sphingosine** **1 mg**
N-C18:0-D-threo-Ceramide C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

1916 **N-Tetracosanoyl-D-erythro-sphingosine** **5 mg**
N-C24:0-D-erythro-Ceramide C₄₂H₈₃NO₃ **CAS#:** 34435-05-7

Source: synthetic **Mol. Wt.:** 650 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

2-Hydroxy ceramides

2042 **N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-sphingosine** **5 mg**
N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-ceramide C₃₀H₅₉NO₄

Source: synthetic **Mol. Wt.:** 498 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, methanol, ethanol, DMSO **Storage:** -20°C

2044 **N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-sphingosine** **5 mg**
N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-ceramide C₃₆H₇₁NO₄

Source: synthetic **Mol. Wt.:** 582 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol/water, 2:1:0.5 **Storage:** -20°C

Ceramide made from sphingosines with sphingoid bases other than C18

1842 **N-Acetyl-D-erythro-sphingosine (C14 sphingoid base)** **5 mg**
N-C2:0 Ceramide of D-erythro-C14-sphingosine C₁₆H₃₁NO₂


Source: synthetic **Mol. Wt.:** 285 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO, DMF (up to 5 mg/ml) **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1856 **N-Hexanoyl-D-erythro-sphingosine (C8 sphingoid base)** **1 mg**
N-C6:0 Ceramide of D-erythro-C8-sphingosine C₁₄H₂₇NO₂

Source: synthetic **Mol. Wt.:** 257 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Dihydroceramides



Catalog number 1834

1834 N-Acetyl-D-erythro-dihydrosphingosine 1 mg

N-C2:0-D-erythro-Dihydroceramide; N-Acetyl-D-erythro-sphinganine
 $C_{20}H_{41}NO_3$

Source: synthetic **Mol. Wt.:** 344 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

Reference:

J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1910 N-Hexanoyl-D-erythro-dihydrosphingosine 1 mg

N-C6:0-D-erythro-Dihydroceramide; N-Hexanoyl-D-erythro-sphinganine
 $C_{24}H_{49}NO_3$

Source: synthetic **Mol. Wt.:** 400 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$

Reference:

J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1854 N-Octanoyl-D-erythro-dihydrosphingosine 1 mg

N-C8:0-D-erythro-Dihydroceramide; N-Octanoyl-D-erythro-sphinganine
 $C_{26}H_{53}NO_3$

Source: synthetic **Mol. Wt.:** 428 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO **Storage:** $-20^{\circ}C$

Reference:

J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

2041 N-Octadecanoyl-D-erythro-dihydrosphingosine 10 mg

N-C18:0-D-erythro-Dihydroceramide; N-Octadecanoyl-D-erythro-sphinganine
 $C_{36}H_{73}NO_3$

Source: synthetic **Mol. Wt.:** 568 **Purity:** 98% by TLC **Appearance:** white solid **Solubility:** warm chloroform/methanol, 5:1; hot ethanol, DMSO **Storage:** $-20^{\circ}C$

2-Hydroxy dihydroceramides

2043 N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-dihydrosphingosine 5 mg

N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-dihydroceramide $C_{30}H_{61}NO_4$

Source: synthetic **Mol. Wt.:** 500 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol/water, 2:1:0.5 **Storage:** $-20^{\circ}C$


2045 N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-dihydrosphingosine 5 mg

N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-dihydroceramide $C_{36}H_{73}NO_4$

Source: synthetic **Mol. Wt.:** 584 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol/water, 2:1:0.5 **Storage:** $-20^{\circ}C$

2047 **N-(R,S)-alpha-Hydroxyhexadecanoyl-D-erythro-dihydrosphingosine** **5 mg**
N-(R,S)-alpha-Hydroxy-C16:0-D-erythro-dihydroceramide C₃₄H₆₉NO₄
Source: synthetic **Mol. Wt.:** 556 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/water, 2:1:0.5 **Storage:** -20°C

Ceramides from natural sources



General ceramide structure

1056 **Ceramides** **25 mg**
Ceramides with hydroxy and non-hydroxy acyl groups
C₄₂H₈₃NO₄ **CAS#:** 104404-17-13
Source: natural, bovine **Mol. Wt.:** 666(2-hydroxy-lignoceroyl) **Purity:** 98+% by TLC
Appearance: off-white solid **Solubility:** chloroform/methanol 2:1
Storage: -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1322 **Ceramides** **10 mg**
1322-05 Ceramides with mostly non-hydroxy acyl groups C₃₆H₇₁NO₃ **50 mg**
Source: natural, bovine **Mol. Wt.:** 566 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1, ethanol
Storage: -20°C


Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1323 **Ceramides** **10 mg**
1323-05 Ceramides with mostly hydroxy acyl groups C₃₆H₇₁NO₄ **50 mg**
Source: natural, bovine **Mol. Wt.:** 582 (2-hydroxy-stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1, methanol
Storage: -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

See Table III in Appendix for typical fatty acid content of products prepared from natural sources.


Phytoceramides



Catalog number 1897

- | | | |
|-------------|---|-------------|
| 1897 | N-Acetyl-phytosphingosine
N-C2:0-Phytoceramide C ₂₀ H ₄₁ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 360 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm), warm DMSO Storage: -20°C | |
| 1895 | N-Hexanoyl-phytosphingosine
N-C6:0-Phytoceramide C ₂₄ H ₄₉ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 416 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm) Storage: -20°C | |
| 1894 | N-Octanoyl-phytosphingosine
N-C8:0-Phytoceramide C ₂₆ H ₅₃ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 444 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm) Storage: -20°C | |
| 2035 | N-Hexadecanoyl-phytosphingosine
N-C16:0-Phytoceramide C ₃₄ H ₆₉ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 556 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C | |
| 2034 | N-Stearoyl-phytosphingosine
N-C18:0-Phytoceramide C ₃₆ H ₇₃ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 584 Purity: 98+% by TLC-MS Appearance: white solid Solubility: chloroform/methanol 1:1 (warm) Storage: -20°C | |
| 2036 | N-Tetracosanoyl-phytosphingosine
N-C24:0-Phytoceramide C ₄₂ H ₈₅ NO ₄ | 5 mg |
| | Source: semi-synthetic, yeast (<i>Pichia ciferri</i>) Mol. Wt.: 668 Purity: 98+% by TLC-MS Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C | |

Fluorescent ceramides



Catalog number 1618

Excitation: 460 nm
Emission: 535 nm

- 1841**
1841-001 **N-Hexanoyl-NBD-D-erythro-sphingosine** **100 µg**
N-C6:0-NBD-ceramide; N-C6:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-sphingosine $C_{30}H_{49}N_5O_6$ **CAS#:** 86701-10-2 **1 mg**
- Source:** synthetic **Mol. Wt.:** 575 **Melting Point (°C):** 85.7-87.9 **Purity:** 98+% by TLC **Appearance:** orange film, vacuum dried **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C
- Reference:**
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994
- 1618**
1618-001 **N-Dodecanoyl-NBD-D-erythro-sphingosine** **100 µg**
N-C12:0-NBD ceramide; N-C12:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-sphingosine $C_{36}H_{61}N_5O_6$ **1 mg**
- Source:** synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 2:1, methanol **Storage:** -20°C
- 1857**
1857-001 **N-Hexanoyl-NBD-L-threo-sphingosine** **100 µg**
N-C6:0-NBD-ceramide; N-C6:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminocaproyl)-L-threo-sphingosine $C_{30}H_{49}N_5O_6$ **1 mg**
- Source:** synthetic **Mol. Wt.:** 575 **Purity:** 98+% by TLC **Appearance:** red-orange solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C
- Reference:**
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994
- 1620**
1620-001 **N-Dodecanoyl-NBD-L-threo-sphingosine** **100 µg**
N-C12:0-NBD-ceramide; N-C12:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-sphingosine $C_{36}H_{61}N_5O_6$ **1 mg**
- Source:** synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 2:1, methanol **Storage:** -20°C
- 1624**
1624-001 **N-Hexanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-L-threo-dihydrosphingosine, fluorescent; N-(NBD-aminocaproyl)-L-threo-dihydrosphingosine $C_{30}H_{51}N_5O_6$ **1 mg**
- Source:** synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 2:1, methanol **Storage:** -20°C
- 1623**
1623-001 **N-Dodecanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-L-threo-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-dihydrosphingosine $C_{36}H_{63}N_5O_6$ **1 mg**
- Source:** synthetic **Mol. Wt.:** 662 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 2:1, methanol **Storage:** -20°C


1626 1626-001	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-dihydrosphingosine C ₃₀ H ₅₁ N ₅ O ₆	100 µg 1 mg
	Source: synthetic Mol. Wt.: 578 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1625 1625-001	N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-dihydrosphingosine C ₃₆ H ₆₃ N ₅ O ₆	100 µg 1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1628 1628-001	N-Hexanoyl-NBD-phytosphingosine N-C6:0-NBD-phytoceramide; N-C6:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminocaproyl)-phytosphingosine C ₃₀ H ₅₁ N ₅ O ₇	100 µg 1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 594 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1627 1627-001	N-Dodecanoyl-NBD-phytosphingosine N-C12:0-NBD-phytoceramide; N-C12:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminolauroyl)-phytosphingosine C ₃₆ H ₆₃ N ₅ O ₇	100 µg 1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 678 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information.

Phosphosphingolipids


Sphingomyelins



Catalog number 1051

1051 1051-1	Sphingomyelin SPM; ceramide-1-phosphorylcholine C ₄₁ H ₈₃ N ₂ O ₆ P CAS#: 85187-10-6	25 mg 1 g
	Source: natural, bovine Mol. Wt.: 731 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	Predominately C18:0 and C24:1 fatty acids	


1328	<p>Sphingomyelin SPM; ceramide-1-phosphorylcholine $C_{47}H_{95}N_2O_6P$ CAS#: 85187-10-6</p> <p>Source: natural, porcine Mol. Wt.: 815 (lignoceroyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol Storage: $-20^{\circ}C$</p> <p>Predominately C16:0 and C24:0 fatty acids</p>	25 mg
1329	<p>Sphingomyelin SPM; ceramide-1-phosphorylcholine $C_{46}H_{93}N_2O_6P$ CAS#: 85187-10-6</p> <p>Source: natural, bovine buttermilk Mol. Wt.: 801 (tricosanoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform, ethanol Storage: $-20^{\circ}C$</p> <p>Approximately equal amounts of C16:0, C22:0, C23:0, and C24:0 fatty acids</p>	25 mg
1332 1332-1	<p>Sphingomyelin Ceramide-1-phosphorylcholine $C_{39}H_{79}N_2O_6P$</p> <p>Source: natural, egg yolk, chicken Mol. Wt.: 703 (palmitate) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, methanol, warm ethanol Storage: $-20^{\circ}C$</p>	25 mg 1 gram
1907	<p>N-Acetyl-sphingosylphosphorylcholine Sphingomyelin with C2:0 fatty acid $C_{25}H_{51}N_2O_6P$</p> <p>Source: semi-synthetic, bovine buttermilk Mol. Wt.: 506 Purity: 98+% by TLC Appearance: vacuum dried Solubility: ethanol, chloroform/methanol 2:1 Storage: $-20^{\circ}C$</p> <p>Mixture of D-erythro and L-threo isomers</p>	5 mg
1909	<p>N-Hexanoyl-sphingosylphosphorylcholine Sphingomyelin with C6:0 fatty acid $C_{29}H_{59}N_2O_6P$</p> <p>Source: semi-synthetic, bovine buttermilk Mol. Wt.: 563 Purity: 98+% by TLC Appearance: solid, vacuum dried Solubility: ethanol, chloroform/methanol 2:1 Storage: $-20^{\circ}C$</p> <p>Mixture of D-erythro and L-threo isomers</p>	5 mg
1911	<p>N-Octadecanoyl-sphingosylphosphorylcholine Sphingomyelin with C18:0 fatty acid $C_{41}H_{83}N_2O_6P$</p> <p>Source: semi-synthetic, bovine buttermilk Mol. Wt.: 731 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: $-20^{\circ}C$</p> <p>Mixture of D-erythro and L-threo isomers</p>	5 mg
1890	<p>N-Heptadecanoyl-sphingosylphosphorylcholine Sphingomyelin with C17:0 fatty acid $C_{40}H_{81}N_2O_6P$</p> <p>Source: semi-synthetic, bovine buttermilk Mol. Wt.: 717 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: $-20^{\circ}C$</p> <p>Mixture of D-erythro and L-threo isomers</p>	5 mg
1917	<p>N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine Sphingomyelin with C20:0 fatty acid $C_{43}H_{87}N_2O_6P$</p> <p>Source: semi-synthetic, bovine buttermilk Mol. Wt.: 759 Purity: 98+% by TLC Appearance: solid, vacuum dried Solubility: chloroform/methanol 14:1, ethanol, methanol Storage: $-20^{\circ}C$</p>	0.5 mg



Catalog number 1918

- 1918 N-Docosanoyl-D-erythro-sphingosylphosphorylcholine** **0.5 mg**
 Sphingomyelin with C22:0 fatty acid $C_{45}H_{91}N_2O_6P$
- Source:** semi-synthetic, bovine buttermilk **Mol. Wt.:** 787 **Purity:** 98+% by TLC
Appearance: solid, vacuum dried **Solubility:** chloroform/methanol 14:1, ethanol, methanol **Storage:** $-20^{\circ}C$
- 2200 N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine** **1 mg**
 D-erythro-Sphingomyelin with 1-¹³C-palmitic acid; SPM with ¹³C labeled fatty acid $^{12}C_{38}^{13}CH_{79}N_2O_6P$
- Source:** semi-synthetic, bovine **Mol. Wt.:** 703 **Purity:** 98+% by TLC **Appearance:** waxy solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$
- 1327 N-Acetyl-sphingosylphosphorylethanolamine** **5 mg**
 Sphingosylphosphorylethanolamine with C2:0 fatty acid side chain (D-erythro) $C_{38}H_{77}N_2O_6P$
- Source:** semi-synthetic, bovine buttermilk **Mol. Wt.:** 689 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

Sphingosylphosphorylcholines (SPC)



Catalog number 1318

- 1318 D-erythro-Sphingosylphosphorylcholine** **5 mg**
 D-erythro-SPC $C_{23}H_{49}N_2O_5P$
- Source:** semi-synthetic, bovine buttermilk **Mol. Wt.:** 464 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$
- 1319 L-threo-Sphingosylphosphorylcholine** **5 mg**
 L-threo-SPC $C_{23}H_{49}N_2O_5P$
- Source:** semi-synthetic, bovine buttermilk **Mol. Wt.:** 464 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$
- 1321 Sphingosylphosphorylcholine** **10 mg**
1321-05 lyso-Sphingomyelin; SPC (mixture of D-erythro and L-threo isomers) **50 mg**
 $C_{23}H_{49}N_2O_5P$ **CAS#:** 82970-80-7
- Source:** semi-synthetic, bovine buttermilk **Mol. Wt.:** 466 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

1913 lyso-Dihydrosphingomyelin **1 mg**
Dihydrosphingosylphosphorylcholine (mixture of D-erythro and L-threo isomers) $C_{23}H_{52}N_2O_5P$

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 485 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

Sphingosine phosphates

1803 D-erythro-Sphingosine-1-phosphate **5 mg**
S-1-P, S-P-A $C_{18}H_{38}NO_5P$ **CAS#:** 26993-30-6

Source: synthetic **Mol. Wt.:** 380 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: warm acetic acid **Storage:** $-20^{\circ}C$

References:

T.K. Ghosh et al., J. Biol. Chem., **269**, 22628, 1994

L.Kindman et al., ibid, **269**, 13088, 1994

A. Olivera et al., ibid, **269**, 17924, 1994

M. Mattie, et. al., J. Biol. Chem. **269**:3181, 1994

Yosuke Osawa, Hiroshi Uchinami, Jacek Bielawski, Robert F. Schwabe, Yusuf A. Hannun, and David A. Brenner. J. Biol. Chem., **280**, Issue 30, 27879-27887, July 29, 2005


1852 D-erythro-Dihydrosphingosine-1-phosphate **5 mg**
 $C_{18}H_{40}NO_5P$ **CAS#:** 19794-97-9

Source: synthetic **Mol. Wt.:** 382 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: warm acetic acid **Storage:** $-20^{\circ}C$

2046 N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate, NH_4^+ salt **5 mg**
N-C16:0-Ceramide-1-phosphate $C_{34}H_{68}NO_6P$

Source: synthetic **Mol. Wt.:** 618 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/methanol/acetic acid, 60:15:25 **Storage:** $-20^{\circ}C$

Fluorescent sphingomyelins



Catalog number 1912

Excitation: 460 nm

Emission: 535 nm

1912 N-Hexanoyl-NBD-sphingosylphosphorylcholine **100 μ g**
1912-001 N-C6:0-NBD-sphingomyelin, fluorescent; N-C6:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminocaproyl)-sphingomyelin
 $C_{35}H_{61}N_6O_9P$ **CAS#:** 94885-04-8 **1 mg**

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 740 **Purity:** 98+% by TLC
Appearance: red-brown solid **Solubility:** chloroform, ethanol, methanol
Storage: $-20^{\circ}C$

Mixture of D-erythro and L-threo isomers

1619
1619-001

N-Dodecanoyl-NBD-sphingosylphosphorylcholine

N-C12:0-NBD-sphingomyelin, fluorescent; N-C12:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminolauroyl)-sphingomyelin C₄₁H₇₃N₆O₄P

100 µg
1 mg

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 825 **Purity:** 98+% by TLC

Appearance: orange solid **Solubility:** chloroform/methanol 2:1, methanol

Storage: -20°C

Mixture of D-erythro and L-threo isomers

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information.


Glycosphingolipids

Glycosphingolipids are widely distributed in animal and plant tissues. They consist of a ceramide (Cer) bound in glycosidic linkage through the primary hydroxyl to a mono- or oligosaccharide which may contain substituents such as a sulfate, acetate, or phosphate group. They are amphiphilic and the less glycosylated compounds can be dispersed in buffers by dissolving them in a detergent or organic solvent (EtOH, DMSO, isoPrOH) and mixing by sonication.


Galactosphingolipids, mainly GalCer (cerebrosides) and its sulfate ester, occur in large amounts in the nervous system. Glucosphingolipids, the simplest of which is GlcCer (glucocerebrosides), are very widely distributed, particularly in nerve cell membranes. GlcCer is isolated from a variety of natural sources including human, bovine, and plant. Each of these sources has a heterogeneity in the fatty acid content of the ceramide as well as an occasional variation in the sphingoid chain. Globosides (containing both glucose and galactose) are a prominent group of glycosphingolipids, they contain an α -linked galactose moiety and are typically located in blood cell membranes. Gangliosides are another prominent group of glycosphingolipids; they are acidic because of substitution with sialic (neuraminic) acid. The glycosphingolipids function in a wide range of enzyme and structural interactions, such as immunological or membrane recognition phenomena, binding of microbial pathogens, hormone and growth factor actions, cancer cell growth and malignancy, atherosclerosis, genetic disease errors, blood group determinants, etc. Tissues change in glycosphingolipid composition during embryogenesis, maturation, aging, and other vital physiological processes. Some glycosphingolipids stimulate cell proliferation, others induce apoptosis, effects of great significance to cancer therapy and maturational development. Marked differences in glycosphingolipid composition are seen in normal and cancerous cells. See references 41-53.

See Literature References on page 96.

Galactosylceramides and glucosylceramides



Galactosylceramide



Glucosylceramide

1050 Cerebrosides **50 mg**
Galactosylceramide, ceramide beta-D-galactoside
 $C_{48}H_{93}NO_8$ **CAS#:** 85305-88-0

Source: natural, bovine **Mol. Wt.:** 812 (lignoceryl form) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

Contains both hydroxy and non-hydroxy fatty acid side chains

1066 Cerebroside, Kerasin (top spot) **10 mg**
Galactosylceramide with mostly non-hydroxy fatty acid side chain
 $C_{42}H_{81}NO_8$ **CAS#:** 536-13-0


Source: natural, bovine **Mol. Wt.:** 728 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.5
Storage: $-20^{\circ}C$

1138 Cerebroside, Phrenosin (bottom spot) **10 mg**
Galactosylceramide with mostly 2-hydroxy fatty acid side chains
 $C_{42}H_{81}NO_9$ **CAS#:** 37211-11-3

Source: natural, bovine **Mol. Wt.:** 743 (2-hydroxystearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.5
Storage: $-20^{\circ}C$

1305 Psychosine, (in free amine form) **10 mg**
lyso-Cerebroside; 1-beta-D-galactosylsphingosine
 $C_{24}H_{47}NO_7$ **CAS#:** 2238-90-6

Source: semi-synthetic, bovine **Mol. Wt.:** 461 **Purity:** 98+% by TLC **Appearance:** off-white solid **Solubility:** ethanol, chloroform/methanol 5:1 **Storage:** $-20^{\circ}C$



Catalog number 1325

1325 N-Acetyl-psychosine **10 mg**
N-C2:0-Cerebroside; cerebroside with C2:0 fatty acid $C_{26}H_{49}NO_8$

Source: semi-synthetic, bovine **Mol. Wt.:** 503 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

1335 N-Pentadecanoyl-psychosine **5 mg**
N-C15:0-Cerebroside $C_{39}H_{75}NO_8$

Source: semi-synthetic, bovine **Mol. Wt.:** 685 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/ methanol, 2:1 **Storage:** $-20^{\circ}C$

1334 N-Octanoyl-β-D-galactosylceramide **10 mg**
1334-50 N-C8:0-Galactosylceramide $C_{32}H_{61}NO_8$ **50 mg**

Source: semi-synthetic, bovine **Mol. Wt.:** 588 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/ methanol, 9:1, ethanol, methanol **Storage:** $-20^{\circ}C$

1621 **N-Hexanoyl-NBD-galactosylceramide** **100 µg**
1621-001 N-C6:0-NBD-beta-D-galactosylsphingosine; N-C6:0-NBD-cerebroside; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-galactosylsphingosine C₃₆H₅₉N₅O₁₁ **1 mg**

Source: semi-synthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol, 5:1, methanol **Storage:** -20°C

1057 **Glucocerebrosides** **5 mg**
 Glucosylceramide; ceramide beta-D-glucoside
 C₄₆H₉₃NO₈ **CAS#:** 85305-87-9


Source: natural, human **Mol. Wt.:** 812 (lignoceroyl) **Purity:** 98+% by TLC
Appearance: white to an off white solid **Solubility:** chloroform/methanol 2:1
Storage: -20°C

Contains 24:1 fatty acid side chain. See Table III (p 90-94) for other fatty acid content

1521 **Glucocerebrosides** **5 mg**
1521-50 Glucosylceramide; ceramide beta-D-glucoside C₄₆H₈₉NO₈ **50 mg**

Source: natural, bovine buttermilk **Mol. Wt.:** 784 (docosanoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

See Table III (p 90-94) for side chain variants



Catalog number 1522

1522 **Glucocerebrosides, plant** **5 mg**
1522-100 Glucosylceramide; ceramide beta-D-glucoside C₄₀H₇₅NO₉ **100 mg**

Source: natural, plant **Mol. Wt.:** 714 (2-hydroxyhexadecanoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C


Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4,8) and most of the fatty acids are of the 2-hydroxy type. See Table III page 90-94.

1622 **N-Hexanoyl-NBD-glucosylceramide** **100 µg**
1622-001 N-C6:0-NBD-beta-D-glucosylsphingosine; N-C6:0-NBD-glucosylceramide, fluorescent C₃₆H₅₉N₅O₁₁ **1 mg**

Source: semi-synthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 5:1, methanol **Storage:** -20°C

1306 **Glucosylsphingosine** **5 mg**
 Glucosylsphingosine; lyso-glucocerebroside; 1-beta-D-glucosylsphingosine
 C₂₄H₄₇NO₇ **CAS#:** 52050-17-6

Source: semi-synthetic, bovine **Mol. Wt.:** 461 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** ethanol, methanol, chloroform/methanol 2:1 **Storage:** -20°C




Catalog number 1310

1310 **Glucopsychosine** **5 mg**
 Glucosylsphingosine; lyso-glucocerebroside; 1-beta-D-glucosylsphingosine
 $C_{24}H_{45}NO_7$ **CAS#:** 52050-17-6

Source: natural, plant **Mol. Wt.:** 459 **Purity:** 98+% by TLC
Appearance: off white solid **Solubility:** chloroform/methanol 4:1 **Storage:** $-20^{\circ}C$
 Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4, 8)

1531 **N-Docosanoyl-glucopsychosine** **1 mg**
 Glucocerebroside with C22:0 fatty acid side chain; N-Docosanoyl- β -
 glucosylsphingosine $C_{46}H_{89}NO_8$

Source: semi-synthetic, bovine **Mol. Wt.:** 784 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform **Storage:** $-20^{\circ}C$



Catalog number 1049

1049 **Sulfatides** **50 mg**
 Ceramide-galactoside-3-sulfate; cerebroside sulfate $C_{42}H_{81}NO_{11}S$
CAS#: 85496-63-5

Source: natural, bovine **Mol. Wt.:** 830 (stearoyl) Na Salt **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1 (if needed, a
 few drops of acetic acid) **Storage:** $-20^{\circ}C$

1904 **lyso-Sulfatide (NH_4^+ salt)** **1 mg**
 Sphingosine-1-galactoside-3-sulfate $C_{24}H_{47}NO_{10}S$ **CAS#:** 38621-58-8

Source: semi-synthetic, bovine **Mol. Wt.:** 542 **Purity:** 98+% by TLC **Appearance:**
 white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

2076 **N-Acetyl-sulfatide** **1 mg**
 N-C2:0-sulfatide; N-acetyl-sphingosyl-beta-D-galactoside-3-sulfatide
 $C_{26}H_{49}NO_{11}S$


Source: semi-synthetic, bovine **Mol. Wt.:** 584 **Purity:** 98+% by TLC **Appearance:**
 solid **Solubility:** chloroform/methanol 1:1, ethanol, methanol **Storage:** $-20^{\circ}C$

1875 **N-Palmitoyl-sulfatide** **1 mg**
 Sulfatide with C16:0 fatty acid side chain; N-palmitoyl-sphingosyl-beta-D-
 galactoside-3-sulfate $C_{40}H_{76}NO_{11}S$

Source: semi-synthetic, bovine **Mol. Wt.:** 780 **Purity:** 98+% by TLC **Appearance:**
 white solid **Solubility:** chloroform/methanol 2:1 **Storage:** $-20^{\circ}C$

1888	N-Tetracosanoyl-sulfatide N-C24:0-Sulfatide; N-tetracosanoyl-sphingosyl-beta-D-galactoside-3-sulfate C ₄₈ H ₉₃ NO ₁₁ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 892 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide N-C18:0-D ₃ -Sulfatide C ₄₂ H ₇₈ D ₃ NO ₁₁ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 833 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/DI water 2:1:0.1 Storage: -20°C	
1632 1632-001	N-Dodecanoyl-NBD-sulfatide N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate C ₄₂ H ₇₁ N ₅ O ₁₄ S	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 901 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

Lactosylceramides



1500	Lactosylceramide LC, lactocerebrosides; CDH, ceramide beta-lactoside C ₄₈ H ₉₁ NO ₁₃ CAS#: 4682-48-8	1 mg
	Source: natural, porcine Mol. Wt.: 890 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1, DMSO Storage: -20°C	
	Contains 2-hydroxy fatty acids (See Table I)	
1507 1507-50	Lactosylceramide LC; lactocerebrosides; CDH, ceramide beta-lactoside C ₅₃ H ₁₀₁ NO ₁₃ CAS#: 4682-48-8	5 mg 50 mg
	Source: natural, bovine buttermilk Mol. Wt.: 960 (tricosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
1517	lyso-Lactosylceramide Lactosylsphingosine; lyso-LC C ₃₀ H ₅₇ NO ₁₂	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 623 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
1532	N-Palmitoyl-lactosylceramide Lactosylceramide with C16:0 fatty acid side chain C ₄₆ H ₈₇ NO ₁₃	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 862 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	


1629 **N-Hexanoyl-NBD-lactosylceramide** **50 ug**
1629-001 N-Hexanoyl-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminocaproyl)-lactosylsphingosine C₄₂H₆₉N₅O₁₆ **1 mg**

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 900 **Purity:** 98+% by TLC
Appearance: orange solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

1630 **N-Dodecanoyl-NBD-lactosylceramide** **50 ug**
1630-001 N-Dodecanoyl-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-lactosylsphingosine C₄₈H₈₁N₅O₁₆ **1 mg**

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 984 **Purity:** 98+% by TLC
Appearance: orange solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

Ceramide trihexosides



Catalog number 1067

1067 **Ceramide trihexosides** **1 mg**
1067-10 CTH; Gb3; globotriaosylceramide C₆₀H₁₀₃NO₁₈ **CAS#:** 71965-57-6 **10 mg**

Source: natural, porcine **Mol. Wt.:** 1126 (tetracosanoyl) **Purity:** 98+% by TLC
Appearance: off white solid **Solubility:** chloroform/methanol 2:1, DMSO, hot methanol **Storage:** -20°C

Contains hydroxy and non-hydroxy fatty acid side chains

1513 **Ceramide trihexosides (top spot)** **0.5 mg**
 CTH with non-hydroxy fatty acid side chain C₅₄H₁₀₁NO₁₈

Source: natural, porcine **Mol. Wt.:** 1052 (stearoyl) **Purity:** 98+% by TLC
Appearance: off white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

1514 **Ceramide trihexosides (bottom spot)** **0.5 mg**
 CTH with hydroxy fatty acid side chain C₅₄H₁₀₁NO₁₉

Source: natural, porcine **Mol. Wt.:** 1068 (hydroxystearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 1:1, DMSO, hot methanol **Storage:** -20°C

References:

S. Ashkenazi, T. C. Cleary, J. Clin. Microbiol., **27**, 1145, 1989
 J. Ghislain et al., J. of Immunol., **153**, 3655, 1995

1520 **lyso-Ceramide trihexoside** **1 mg**
 lyso-CTH; lyso-globotriaosylsphingosine C₃₆H₆₇NO₁₇ **CAS#** 126550-86-5


Source: semi-synthetic, porcine **Mol. Wt.:** 786 **Purity:** 98+% by TLC **Appearance:** film, vacuum dried **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

1523 **N-Heptadecanoyl ceramide trihexoside** **0.5 mg**
 N-C17:0-Ceramide trihexoside; N-heptadecanoyl globotriaosylceramide C₅₃H₉₉NO₁₈

Source: semi-synthetic, porcine **Mol. Wt.:** 1038 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol 2:1, DMSO, hot methanol **Storage:** -20°C

1524	N-Tricosanoyl ceramide trihexoside N-C23:0-Ceramide trihexoside; N-tricosanoyl globotriaosylceramide $C_{59}H_{111}NO_{18}$	0.5 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1122 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 2:1, DMSO, hot methanol Storage: -20°C	
1631 1631-001	N-Dodecanoyl-NBD-ceramide trihexoside N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl) ceramide trihexoside $C_{54}H_{91}N_5O_{21}$	100 µg 1 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1145 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1; DMSO; hot methanol Storage: -20°C	
1537	N-Octadecanoyl-D₃-ceramide trihexoside C18:0-D ₃ -CTH; C18:0-D ₃ -Gb3; N-Octadecanoyl-D ₃ -globotriaosylceramide $C_{54}H_{98}D_3NO_{18}$	0.5 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1059 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol 2:1, DMSO Storage: -20°C	

Globosides



1068	Globosides Gb4; globotetrahexosylceramide $C_{62}H_{114}N_2O_{23}$ CAS#: 11034-93-8	5 mg
	Source: natural, porcine Mol. Wt.: 1254 (tetracosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, DMSO, hot methanol Storage: -20°C	

Labeled glycolipids

Stable isotopes

1914	N-Stearoyl-D₃₅-psychosine, perdeuterated Cerebrosides with N-C18:0-D ₃₅ fatty acid side chain $C_{42}H_{46}D_{35}NO_8$	5 mg
	Source: semi-synthetic, bovine Mol. Wt.: 762 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, hot ethanol, chloroform/methanol 2:1 Storage: -20°C	
	Deuterium labeled stearoyl-sidechain	
1533	N-Palmitoyl-D₃-glucopsychosine, deuterated N-C16:0-D ₃ -Glucopsychosine; glucocerebroside with C16:0-D ₃ fatty acid side chain $C_{40}H_{74}D_3NO_8$	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 703 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

1534	N-Palmitoyl-D₃-lactosylceramide, deuterated N-C16:0-D ₃ -Lactosylceramide; lactosylceramide with C16:0-D ₃ fatty acid side chain C ₄₆ H ₈₄ D ₃ NO ₁₃	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 864 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide N-C18:0-D ₃ -Sulfatide C ₄₂ H ₇₈ D ₃ NO ₁₁ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 833 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/DI water 2:1:0.1 Storage: -20°C	
1537	N-Octadecanoyl-D₃-ceramide trihexoside C18:0-D ₃ -CTH; C18:0-D ₃ -Gb3; N-Octadecanoyl-D ₃ -globotriaosylceramide C ₅₄ H ₉₈ D ₃ NO ₁₈	0.5 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1059 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol 2:1, DMSO Storage: -20°C	

Fluorescent compounds

1621 1621-001	N-Hexanoyl-NBD-galactosylceramide N-C6:0-NBD-beta-D-galactosylsphingosine; N-C6:0-NBD-cerebroside; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-galactosylsphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1622 1622-001	N-Hexanoyl-NBD-glucosylceramide N-C6:0-NBD-beta-D-glucosylsphingosine; N-C6:0-NBD-glucosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-glucosylsphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1629 1629-001	N-Hexanoyl-NBD-lactosylceramide N-Hexanoyl-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminocaproyl)-beta-D-lactosylsphingosine C ₄₂ H ₆₉ N ₅ O ₁₆	50 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1630 1630-001	N-Dodecanoyl-NBD-lactosylceramide N-Dodecanoyl-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-beta-D-lactosylsphingosine C ₄₈ H ₈₁ N ₅ O ₁₆	50 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

1631 **N-Dodecanoyl-NBD-ceramide trihexoside** **100 µg**
1631-001 N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl) ceramide trihexoside $C_{54}H_{91}N_5O_{21}$ **1 mg**

Source: semi-synthetic, porcine **Mol. Wt.:** 1145 **Purity:** 98+% by TLC
Appearance: red-orange solid **Solubility:** chloroform/methanol 2:1; DMSO; hot methanol **Storage:** -20°C

1632 **N-Dodecanoyl-NBD-sulfatide** **100 µg**
1632-001 N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate; N-(NBD-amniolauroyl) sulfatide $C_{42}H_{71}N_5O_{14}S$ **1 mg**


Source: semi-synthetic, bovine **Mol. Wt.:** 901 **Purity:** 98+% by TLC
Appearance: red-orange solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical service for more information.

Gangliosides

The diagram below can be used with the general formulas given in the ganglioside descriptions to construct the individual structures.



1064

Gangliotetraosylceramide

Asialo GM₁; Gg4 C₆₂H₁₁₆N₂O₂₃ CAS#: 71012-19-6

1 mg

Source: semi-synthetic, bovine **Mol. Wt.:** 1240 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,4

1512

Gangliotriosylceramide

Asialo-GM₂; Gg3 C₅₆H₁₀₄N₂O₁₈

100 µg

Source: semi-synthetic, human **Mol. Wt.:** 1078 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3

- 1061** **Monosialoganglioside GM₁ (NH₄⁺ salt)** **5 mg**
1061-50 GM₁ C₇₃H₁₃₁N₃O₃₁ CAS#: 37758-47-7 **50 mg**
- Source:** natural, bovine **Mol. Wt.:** 1545 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid, **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C
- General formula: 1,2,3,4,5
- References:**
Qing Zhang, Keiko Furukawa, Ho-Hsiang Chen, Takumi Sakakibara, Takeshi, and Koichi Furukawa. *J. Biol. Chem.*, **281**, Issue 26, 18145-18155, *June 30, 2006*
Birkles. Zeng G, Gaol, Yu R.K., Aubry J. *Biochimie*, **85**:455-63, 2003
Inokuchi J, Momosaki K., Shimeno H., Nagamatsu A., Radin NS. *J. Cell Physiol*, **141**: 573-83, 1989
- 2050** **N-Octadecanoyl-D₃-monosialoganglioside GM₁** **0.5 mg**
N-D₃-Stearoyl-GM₁ C₇₃H₁₂₈N₃O₃₁D₃
- Source:** semi-synthetic, bovine **Mol. Wt.:** 1548 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C
- Reference:**
Qing Zhang, Keiko Furukawa, Ho-Hsiang Chen, Takumi Sakakibara, Takeshi, and Koichi Furukawa. *J. Biol. Chem.*, **281**, Issue 26, 18145-18155, *June 30, 2006*
- 1502** **Monosialoganglioside GM₂ (NH₄⁺ salt)** **500 µg**
GM₂ C₆₇H₁₂₁N₃O₂₆ CAS#: 19600-01-02
- Source:** natural, human **Mol. Wt.:** 1383 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C
- General formula: 1,2,3,5
- 1503** **Monosialoganglioside GM₃ (NH₄⁺ salt)** **500 µg**
GM₃ C₆₄H₁₁₈N₂O₂₁ CAS#: 54827-14-4
- Source:** natural, bovine buttermilk **Mol. Wt.:** 1250 (tricosanoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1, forms micellar solution in water **Storage:** -20°C
- General formula: 1,2,5
- References:**
Inokuchi J, Momosaki K., Shimeno H., Nagamatsu A., Radin NS. *J. Cell Physiol*, **141**: 573-83, 1989
Lovat P.E., Corazzari M., Disano F., Piacentini M., Redfern C.P. *Cancer Lett.*, **228**: 105-110, 2005
Malisan R., Testi R., *IUBMB Life*, **57**:477-482, 2005
- 1535** **Monosialoganglioside GM₄, egg (NH₄⁺ salt)** **0.5 mg**
GM₄ C₅₃H₈₈N₂O₁₆
- Source:** natural, egg, chicken **Mol. Wt.:** 1008 (oleoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol 2:1, forms micellar solution in water **Storage:** -20°C
- General formula: 1,5
- References:**
Ledeen, R.W., R.K. Yu, and L.F. Eng. *J. Neurochem*, **21**:829-839, 1978
Li Y., E. Sugiyama, T. Ariga, J. Nakayama, M. Hayama, Y. Hama, H. Nakagawa, T. Tai, S. Li, and T. Ksama. *J. Lipid Res.*, **43**:1019-1025, 2002

1062	<p>Disialoganglioside GD_{1a} (NH₄⁺ salt) GD_{1a} C₈₄H₁₄₈N₄O₄₀ CAS#: 12707-58-3</p> <p>Source: natural, bovine Mol. Wt.: 1852 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p> <p>General formula: 1,2,3,4,5,7</p>	5 mg
1501	<p>Disialoganglioside GD_{1b} (NH₄⁺ salt) GD_{1b} C₈₄H₁₄₈N₄O₄₀ CAS#: 19553-76-5</p> <p>Source: natural, bovine Mol. Wt.: 1852 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C</p> <p>General formula: 1,2,3,4,5,6</p>	1 mg
1504	<p>Disialoganglioside GD₃ (NH₄⁺ salt) GD₃ C₇₅H₁₂₅N₃O₂₉ CAS#: 62010-37-1</p> <p>Source: natural, bovine buttermilk Mol. Wt.: 1541 (tricosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, forms micellar solution in water Storage: -20°C</p> <p>General formula: 1,2,5,6</p> <p>References: Lovat P.E., Corazzari M., Disano F., Piacentini M., Redfern C.P. <i>Cancer Lett.</i>, 228: 105-110, 2005 Malisan R., Testi R., <i>IUBMB Life</i>, 57:477-482, 2005 Reimer A.B., Forster-Waldl E., Bramswig K.H., Pollak A., Zielinski C.C., Pehamberger H., Lode H.N., Scheiner O., Jensen-Jarolim E., <i>Eur. J. Immunol.</i>, 36:1267-1270, 2006</p>	1 mg
1063	<p>Trisialoganglioside GT_{1b} (NH₄⁺ salt) GT_{1b} C₉₅H₁₆₅N₅O₄₈ CAS#: 59247-13-1</p> <p>Source: natural, bovine Mol. Wt.: 2144 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p> <p>General Formula: 1,2,3,4,5,6,7</p>	5 mg
1516	<p>Tetrasialoganglioside GQ_{1b} (NH₄⁺ salt) GQ_{1b} C₁₀₆H₁₈₂N₆O₅₆ CAS#: 68652-37-9</p> <p>Source: natural, bovine Mol. Wt.: 2435 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p> <p>General formula: 1,2,3,4,5,6,7,8</p> <p>References: Birkles. Zeng G, Gaol, Yu R.K., Aubry J. <i>Biochimie</i>, 85:455-63, 2003 Overell J.R., Willison H.J., <i>Curr. Opin. Neurol.</i>, 18:562-566, 2005</p>	100 µg
1526	<p>Fucosylated monosialoganglioside GM₁ (NH₄⁺ salt) Fucosyl-GM₁ C₇₉H₁₄₁N₃O₃₅</p> <p>Source: natural, porcine Mol. Wt.: 1691 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p>	500 µg

1518	lyso-Monosialoganglioside GM₁ (NH₄⁺ salt) lyso-GM ₁ C ₅₅ H ₉₇ N ₃ O ₃₀ CAS#: 171483-40-2	500 µg
	<p>Source: semi-synthetic, bovine Mol. Wt.: 1279 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.2 Storage: -20°C</p>	
1065	Purified mixed gangliosides, bovine (NH₄⁺ salt) Mixed gangliosides	25 mg
	<p>Source: natural, bovine Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p> <p>Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides</p>	
1525	Purified mixed gangliosides, porcine, (NH₄⁺ salt)	25 mg
	<p>Source: natural, porcine Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C</p> <p>Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides</p>	

Glycosphingolipid reference mixes for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505	Neutral glycosphingolipid qualmix Glycosylceramides, qualitative mix	1 mg/ml, 1 ml
	<p>Source: natural, bovine and porcine Appearance: liquid Solvent: chloroform/methanol 2:1 Solubility: chloroform/methanol 2:1 Storage: -20°C</p> <p>Contains: cerebrosides, lactosylceramide, ceramide trihexoside, globoside</p>	
1508	Monosialoganglioside mix GM ₃ , GM ₂ , GM ₁ qualitative mix	0.5 mg/ml, 1 ml
	<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C</p> <p>Contains: GM₃, GM₂, GM₁</p>	
1509	Disialoganglioside mix GD ₃ , GD _{1a} , GD _{1b} , qualitative mix	0.5 mg/ml, 1 ml
	<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C</p> <p>Contains: GD₃, GD_{1a}, GD_{1b}</p>	
1510	Lactosylceramide and sialosyl derivatives mix LC, GM ₃ , GD ₃ qualitative mix	0.5 mg/ml, 1 ml
	<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C</p> <p>Contains: LC, GM₃, GD₃</p>	

1511 **Gangliotetraosylceramide and sialosyl derivatives mix** **0.5 mg/ml, 1 ml**
asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b} qualitative mix

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water
2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b}

Antibodies directed against glycolipids

These monoclonal and polyclonal antibodies are directed against the carbohydrate chains of Matreya's glycolipids. The same carbohydrate moieties are found on many glycoproteins. The antibodies are for use in ELISA or TLC immunoblotting applications (12). All antibodies are quality tested by actual performance in ELISA and TLC immunoblotting. The antibodies contain no preservatives and are shipped on dry ice.

See Literature References on page 96.

1977 **Anti-ganglioside GD₃** **50 µl**
Monoclonal antibody to GD₃, isotype IgG

Source: natural, mouse hybridoma R-24 cell line **Appearance:** liquid
Solubility: water **Storage:** -20°C **Dry Ice Charge Applies**

Suitable for TLC immunoblotting, ELISA

References:

Kusunoki, A. et al., *Neurology*, **37**:1795 1987
Pukel, C. S. et al., *J. Exptl. Med.*, **155**:1137, 1982
Ren, S. et al., *Cancer Res.*, **49**:7051, 1989

1950 **Anti-ganglioside asialo GM₁** **100 µl**
Polyclonal antibody to asialo-GM₁, isotype IgG

Source: natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to GM₁

References:

Kusunoki, A. et al., *Neurology*, **37**:1795 1987
Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993

1951 **Anti-ganglioside asialo-GM₂** **50 µl**
Polyclonal antibody to asialo-GM₂, isotype IgG, IgM

Source: natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting

References:

Kusunoki, A. et al., *Neurology*, **37**:1795 1987
Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993

1954 **Anti-ganglioside GM₁** **100 µl**
Polyclonal antibody to GM₁, isotype IgG

Source: natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to asialo-GM₁

References:

Kusunoki, A. et al., *Neurology*, **37**:1795 1987
Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993

- 1961** **Anti-ganglioside GM₂ (NANA)** **50 µl**
 Polyclonal antibody to GM₂ (NANA), isotype IgG, IgM
- Source:** natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** –20°C
Dry Ice Charge Applies
- Suitable for ELISA, TLC-immunoblotting
- References:**
 Kusunoki, A. et al., *Neurology*, **37**;1795 1987
 Kusunoki, A. et al. *Arch. Biochem. Biophys.*, 255-226, 1987
 Saito, M. et al. *Biochem. Biophys. Res. Comm.*, **127**:1, 1985
 Pukel, C. S. et al., *J. Exptl. Med.*, **155**:1137, 1982
 Ren, S. et al., *Cancer Res.*, **49**:7051, 1989
 Yu, R. K. et al., *Ann. Neurol.*, **27**:530, 1990
 Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993
- 1962** **Anti-ganglioside GM₂ (NGNA)** **50 µl**
 Polyclonal antibody to GM₂ (NGNA), isotype IgG, IgM
- Source:** natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** –20°C
Dry Ice Charge Applies
- Suitable for TLC immunoblotting, ELISA
- References:**
 Kusunoki, A. et al., *Neurology*, **37**;1795 1987
 Kusunoki, A. et al. *Arch. Biochem. Biophys.*, 255-226, 1987
 Saito, M. et al. *Biochem. Biophys. Res. Comm.*, **127**:1, 1985
 Pukel, C. S. et al., *J. Exptl. Med.*, **155**:1137, 1982
 Ren, S. et al., *Cancer Res.*, **49**:7051, 1989
 Yu, R. K. et al., *Ann. Neurol.*, **27**:530, 1990
- 1957** **Anti-ganglioside GM₄** **50 µl**
 Polyclonal antibody to GM₄, isotype IgG
- Source:** natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** –20°C
Dry Ice Charge Applies
- Suitable for ELISA, TLC-immunoblotting
- References:**
 Kusunoki, A. et al., *Neurology*, **37**;1795 1987
 Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993
- 1960** **Anti-globoside GL-4** **50 µl**
 Polyclonal antibody to GL-4, isotype IgG, IgM
- Source:** natural, rabbit **Appearance:** liquid **Solubility:** water **Storage:** –20°C
Dry Ice Charge Applies
- Suitable for ELISA, TLC-immunoblotting
- References:**
 Kusunoki, A. et al., *Neurology*, **37**;1795 1987
 Kusunoki, A. et al. *Arch. Biochem. Biophys.*, 255-226, 1987
 Saito, M. et al. *Biochem. Biophys. Res. Comm.*, **127**:1, 1985
 Pukel, C. S. et al., *J. Exptl. Med.*, **155**:1137, 1982
 Ren, S. et al., *Cancer Res.*, **49**:7051, 1989
 Yu, R. K. et al., *Ann. Neurol.*, **27**:530, 1990
 Yoship, H. et al., *J. Neurochemistry*, **61**:658, 1993

Enzyme Inhibitors


Ceramide: UDPglucose Transferase. PDMP (D,L-threo-1-phenyl-2-decanoylamino-3-morpholino-1-propanol-HCl) closely resembles the natural sphingolipid substrate of brain glucosyl transferase and is a very potent and competitive inhibitor of the enzyme (13). It has been shown to block outgrowth of neurites in cultured retina and to block glucolipid synthesis in cultured 3T3 cells (14). N.S. Radin and co-workers have shown (15) that PPMP has activity equivalent to that of PDMP when cell homogenates and brain and liver microsomes are used, but it is about 20 times more potent when used with intact cells. In another paper (16), Radin's group has shown that PDMP has substantial activity against Ehrlich ascites tumors in mice. Recent publications from the laboratory of Myles Cabot (17, 18) show that PPMP can reverse multi-drug resistance in cancer cells by causing a build-up of ceramide and preventing the synthesis of glycosylated ceramides. **See Literature References on page 96.**

Matreya also offers the resolved D- and L-threo-isomers of PDMP and PPMP.

Protein Kinase C Inhibitor. Sphingosine is a potent and reversible inhibitor of protein kinase C (19); it also has been shown at low concentrations to stimulate DNA synthesis and act synergistically with known growth factors (20). Note that Safingol (our L-threo-dihydrosphingosine) has also been shown to partially reverse multi-drug resistance in cancer cells (18) *via* inhibition of protein kinase C.

Dihydroceramide desaturase Inhibitor. Cyclopropenylceramide is the first known inhibitor of this enzyme and may allow significant studies on the role of ceramide in apoptosis. Matreya is the only source for this inhibitor. (58)

Ceramidase Inhibitors. N-Oleoylethanolamine has been shown to be an efficacious inhibitor of the ceramidase found in human kidney and cerebellum (21). It is specifically an inhibitor of acid ceramidase (22) with an IC₅₀ of ca. 500 μM. N-Hexadecanoylethanolamine can be used as an inactive control. D-MAPP is a potent (IC₅₀ approximately 5 μM) inhibitor of alkaline ceramidase. Its enantiomer L-MAPP is inactive as an inhibitor and acts as a substrate for this enzyme (22,23). **See Literature References on page 96.**



Catalog number 1751

1751 N-Oleoylethanolamine 100 mg

NOE C₂₀H₃₉NO₂ CAS#: 111-58-0

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, ethyl ether, DMSO **Storage:** -20°C

Activity: acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. *J. Lipid. Res.*, **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. *BBA* **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. *BBRC*, **255**: 456-459, 1999

1786 N-Hexadecanoylethanolamine 100 mg

C₁₈H₃₇NO₂ CAS# 544-31-0

Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: inactive as acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. *J. Lipid. Res.*, **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. *BBA* **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. *BBRC*, **255**: 456-459, 1999

1757 **Anandamide** **10 mg/ml, 1 ml**

Arachidonylethanolamide; 5,8,11,14(Z,Z,Z,Z)-Eicosatetraenoyl 2'-hydroxy-ethyl-amide $C_{22}H_{37}NO_2$ **CAS#:** 94421-68-8

Source: synthetic **Mol. Wt.:** 347 **Purity:** 98+% by TLC **Appearance:** liquid

Solvent: chloroform **Solubility:** chloroform, ethanol **Storage:** $-20^{\circ}C$

Induces apoptosis, endocannabinoid

References:

M. van der Stelt and V. DiMarzo; Prostaglandins Other Lipid Mediat. **77**, 111, 2005

Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. BBA **1657**: 151-163, 2004

C. Grimaldi, et al.; Exp. Cell Res. **312**, 363, 2006

1807 **L-threo-Dihydrosphingosine (Safingol)** **5 mg**
1807-025 L-threo-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ **CAS#:** 15639-50-6 **25 mg**

Source: synthetic **Mol. Wt.:** 301 **Melting Point ($^{\circ}C$):** 103-114 **Purity:** 98+% by


TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol

Storage: $-20^{\circ}C$

References:

C.W. Sachs et al., *ibid.*, **270**, 26639, 1995

G.K. Schwartz et al., J. Natl. Cancer Inst., **87**, 1394, 1995



Catalog number 1859

1859 **D-MAPP** **100 mg**

D-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol $C_{23}H_{39}NO_2$

CAS#: 143492-39-1

Source: synthetic **Mol. Wt.:** 361 **Purity:** 98+% by TLC **Appearance:** white solid

Solubility: ethanol **Storage:** $-20^{\circ}C$

Activity: alkaline ceramidase inhibitor

1860 **L-MAPP** **100 mg**


L-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol $C_{23}H_{39}NO_2$

CAS#: 143492-38-0

Source: synthetic **Mol. Wt.:** 361 **Purity:** 98+% by TLC **Appearance:** white solid

Solubility: ethanol **Storage:** $-20^{\circ}C$

Activity: inactive as alkaline ceramidase inhibitor



Catalog number: 1886

1886
1886-005

N-C8:0-Cyclopropenylceramide

1 mg
5 mg

N-C8:0-CPPC; N-[(1R, 2S)-2-hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] octanamide; GT₁₁ C₂₇H₅₁NO₃

Source: synthetic **Mol. Wt.:** 437 **Melting Point (°C):** 69-70 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** off white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Dihydroceramide desaturase inhibitor

References:

Jacqueline M. Kravcka, Li Li, Zdzislaw M. Szulc, Jacek Bielawski, Besium Ogretmen, Yusuf A. Hannun, Lina M. Obeid, and Alicja Bielawska. *J. Biol. Chem.*, **10**, 1074/jbc. M700647200, *February 5, 2007*
G. Triola, G. Fabrias, and A. Liebaria. *Agnew. Chem. Int. Ed.*, 40, No. 10, 1960-1962, 2001
Triola G., Fabrias G., Casas J., and Liebaria A. *J. Org. Chem.* 68 (26), 9924-9932, 2003
Bedia C., Triola G., Casas J., Liebaria A., Fabrias G. *Or. Biomol. Chem.* 3 (20), 3707-3712, 2005

1887
1887-005

N-C16:0-Cyclopropenylceramide

1 mg
5 mg


N-C16:0-CPPC; N-[(1R, 2S)-2-hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] hexadecamide C₃₅H₆₇NO₃

Source: synthetic **Mol. Wt.:** 550 **Melting Point (°C):** 156-157 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** off white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Dihydroceramide desaturase inhibitor

References:

G. Triola, G. Fabrias, and A. Liebaria. *Agnew. Chem. Int. Ed.*, 40, No. 10, 1960-1962, 2001
Triola G., Fabrias G., Casas J., and Liebaria A. *J. Org. Chem.* 68 (26), 9924-9932, 2003
Bedia C., Triola G., Casas J., Liebaria A., Fabrias G. *Or. Biomol. Chem.* 3 (20), 3707-3712, 2005



Catalog number 1889

1889

Conduritol B epoxide

25 mg

C₆H₁₀O₅ CAS#: 6090-95-5

Source: synthetic **Mol. Wt.:** 162 **Melting Point (°C):** 164-166 **Purity:** 98+% by TLC, NMR **Appearance:** solid **Solubility:** water, DMSO, methanol (slightly) **Storage:** -20°C

Inhibits α-glucosidase activity; specific inhibitor of glucocerebrosidase in cultured cells.

1719 **D,L-threo-PDMP** 100 mg

D,L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
C₂₃H₃₈N₂O₃•HCl CAS#: 80938-69-8

Source: synthetic Mol. Wt.: 427 Melting Point (°C): 158-161 Purity: 98+% by TLC
Appearance: white solid Solubility: ethanol, methanol, chloroform, DMSO
Storage: -20°C


Activity: glucosyl ceramide synthase inhibitor

1720 **D,L-threo-PPMP** 100 mg

D,L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
C₂₉H₅₀N₂O₃•HCl CAS#: 149022-18-4

Source: synthetic Mol. Wt.: 511 Purity: 98+% by TLC Appearance: white solid
Solubility: alcohols, chloroform Storage: -20°C

Activity: glucosyl ceramide synthase inhibitor



1749 **L-threo-PDMP** 10 mg

L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
C₂₃H₃₈N₂O₃•HCl CAS#: 109836-81-9

Source: synthetic Mol. Wt.: 427 Melting Point (°C): 89-92 Purity: 98+% by TLC
Appearance: white solid Solubility: ethanol, methanol Storage: -20°C

1753 **D,L-erythro-PPMP** 100 mg

D,L-erythro-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
C₂₉H₅₀N₂O₃•HCl

Source: synthetic Mol. Wt.: 511 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

1755 **D,L-erythro-PDMP** 100 mg

D,L-erythro-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
C₂₃H₃₈N₂O₃•HCl CAS#: 109760-77-2

Source: synthetic Mol. Wt.: 427 Melting Point (°C): 112-115 Purity: 98+% by TLC
Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO
Storage: -20°C

1756

D-threo-PDMP

10 mg

D-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
 $C_{23}H_{38}N_2O_3 \cdot HCl$ CAS#: 109836-82-0

Source: synthetic **Mol. Wt.:** 427 **Melting Point (°C):** 94-97 **Purity:** 98+% by TLC

Appearance: white solid **Solubility:** ethanol, methanol **Storage:** -20°C

Activity: glucosyl ceramide synthase inhibitor


References:

Nicholson K.M., Quinn D.M., Kellett G. L., Warr J.R. *Br. J. Cancer* **81**: 423-430, 1999

Sietsma H., Veldman R.J., Vander Kolk D., Ausema B., Nijhof W., Kamps W., Vellenga E., Kok J.W. *Clin. Cancer Res.* **6**:942-948, 2000

Basu S., Ma R., Mikulla B., Bradley M., Moulton C., Basu M., Banerjee S., Inokuchi J. J. *Glycoconj.* **20**:157-168, 2003

Radin N.S. *Biochem Pharmacol* **57**:589-595, 1999



Catalog number 1865

1865

D-threo-PPMP

10 mg

D-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
 $C_{29}H_{50}N_2O_2 \cdot HCl$

Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 94-98 **Purity:** 98+% by TLC

Appearance: white solid **Solubility:** ethanol, methanol **Storage:** -20°C

Activity: glucosyl ceramide synthase inhibitor

References:

Abe A., Inokuchi J., Jimbo M., Shimeno H., Nagamatsu A., Shayman J.A., Shukla G.S., Radin N.S. *J. Biochem (Tokyo)* **111**:191-196, 1992

Maurer B.J., Melton L., Billups C., Cabot M.C., Reynolds C.P. *J. Natl. Cancer Inst.* **92**:1897-1909, 2000

Puri A., Hug P., Munoz-Barroso I., Blumenthal R. *Biochem. Biophys. Res. Commun* **242**:219-225, 1998

Couto A.S., Caffaro C., Uhrig M.L., Kimura E., Peres V.J., Merino E.F., Katzin A.M., Nishioka M., Nonami H., Era-Balsells R. *Eur. J. Biochem.* **271**: 2204-2214, 2004

Morjani H., Aouali N., Belhoussine R., Veldman R.J., Levade T., Manfait M. *Int. J. Cancer* **94**:157-165, 2001

1868

L-threo-PPMP

10 mg

L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
 $C_{29}H_{50}N_2O_3 \cdot HCl$

Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 89-94 **Purity:** 98+% by TLC

Appearance: white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1800

Castanospermine


25 mg

1,6,7,8-tetrahydroxyoctahydroindolizine $C_8H_{15}NO_4$ CAS#: 79831-76-8

Source: natural, plant **Mol. Wt.:** 189 **Melting Point (°C):** 210-215 **Purity:** 98+% by TLC, NMR

Appearance: solid **Solubility:** water, methanol/water, 90:10

Storage: -20°C



Catalog number 1717

1717 2-Fluoropalmitic acid 25 mg
 $C_{16}H_{31}FO_2$ CAS#: 89270-22-4

Source: synthetic **Mol. Wt.:** 274 **Melting Point (°C):** 83-85 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Acyl-CoA synthase inhibitor

1718 Methyl 2-fluoropalmitate 10 mg
 $C_{17}H_{33}FO_2$

Source: synthetic **Mol. Wt.:** 288 **Melting Point (°C):** 36-38 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: inactive ester of 2-fluoropalmitic acid

1750 2,2-Difluoropalmitic acid 25 mg
 $C_{16}H_{30}F_2O_2$


Source: synthetic **Mol. Wt.:** 292 **Melting Point (°C):** 50.8-53 **Purity:** 98+% by TLC, GC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C

1858 2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole 1 mg
 THI $C_9H_{14}N_2O_5$ CAS#: 94944-70-4

Source: synthetic **Mol. Wt.:** 230 **Melting Point (°C):** n/a **Purity:** 99% by HPLC, MS, NMR
Appearance: white solid **Solubility:** water **Storage:** -20°C

Reference:

S.R. Schwab, J.P. Pereira, M. Matloubian, Y. Xu, Y. Huang, and J.G. Cyster. *Science* **309**: 1735, 2005



Catalog number 1752

1752 EST 5 mg
 E-64-d; Loxastatin $C_{17}H_{30}N_2O_5$ CAS#: 88321-09-9

Source: synthetic **Mol. Wt.:** 342 **Melting Point (°C):** 125-127 **Purity:** 98+% by TLC, GC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C

Activity: cystein protease inhibitor


Reference:

S. Mehdi, TIBS, **16**, April 1991


Glycerolipids

Glycerophospholipids

Natural phospholipids



Catalog number 1044



Catalog number 1046

1044 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC C₄₄H₈₄NO₈P **CAS#:** 8002-43-5

Source: natural, egg **Mol. Wt.:** 787 (oleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether, ethanol
Storage: -20°C

See Table III page 90-94 for fatty acid content

1070 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC C₄₄H₈₄NO₈P **CAS#:** 8002-43-5

Source: natural, bovine **Mol. Wt.:** 787 (oleoyl) **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform, ethyl ether **Storage:** -20°C

See Table III page 90-94 for fatty acid content

1302 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC C₄₄H₈₀NO₃P **CAS#:** 8002-43-5

Source: natural, plant **Mol. Wt.:** 783 (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C

See Table III page 90-94 for fatty acid content

1046 **lyso-Lecithin** **50 mg**
lyso-Phosphatidylcholine C₂₄H₅₂NO₇P **CAS#:** 9008-30-4

Source: natural, egg **Mol. Wt.:** 496 (palmitoyl) **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol 2:1 **Storage:** -20°C

See Table III page 90-94 for fatty acid content

1047 **Phosphatidylserine** **50 mg/ml, 1 ml**
PS C₄₂H₇₈NO₁₀P

Source: natural, bovine **Mol. Wt.:** 788 (oleoyl) **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform, toluene
Storage: -20°C


See Table III page 90-94 for fatty acid content

- 1048 Phosphatidylinositol (Na⁺ salt)** **10 mg/ml, 1 ml**
 PI C₄₅H₇₈O₁₃P Na CAS# 383907-36-6
Source: natural, plant **Mol. Wt.:** 880 (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C
 See Table III page 90-94 for fatty acid content
- 1336 Phosphatidylinositol, plant, soy, (K⁺ salt)** **50 mg/ml, 1 ml**
 C₄₃H₇₈O₁₃P K CAS# 383907-36-6
Source: natural, plant, soy **Mol. Wt.:** 873 (linoleoyl and pamitoyl) **Purity:** 98+% by TLC
Appearance: tinted liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C
- 1053 Phosphatidic acid (NH₄⁺ salt)** **50 mg**
 PA C₃₉H₇₂O₈P NH₄
Source: semi-synthetic, egg **Mol. Wt.:** 744 (oleoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethyl ether **Storage:** -20°C
 See Table III page 90-94 for fatty acid content
- 1045 Phosphatidylethanolamine** **50 mg/ml, 1 ml**
 PE C₄₁H₇₈NO₈P CAS#: 39382-08-6
Source: natural, egg **Mol. Wt.:** 744 (oleoyl) **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform **Storage:** -20°C
 See Table III page 90-94 for fatty acid content
- 1069 Phosphatidylethanolamine** **50 mg/ml, 1 ml**
 PE C₄₁H₇₈NO₈P CAS#: 90989-93-8
Source: natural, bovine **Mol. Wt.:** 744 (oleoyl) **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform **Storage:** -20°C
- 1301 Phosphatidylethanolamine** **50 mg/ml, 1 ml**
 PE C₄₁H₇₄NO₃P CAS#: 90989-93-8
Source: natural, plant **Mol. Wt.:** 740 (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C
 See Table III page 90-94 for fatty acid content
- 1052 Phosphoglycerides kit** **1 each**
Source: natural, egg, bovine, plant **Purity:** 98+% by TLC **Appearance:** liquid/solid
Solvent: various **Storage:** -20°C
 Individually packed in ampules and vials (Purity 98+%): Phosphatidic acid NH₄⁺ salt 10mg; Phosphatidylethanolamine, egg (in 1 ml CHCl₃) 10mg; Sphingomyelin, bovine 10mg; Phosphatidylserine, bovine (in 1 ml CHCl₃) 10 mg; Lecithin, egg (in 1 ml CHCl₃) 10 mg; lyso-Lecithin, egg 10 mg; Cerebrosides, bovine 10mg; Sulfatides, bovine 10mg; Phosphatidylinositol, Na⁺ salt, plant (in 1 ml CHCl₃) 3mg

Synthetic phospholipids

These phospholipids have 98+% chemical purity except where stated and 99% fatty acid chain purity. Store at -20°C. Solubility: see individual entries


Phosphatidic acid derivatives



Catalog number 1428

- | | | |
|-------------|--|---------------|
| 1428 | 1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid
DMPA C ₃₁ H ₆₀ O ₈ P•Na CAS#: 80724-31-8 | 100 mg |
| | Source: synthetic Mol. Wt.: 615 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C | |
| 1429 | 1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid
DPPA C ₃₅ H ₆₈ O ₈ P•Na CAS#: 70240-64-1 | 100 mg |
| | Source: synthetic Mol. Wt.: 671 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C | |
| 1430 | 1,2-Distearoyl-sn-glycero-3-phosphatidic acid
DSPA C ₃₉ H ₇₆ O ₈ P•Na CAS#: 108321-18-2 | 100 mg |
| | Source: synthetic Mol. Wt.: 727 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C | |

Phosphatidylcholines




Catalog number 1425

- | | | |
|-------------|--|---------------|
| 1442 | 1,2-Dilauroyl-sn-glycero-3-phosphorylcholine
DLPC C ₃₂ H ₆₄ NO ₈ P CAS#: 18194-25-7 | 100 mg |
| | Source: synthetic Mol. Wt.: 622 Purity: 98+% by TLC Appearance: white solid
Solubility: methylene chloride, methanol Storage: -20°C | |
| 1425 | 1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine
DMPC C ₃₆ H ₇₂ NO ₈ P CAS#: 18194-24-6 | 100 mg |
| | Source: synthetic Mol. Wt.: 678 Purity: 98+% by TLC Appearance: white solid
Melting Point: 130-139°C Solubility: methylene chloride, methanol Storage: -20°C | |

1426	1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine DPPC C ₄₀ H ₈₀ NO ₈ P CAS#: 63-89-8	100 mg
	Source: synthetic Mol. Wt.: 734 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1400	1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine DHDPC C ₄₂ H ₈₄ NO ₈ P CAS#: 70897-27-7	50 mg
	Source: synthetic Mol. Wt.: 762 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1427	1,2-Distearoyl-sn-glycero-3-phosphorylcholine DSPC C ₄₄ H ₈₈ NO ₈ P CAS#: 816-94-4	100 mg
	Source: synthetic Mol. Wt.: 790 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1437	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine POPC C ₄₂ H ₈₄ NO ₈ P CAS#: 26853-31-6	100 mg
	Source: synthetic Mol. Wt.: 760 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1445	1-Palmitoyl-sn-glycero-3-phosphorylcholine lyso-PPC C ₂₄ H ₅₀ NO ₇ P CAS#: 17364-16-8	100 mg
	Source: synthetic Mol. Wt.: 496 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	


Phosphatidylglycerols



1443	1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol DLPG C ₃₀ H ₅₈ O ₁₀ P•Na CAS#: 73548-69-3	100 mg
	Source: synthetic Mol. Wt.: 632 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1431	1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol DMPG C ₃₄ H ₆₆ O ₁₀ P•Na CAS#: 67232-80-8	100 mg
	Source: synthetic Mol. Wt.: 689 Purity: 98+% by TLC Appearance: white solid Melting Point: 120-129°C Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1432	1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol DPPG C ₃₈ H ₇₄ O ₁₀ P•Na CAS#: 67232-81-9	100 mg
	Source: synthetic Mol. Wt.: 745 Purity: 98+% by TLC Appearance: white solid Melting Point: 122-127°C Solubility: chloroform/methanol, 5:1 Storage: -20°C	

- 1433** **1,2-Distearoyl-sn-glycero-3-phosphorylglycerol** **100 mg**
 DSPG $C_{42}H_{82}O_{10}P \cdot Na$ **CAS#:** 4537-78-4
- Source:** synthetic **Mol. Wt.:** 801 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/methanol, 5:1 **Storage:** $-20^{\circ}C$
- 1438** **1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol** **100 mg**
 POPG $C_{40}H_{76}O_{10}P \cdot Na$ **CAS#:** 81490-05-3
- Source:** synthetic **Mol. Wt.:** 771 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/methanol, 5:1 **Storage:** $-20^{\circ}C$

Phosphatidylethanolamines




- 1444** **1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine** **100 mg**
 DLPE $C_{29}H_{58}NO_8P$ **CAS#:** 59752-57-7
- Source:** synthetic **Mol. Wt.:** 579 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform + methanol mixture **Storage:** $-20^{\circ}C$
- 1434** **1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine** **100 mg**
 DMPE $C_{33}H_{66}NO_8P$ **CAS#** 998-07-2
- Source:** synthetic **Mol. Wt.:** 636 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/acetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 **Storage:** $-20^{\circ}C$
- 1435** **1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine** **100 mg**
 DPPE $C_{37}H_{74}NO_8P$ **CAS#:** 923-61-5
- Source:** synthetic **Mol. Wt.:** 692 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/acetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 **Storage:** $-20^{\circ}C$
- 1436** **1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine** **100 mg**
 DSPE $C_{41}H_{82}NO_8P$ **CAS#:** 1069-79-0
- Source:** synthetic **Mol. Wt.:** 748 **Purity:** 98+% by TLC **Appearance:** white solid
Solubility: chloroform/acetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 **Storage:** $-20^{\circ}C$
- 1439** **1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000 (Na⁺ salt)** **100 mg**
 DSPE-MPEG-2000 **CAS#:** 147867-65-0
- Source:** synthetic **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform **Storage:** $-20^{\circ}C$

Phosphatidylinositols

The metabolism of inositol lipids is involved in the signal transduction of many hormones, neurotransmitters and growth factors (1,2). In the classical pathway, phosphatidylinositol-specific phospholipase C (PI-PLC) hydrolyzes phosphatidyl 4,5-biphosphate (PIP₂) to yield 1,2-diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃). The role of IP₃ and DAG as second messengers is well recognized.

In a second, more recently discovered pathway, the activation of phosphoinositide (PI) 3-kinase results in the formation of three novel phosphatidyl (PI) lipids phosphorylated at the D3 position of the inositol ring: PI-3-P, PI-3,4-P₂ and PI-3,4,5-P₃ (3). These D3 lipids are not known substrates for any of the phospholipase C enzymes and function as second messengers. PI 3-kinase activity is correlated with many cellular processes, including the regulation of cell growth, oncogenic transformation, chemotaxis and receptor down-regulation among others (4,5,6). The recent paper on the effect of PI3,4-P₂ on the *Akt* proto-oncogene product (9) also contains protocols for applying PIP's to cell cultures. Matreya's synthetic phosphatidylinositols and inositol phosphates are excellent tools for investigating these second messengers, understanding the enzyme mechanisms involved in phosphoinositide metabolism (7,8) and for designing therapeutic pharmacological agents. The compounds are evaluated by ¹H and ³¹P NMR to guarantee enantiomeric purity of >98%. **See Literature References on page 96.**

Phosphatidylinositols



Catalog number 1779

1779 1779-1	Phosphatidylinositol, dipalmitoyl, (NH₄⁺ salt) PI; DPPI (NH ₄ ⁺ salt) C ₄₁ H ₇₈ O ₁₃ P•NH ₄	0.5 mg 1 mg
	Source: synthetic Mol. Wt.: 828 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
1773 1773-1 1773-5	Phosphatidylinositol 3-phosphate, dipalmitoyl, (NH₄⁺ salt) DPPI-3-P; PI-3-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	100 µg 1 mg 5 mg
	Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR, Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C	
	References: B.A. Fenderson, E.M. Eddy, S.Hakomori, <i>BioEssays</i> 12 , 173, 1990 R. T. Dobrowsky et al., <i>ibid</i> , 268 , 15523, 1993 Berridge, M. J., <i>Nature</i> 361 :315, 1993 Bhamare, N. et al., 1996 <i>Phosphorus, Sulfur and Silicon</i> XXX, Overseas Publishers Association, Amsterdam B.V. 109-110:317	
1780 1780-1 1780-5	Phosphatidylinositol 3-phosphate, dipalmitoyl, (Na⁺ salt) DPPI-3-P; PI-3-P dipalmitoyl (Na ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3Na	100 µg 1 mg 5 mg
	Source: synthetic Mol. Wt.: 957 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C	


1922	Phosphatidylinositol 4-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	
1922-1	DOPI-4-P; PI-4-P dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₅ O ₁₆ P ₂ •3NH ₄	1 mg	
1922-5		5 mg	
	Source: synthetic Mol. Wt.: 718 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: soluble in water; slightly soluble in methanol; slightly soluble in chloroform/methanol/DI water, 1:1:0.3 Storage: -20°C		
1919	Phosphatidylinositol 4-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1919-1	DPPI-4-P; PI-4-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	1 mg	
1919-5		5 mg	
	Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: methanol, chloroform/methanol/water 1:1:0.3, slightly soluble in water Storage: -20°C		
1923	Phosphatidylinositol 5-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	0
1923-1	DOPI-5-P; PI-5-P dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₅ O ₁₆ P ₂ •3NH ₄	1 mg	
1923-5		5 mg	
	Source: synthetic Mol. Wt.: 718 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: soluble in water; slightly soluble in methanol; slightly soluble in chloroform/methanol/DI water, 1:1:0.3 Storage: -20°C		
1920	Phosphatidylinositol 5-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1920-1	DPPI-5-P; PI-5-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	1 mg	
1920-5		5 mg	
	Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: methanol, chloroform/methanol/water 1:1:0.3, slightly soluble in water Storage: -20°C		
1781	Phosphatidylinositol bis-3,4-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1781-1	DPPI-3,4-P2; PI-3,4-P2 dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5NH ₄	1 mg	
1781-5		5 mg	
	Source: synthetic Mol. Wt.: 1056 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		
1774	Phosphatidylinositol bis-3,4-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg	
1774-1	DPPI-3,4-P2; PI-3,4-P2 dipalmitoyl (Na ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5Na	1 mg	
1774-5		5 mg	
	Source: synthetic Mol. Wt.: 1081 Purity: 98+% by ¹ H NMR, ³¹ P NMR, Appearance: white solid Solubility: water Storage: -20°C		
	References: Carpenter, C. L. and L.C. Cantley, <i>Curr. Opin. Cell Biol.</i> 8:153, 1996 Ireton, K. et al., <i>Science</i> 274:80, 1996		
1784	Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	
1784-1	DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₉ O ₁₉ P ₃ •5NH ₄	1 mg	
1784-5		5 mg	
	Source: synthetic Mol. Wt.: 831 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		
1778	Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (Na⁺ salt)	100 µg	
1778-1	DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (Na ⁺ salt) C ₂₅ H ₄₄ O ₁₉ P ₃ •5Na	1 mg	
1778-5		5 mg	
	Source: synthetic Mol. Wt.: 856 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C		
1777	Phosphatidylinositol bis-4,5-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1777-1	DPPI-4,5-P2; PI-4,5-P2 dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5NH ₄	1 mg	
1777-5		5 mg	
	Source: synthetic Mol. Wt.: 1056 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		

1782	Phosphatidylinositol bis-4,5-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg
1782-1	DPPI-4,5-P2; PI-4,5-P2 dipalmitoyl (Na ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5Na	1 mg
1782-5		5 mg
	Source: synthetic Mol. Wt.: 1081 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C	
1921	Phosphatidylinositol tris-3,4,5-phosphate, dioctanoyl, (Na⁺ salt)	100 µg
1921-1	DOPI-3,4,5-P3; PI-3,4,5-P3 dioctanoyl (Na ⁺ salt) C ₂₅ H ₄₃ O ₂₂ P ₄ •7Na	1 mg
1921-5		5 mg
	Source: synthetic Mol. Wt.: 980 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C	
1783	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg
1783-1	DPPI-3,4,5-P3; PI-3,4,5-P3 dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₅ O ₂₂ P ₄ •7NH ₄	1 mg
1783-5		5 mg
	Source: synthetic Mol. Wt.: 1170 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C	
1775	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg
1775-1	DPPI-3,4,5-P3; PI-3,4,5-P3, dipalmitoyl (Na ⁺ salt) C ₄₁ H ₇₅ O ₂₂ P ₄ •7Na	1 mg
1775-5		5 mg
	Source: synthetic Mol. Wt.: 1205 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C	
	References: Carpenter, C. L. and L.C. Cantley, Curr. Opin. Cell Biol. 8 :153, 1996 Ireton, K. et al., Science 274 :80, 1996	

Bacterial tetraethers

1303	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>95% pure)	5 mg
	Purified MPL of <i>Thermoplasma acidophilum</i> (>95% pure) C ₉₅ H ₁₈₈ O ₁₆ P	
	Source: natural, Archaeobacteria Mol. Wt.: 1618 Purity: >95% by TLC, HPLC Appearance: yellow solid Solubility: chloroform/methanol 2:1, hexane/2-propanol/DI water 30:40:5 Storage: 4-8°C	
	References: H.-J. Freisleben et al., J. Liposome Res. 3 (3), 817, 1993 H.-J. Freisleben et al., <i>ibid.</i> , 5 (1), 215, 1995 H.-J. Freisleben et al., Chem. Phys. Lipids 78 , 137, 1995 H.-J. Freisleben et al., Archives Biochem. Biophys. 294 (2), 418, 1992	
1303-2	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>50% pure)	50 mg
	MPL of <i>Thermoplasma acidophilum</i> (>50% pure) C ₉₅ H ₁₈₈ O ₁₆ P	
	Source: natural, Archaeobacteria Mol. Wt.: 1618 Purity: >50% by TLC Appearance: brown viscous liquid Solubility: chloroform/methanol 2:1, hexane/2-propanol/DI water 30:40:5 Storage: 4-8°C highly hygroscopic	
	References: H.-J. Freisleben et al., J. Liposome Res. 3 (3), 817, 1993 H.-J. Freisleben et al., <i>ibid.</i> , 5 (1), 215, 1995 H.-J. Freisleben et al., Chem. Phys. Lipids 78 , 137, 1995 H.-J. Freisleben et al., Archives Biochem. Biophys. 294 (2), 418, 1992	

Glycosyl glycerides



Catalog number 1058

- | | | |
|-------------|---|--------------|
| 1058 | Monogalactosyldiglyceride
MGDG (hydrogenated) $C_{45}H_{86}O_{10}$ CAS#: 41670-62-6 | 10 mg |
| | Source: natural, plant Mol. Wt.: 787 (stearoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform/methanol/water 4:1:0.1 Storage: $-20^{\circ}C$ | |
| 1059 | Digalactosyldiglyceride
DGDG (hydrogenated) $C_{51}H_{96}O_{15}$ CAS#: 92457-02-8 | 5 mg |
| | Source: natural, plant Mol. Wt.: 949 (stearoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform/methanol/water 4:1:0.1 Storage: $-20^{\circ}C$ | |

Fatty acids

Simple fatty acids

Saturated fatty acids and methyl esters

These products are 99% pure by GC. They are stable at room temperature and are supplied neat.

- | | | |
|-------------|---|------------|
| 1200 | Methyl hexanoate
Methyl caproate; C6:0 methyl ester $C_7H_{14}O_2$ CAS#: 106-70-7 | 1 g |
| | Source: natural, plant Mol. Wt.: 130 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature | |
| 1196 | Heptanoic acid
C7:0 fatty acid $C_7H_{14}O_2$ CAS#: 111-14-8 | 1 g |
| | Source: natural, plant Mol. Wt.: 130 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature | |
| 1197 | Methyl heptanoate
C7:0 fatty acid methyl ester $C_8H_{16}O_2$ CAS#: 106-73-0 | 1 g |
| | Source: natural, plant Mol. Wt.: 144 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature | |
| 1198 | Octanoic acid
Caprylic acid; C8:0 acid $C_8H_{16}O_2$ CAS#: 124-07-2 | 1 g |
| | Source: natural, plant Mol. Wt.: 144 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature | |

1199	<p>Methyl octanoate Methyl caprylate; C8:0 methyl ester $C_9H_{18}O_2$ CAS#: 111-11-5</p> <p>Source: natural, plant Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1163	<p>Nonanoic acid C9:0 fatty acid; pelargonic acid $C_9H_{18}O_2$ CAS#: 112-05-0</p> <p>Source: synthetic Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg
1164	<p>Methyl nonanoate C9:0 methyl ester $C_{10}H_{20}O_2$ CAS#: 1731-84-6</p> <p>Source: synthetic Mol. Wt.: 172 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg
1261	<p>Methyl decanoate Methyl caprate; C10:0 methyl ester $C_{11}H_{22}O_2$ CAS#: 110-42-9</p> <p>Source: natural, plant Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane Storage: room temperature</p>	500 mg
1165	<p>Undecanoic acid C11:0 fatty acid $C_{11}H_{22}O_2$ CAS#: 112-37-8</p> <p>Source: synthetic Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg
1166	<p>Methyl undecanoate C11:0 methyl ester $C_{12}H_{24}O_2$ CAS#: 1731-86-8</p> <p>Source: synthetic Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg
1008	<p>Dodecanoic acid Lauric acid; C12:0 acid $C_{12}H_{24}O_2$ CAS#: 143-07-7</p> <p>Source: natural, plant Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1009	<p>Methyl dodecanoate Methyl laurate; C12:0 methyl ester $C_{13}H_{26}O_2$ CAS#: 111-82-0</p> <p>Source: natural, plant Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1161	<p>Tridecanoic acid C13:0 fatty acid $C_{13}H_{26}O_2$ CAS#: 638-53-9</p> <p>Source: synthetic Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg
1162	<p>Methyl tridecanoate C13:0 methyl ester $C_{14}H_{28}O_2$ CAS#: 1731-88-0</p> <p>Source: synthetic Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	100 mg

1010	<p>Tetradecanoic acid Myristic acid; C14:0 acid $C_{14}H_{28}O_2$ CAS#: 544-63-8</p> <p>Source: natural, plant Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1011	<p>Methyl tetradecanoate Methyl myristate; C14:0 methyl ester $C_{15}H_{30}O_2$ CAS#: 124-10-7</p> <p>Source: natural, plant Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1012	<p>Pentadecanoic acid C15:0 fatty acid $C_{15}H_{30}O_2$ CAS#: 1002-84-2</p> <p>Source: synthetic Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1013	<p>Methyl pentadecanoate C15:0 methyl ester $C_{16}H_{32}O_2$ CAS#: 7132-64-1</p> <p>Source: synthetic Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1014	<p>Hexadecanoic acid Palmitic acid; C16:0 fatty acid $C_{16}H_{32}O_2$ CAS#: 57-10-3</p> <p>Source: natural, plant Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1015	<p>Methyl hexadecanoate Methyl palmitate; C16:0 methyl ester $C_{17}H_{34}O_2$ CAS#: 112-39-0</p> <p>Source: natural, plant Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1018	<p>Heptadecanoic acid Margaric acid; C17:0 fatty acid $C_{17}H_{34}O_2$ CAS#: 506-12-7</p> <p>Source: synthetic Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1019	<p>Methyl heptadecanoate Methyl margarate; C17:0 methyl ester $C_{18}H_{36}O_2$ CAS#: 1731-92-6</p> <p>Source: synthetic Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1020	<p>Octadecanoic acid Stearic acid; C18:0 fatty acid $C_{18}H_{36}O_2$ CAS#: 57-11-4</p> <p>Source: natural, plant Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g
1021	<p>Methyl octadecanoate Methyl stearate; C18:0 methyl ester $C_{19}H_{38}O_2$ CAS#: 112-61-8</p> <p>Source: natural, plant Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature</p>	1 g

1028	<p>Nonadecanoic acid C19:0 fatty acid C₁₉H₃₈O₂ CAS#: 646-30-0</p> <p>Source: synthetic Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1029	<p>Methyl nonadecanoate C19:0 methyl ester C₂₀H₄₀O₂ CAS#: 1731-94-8</p> <p>Source: synthetic Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1030	<p>Eicosanoic acid Arachidic acid; C20:0 fatty acid C₂₀H₄₀O₂ CAS#: 506-30-9</p> <p>Source: natural, plant Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	500 mg
1031	<p>Methyl eicosanoate Methyl arachidate; C20:0 methyl ester C₂₁H₄₂O₂ CAS#: 1120-28-1</p> <p>Source: natural, plant Mol. Wt.: 326 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	500 mg
1241	<p>Heneicosanoic acid C21:0 fatty acid C₂₁H₄₂O₂ CAS#: 2363-71-5</p> <p>Source: synthetic Mol. Wt.: 326 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1242	<p>Methyl heneicosanoate C21:0 methyl ester C₂₂H₄₄O₂ CAS#: 6064-90-0</p> <p>Source: synthetic Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1035	<p>Docosanoic acid Behenic acid; C22:0 fatty acid C₂₂H₄₄O₂ CAS#: 112-85-6</p> <p>Source: natural, plant Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	500 mg
1036	<p>Methyl docosanoate Methyl behenate; C22:0 methyl ester C₂₃H₄₆O₂ CAS#: 929-77-1</p> <p>Source: natural, plant Mol. Wt.: 354 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	500 mg
1186	<p>Tricosanoic acid C23:0 fatty acid C₂₃H₄₆O₂ CAS#: 2433-96-7</p> <p>Source: synthetic Mol. Wt.: 355 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1187	<p>Methyl tricosanoate C23:0 methyl ester C₂₄H₄₈O₂ CAS#: 2433-97-8</p> <p>Source: synthetic Mol. Wt.: 368 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg

1037	<p>Tetracosanoic acid Lignoceric acid; C24:0 fatty acid $C_{24}H_{48}O_2$ CAS#: 557-59-5</p> <p>Source: synthetic Mol. Wt.: 369 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1038	<p>Methyl tetracosanoate Methyl lignocerate; C24:0 methyl ester $C_{25}H_{50}O_2$ CAS#: 2442-49-1</p> <p>Source: synthetic Mol. Wt.: 382 Purity: 99% by GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	100 mg
1251	<p>Hexacosanoic acid Cerotic acid; C26:0 acid $C_{26}H_{52}O_2$ CAS#: 506-46-7</p> <p>Source: synthetic Mol. Wt.: 370 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	25 mg
1252	<p>Methyl hexacosanoate Methyl cerotate; C26:0 methyl ester $C_{27}H_{54}O_2$ CAS#: 5802-85-4</p> <p>Source: synthetic Mol. Wt.: 411 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature</p>	25 mg

Unsaturated fatty acids and methyl esters

Unsaturated fatty acids are easily oxidized. Flush open containers with argon or nitrogen and store at -20°C, in dark.

1157	<p>Myristoleic acid C14:1 (cis-9) fatty acid $C_{14}H_{26}O_2$ CAS#: 544-64-9</p> <p>Source: natural, plant Mol. Wt.: 226 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1040	<p>Methyl myristoleate C14:1 (cis-9) methyl ester $C_{15}H_{28}O_2$ CAS#: 56219-06-8</p> <p>Source: natural, plant Mol. Wt.: 240 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1243	<p>cis-6-Hexadecenoic acid Sapienic acid $C_{16}H_{30}O_2$</p> <p>Source: synthetic Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethanol, methanol, chloroform, ethyl ether Storage: -20°C</p>	25 mg
1016	<p>Palmitoleic acid C16:1 (cis-9) fatty acid $C_{16}H_{30}O_2$ CAS#: 373-49-9</p> <p>Source: natural, plant Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1017	<p>Methyl palmitoleate C16:1 (cis-9) methyl ester $C_{17}H_{32}O_2$ CAS#: 1120-25-8</p> <p>Source: natural, plant Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1147	<p>Palmitelaidic acid C16:1 (trans-9) acid $C_{16}H_{30}O_2$ CAS#: 10030-73-6</p> <p>Source: synthetic Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg

1148	<p>Methyl palmitelaidate C16:1 (trans-9) methyl ester C₁₇H₃₂O₂ CAS#: 10030-74-7</p> <p>Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1208	<p>11-Hexadecenoic acid, (92% cis, 8% trans) C16:1 (cis-11) acid C₁₆H₃₀O₂</p> <p>Source: synthetic Mol. Wt.: 254 Purity: >98%, by TLC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C</p> <p>92% cis, 8% trans by GC</p>	50 mg
1204	<p>Heptadecenoic acid C17:1 (cis-10) acid C₁₇H₃₂O₂ CAS#: 29743-97-3</p> <p>Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1203	<p>Methyl heptadecenoate C17:1 (cis-10) methyl ester C₁₈H₃₄O₂ CAS#: 75190-82-8</p> <p>Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1022	<p>Oleic acid C18:1 (cis-9) acid C₁₈H₃₄O₂ CAS#: 112-80-1</p> <p>Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	1 g
1023	<p>Methyl oleate C18:1 (cis-9) methyl ester C₁₉H₃₆O₂ CAS#: 112-62-9</p> <p>Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	1 g
1149	<p>Elaidic acid C18:1 (trans-9) acid C₁₈H₃₄O₂ CAS#: 112-79-8</p> <p>Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	1 g
1150	<p>Methyl elaidate C18:1 (trans-9) methyl ester C₁₉H₃₆O₂ CAS#: 1937-62-8</p> <p>Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	1 g
1262	<p>trans 11-Octadecenoic acid C18:1 (trans-11) acid, trans vaccenic acid C₁₈H₃₄O₂ CAS#: 693-72-1</p> <p>Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1263	<p>Methyl trans 11-octadecenoate Methyl trans vaccenate; C18:1 (trans-11) methyl ester C₁₉H₃₆O₂ CAS#: 6198-58-9</p> <p>Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg

1024	<p>Linoleic acid C18:2 (cis,cis-9,12) acid C₁₈H₃₂O₂ CAS#: 60-33-3</p> <p>Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: ethyl ether, ethanol, hexane Storage: -20°C</p>	1 g
1025	<p>Methyl linoleate C18:2 (cis,cis-9,12) methyl ester C₁₉H₃₄O₂ CAS#: 112-63-0</p> <p>Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	1 g
1151	<p>Linoelaidic acid C18:2 (trans, trans-9, 12) acid C₁₈H₃₂O₂ CAS#: 506-21-8</p> <p>Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1152	<p>Methyl linoelaidate C18:2 (trans, trans-9,12) methyl ester C₁₉H₃₄O₂ CAS#: 2566-97-4</p> <p>Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1266	<p>cis-11-Octadecenoic acid cis-vaccenic acid; C18:1(cis-11) acid C₁₈H₃₄O₂ CAS#: 506-17-2</p> <p>Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1267	<p>Methyl cis-11-octadecenoate Methyl cis-vaccenate; C18:1(cis-11) methyl ester C₁₉H₃₆O₂ CAS#: 1937-63-9</p> <p>Source: semi-synthetic, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1026	<p>Linolenic acid C18:3 (all cis-9,12,15) acid C₁₈H₃₀O₂ CAS#: 463-40-1</p> <p>Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	500 mg
1027	<p>Methyl linolenate C18:3 (all cis-9,12,15) methyl ester C₁₉H₃₂O₂ CAS#: 301-00-8</p> <p>Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	500 mg
1153	<p>gamma-Linolenic acid C18:3 (all cis-6,9,12) acid C₁₈H₃₀O₂ CAS#: 506-26-3</p> <p>Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg
1154	<p>Methyl gamma-linolenate C18:3 (all cis-6,9,12) methyl ester C₁₉H₃₂O₂ CAS#: 16326-32-2</p> <p>Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg

1205	Nonadecenoic acid C19:1 (cis-10) acid C ₁₉ H ₃₂ O ₂ CAS#: 73033-09-7	100 mg
	Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1206	Methyl nonadecenoate C19:1 (cis-10) methyl ester C ₂₀ H ₃₄ O ₂ CAS#: 19788-74-0	100 mg
	Source: synthetic Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1032	Eicosenoic acid C20:1 (cis-11) acid C ₂₀ H ₃₈ O ₂ CAS#: 5561-99-9	100 mg
	Source: natural, plant Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1033	Methyl eicosenoate C20:1 (cis-11) methyl ester C ₂₁ H ₄₀ O ₂ CAS#: 2390-09-2	100 mg
	Source: natural, plant Mol. Wt.: 324 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1192	Eicosadienoic acid C20:2 (cis, cis-11, 14) acid C ₂₀ H ₃₆ O ₂ CAS#: 2091-39-6	100 mg
	Source: synthetic Mol. Wt.: 322 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1193	Methyl eicosadienoate C20:2 (cis, cis-11, 14) methyl ester C ₂₁ H ₃₈ O ₂ CAS#: 2463-02-7	100 mg
	Source: synthetic Mol. Wt.: 322 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1179	Methyl 5,8,11-eicosatrienoate C20:3 (all cis-5,8,11) methyl ester, mead acid methyl ester C ₂₁ H ₃₆ O ₂ CAS#: 14602-39-2	1 mg/ml, 1 ml
	Source: natural, plant Mol. Wt.: 320 Purity: 90% by TLC, GC Appearance: liquid Solvent: hexane Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1269	Methyl homogamma linolenate, C20:3n6 Methyl 8,11,14-eicosatrienoate C ₂₁ H ₃₆ O ₂	50 mg
	Source: semi-synthetic, plant Mol. Wt.: 320 Purity: 99% by TLC, GC Appearance: liquid Solubility: hexane, ethyl ether, chloroform Storage: -20°C	
1042	Arachidonic acid C20:4 (all cis-5,8,11,14) acid C ₂₀ H ₃₂ O ₂ CAS#: 506-32-1	100 mg
	Source: natural, plant Mol. Wt.: 304 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride Storage: -20°C Dry Ice Charge Applies	
1034	Methyl arachidonate C20:4 (all cis-5,8,11,14) methyl ester C ₂₁ H ₃₄ O ₂ CAS#: 2566-89-4	100 mg
	Source: natural, plant Mol. Wt.: 318 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	

1167	<p>Eicosapentaenoic acid EPA ; omega-3 fatty acid; C20:5 (all cis-5,8,11,14,17) acid $C_{20}H_{30}O_2$ CAS#: 10417-94-4</p> <p>Source: natural, fish oil Mol. Wt.: 302 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies</p> <p>Anti-hyperlipoproteinemic agent; 5-LOX inhibitor</p>	25 mg
1194	<p>Methyl eicosapentaenoate Methyl ester of omega-3 fatty acid; C20:5 (all cis-5,8,11,14,17) methyl ester $C_{21}H_{32}O_2$ CAS#: 2734-47-6</p> <p>Source: natural, fish oil Mol. Wt.: 316 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies</p>	25 mg
1264	<p>Docosenoic acid C22:1 (cis-13), erucic acid $C_{22}H_{42}O_2$ CAS#: 112-86-7</p> <p>Source: natural, plant Mol. Wt.: 339 Purity: >99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, hexane Storage: -20°C</p>	100 mg
1265	<p>Methyl docosenoate C22:1 (cis-13) methyl ester; methyl erucate $C_{23}H_{44}O_2$ CAS#: 1120-34-9</p> <p>Source: plant Mol. Wt.: 352 Purity: >99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, hexane Storage: -20°C</p>	100 mg
1175	<p>Docosapentaenoic acid C22:5 (all cis-7,10,13,16,19) acid $C_{22}H_{34}O_2$ CAS#: 24880-45-3</p> <p>Source: semi-synthetic Mol. Wt.: 330 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies</p>	25 mg
1244	<p>Methyl docosapentaenoate C22:5 (all cis-7,10,13,16,19) methyl ester $C_{23}H_{36}O_2$ CAS#: 108698-02-8</p> <p>Source: semi-synthetic Mol. Wt.: 344 Purity: 98+% by TLC, GC Appearance: liquid Solubility: ethyl ether, ethanol, hexane, Storage: -20°C Dry Ice Charge Applies</p>	25 mg
1136	<p>Docosahexaenoic acid DHA; C22:6, (all cis-4,7,10,13,16,19) omega-3 fatty acid $C_{22}H_{32}O_2$ CAS#: 6217-54-5</p> <p>Source: natural, plant Mol. Wt.: 328 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies</p>	100 mg
1041	<p>Methyl docosahexaenoate C22:6 (all cis-4,7,10,13,16,19) methyl ester; methyl ester of omega-3 fatty acid $C_{23}H_{34}O_2$ CAS#: 2566-90-7</p> <p>Source: natural, plant Mol. Wt.: 342 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies</p>	100 mg
1155	<p>Nervonic acid C24:1 (cis-15) acid $C_{24}H_{46}O_2$ CAS#: 506-37-6</p> <p>Source: synthetic Mol. Wt.: 367 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C</p>	100 mg

1156 **Methyl nervonate** **100 mg**
C24:1 (cis-15) methyl ester C₂₅H₄₈O₂ **CAS#:** 2733-88-2
Source: synthetic **Mol. Wt.:** 381 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

Trans fatty acids and methyl esters

1147 **Palmitelaidic acid** **100 mg**
C16:1 (trans-9) acid C₁₆H₃₀O₂ **CAS#:** 10030-73-6
Source: synthetic **Mol. Wt.:** 254 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1148 **Methyl palmitelaidate** **100 mg**
C16:1 (trans-9) methyl ester C₁₇H₃₂O₂ **CAS#:** 10030-74-7
Source: synthetic **Mol. Wt.:** 268 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1149 **Elaidic acid** **1 g**
C18:1 (trans-9) acid C₁₈H₃₄O₂ **CAS#:** 112-79-8
Source: synthetic **Mol. Wt.:** 282 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1150 **Methyl elaidate** **1 g**
C18:1 (trans-9) methyl ester C₁₉H₃₆O₂ **CAS#:** 1937-62-8
Source: natural, plant **Mol. Wt.:** 296 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1262 **trans 11-Octadecenoic acid** **100 mg**
C18:1 (trans-11) acid; trans vaccenic acid C₁₈H₃₄O₂ **CAS#:** 693-72-1
Source: synthetic **Mol. Wt.:** 282 **Purity:** 99% by TLC, GC **Appearance:** solid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1263 **Methyl trans 11-octadecenoate** **100 mg**
Methyl trans vaccenate; C18:1 (trans-11) methyl ester C₁₉H₃₆O₂
CAS#: 6198-58-9
Source: synthetic **Mol. Wt.:** 296 **Purity:** 99% by TLC, GC **Appearance:** clear liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1151 **Linoelaidic acid** **100 mg**
C18:2 (trans, trans-9, 12) acid C₁₈H₃₂O₂ **CAS#:** 506-21-8
Source: natural, plant **Mol. Wt.:** 280 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** -20°C

1152 **Methyl linoelaidate** **100 mg**
C18:2 (trans, trans-9,12) methyl ester C₁₉H₃₄O₂ **CAS#:** 2566-97-4
Source: natural, plant **Mol. Wt.:** 294 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** -20°C

1131 **Cis-trans isomer standard** **5 mg/ml, 5 ml**
Qualitative mix

Source: margarine **Appearance:** liquid **Solvent:** 5ml methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Analysis of positional cis-trans fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of cis-trans fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 trans isomers (4 peaks), C18:1 cis & trans isomers (2 peaks), C18:1 cis isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

1181 **9(E),11(E)-Octadecadienoic acid** **25 mg**
9-trans, 11-trans CLA $C_{18}H_{32}O_2$ **CAS#:** 544-71-8



Source: synthetic **Mol. Wt.:** 280 **Melting Point (°C):** 55-57 **Purity:** 98+% by TLC, GC **Appearance:** off-white solid **Solubility:** chloroform, ethanol, hexane, methanol
Storage: -20°C

Conjugated linoleic acid isomers (CLA)

Linoleic acid is an essential fatty acid (18:2 ω6) of which several naturally occurring conjugated derivatives have been identified. These derivatives, called “conjugated linoleic acid” or CLA can have the two double bonds mainly in the 9 and 11 or in the 10 and 12 positions, resulting in eight possible geometric isomers. CLA occurs in meat (24) and dairy products (25,35). In both cases, the 9(Z),11(E)-isomer is predominant and is thought to be the biologically active form. CLA assimilated through the diet of animals is found in the intestinal musosa, liver and adipose tissue (26). See also review article by Parodi (35). CLA has several biological properties. It’s anti-carcinogenic activity has been demonstrated by its ability to inhibit chemically induced tumor formation in animal models of carcinogenesis (24,27,28,29). The addition of CLA to culture medium suppresses the *in vitro* growth of human melanoma, colorectal and breast cancer cells (30). CLA also exhibits anti-atherogenic activity. Addition of CLA to a controlled atherogenic diet significantly reduced the development of atherosclerosis in hamsters and rabbits (31,32). Animals fed a diet containing CLA also had lower levels of low-density-lipoprotein (LDL) cholesterol. CLA may be involved in regulating fat and protein metabolism (33,34). Several species of animals fed CLA-supplemented diets showed improved feed efficiency. Lean body mass increased while body fat was reduced. This seems to be due, mainly or exclusively, to the 10(E),12(Z)-isomer (catalog # 1249, see below). CLA competes with linoleate for Δ6 desaturase (36). Dietary CLA normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat (40) *via* activation of PPAR γ, a result which bears on the possible ameliorization or prevention of NIDDM. The 11(Z),13(E)-isomer (catalog # 1259) has been shown to be concentrated in the heart and in mitochondria. **See Literature References on page 96.**

CLA Research is Being Redone With Our Highly Pure Isomers


Most studies to date have utilized a mixture of CLA isomers containing less than 30% of the presumed active 9(Z),11(E)-isomer (37,38). In addition to the 9,11- and 10,12-isomers, 8,10- and 11,13-isomers have recently been identified in the widely used mixture (38,39). Matreya offers a highly pure CLA which is 98+% the active 9,11-“cis, trans” isomer. The corresponding “trans,trans” and “cis,cis” isomers are also available. In addition, we now offer the pure 10(E),12(Z)-isomer, which has been widely sought for comparison studies. **See Literature References on page 96.**



1245 1245-1 1245-10	9(Z),11(E)-Octadecadienoic acid 9-cis, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	25 mg 1 g 10 g
	Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol, DMSO Storage: -20°C	
1255	Methyl 9(Z), 11(E)-octadecadienoate Methyl ester of CLA (9-cis, 11-trans) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1181	9(E),11(E)-Octadecadienoic acid 9-trans, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-71-8	25 mg
	Source: synthetic Mol. Wt.: 280 Melting Point (°C): 55-57 Purity: 98+% by TLC, GC Appearance: off-white solid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1257	Methyl 9(E),11(E)-octadecadienoate Methyl ester of CLA (9-trans, 11-trans) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	

1248 1248-1	9(Z),11(Z)-Octadecadienoic acid 9-cis, 11-cis CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-40-7	25 mg 1 g
	Source: synthetic Mol. Wt.: 280 Melting Point (°C): 40-42 Purity: 96+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, ethyl ether Storage: -20°C	
1256	Methyl 9(Z), 11(Z)-octadecadienoate Methyl ester of CLA (9-cis, 11-cis) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 96+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1249 1249-1 1249-10	10(E),12(Z)-Octadecadienoic acid 10-trans, 12-cis CLA C ₁₈ H ₃₂ O ₂ CAS#: 2420-44-2	25 mg 1 g 10 g
	Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1254	Methyl 10(E), 12(Z)-octadecadienoate Methyl ester of CLA (10-trans, 12-cis) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1259	11(Z), 13(E)-Octadecadienoic acid 11-cis, 13-trans CLA C ₁₈ H ₃₂ O ₂	25 mg
	Source: synthetic Mol. Wt.: 280 Purity: 77% cis, trans; 2 % cis, cis; 6% trans, trans by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1247-1 1247-10	9(Z),11(E)-Octadecadienoic acid 9-cis, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	1 g 10 g
	Source: synthetic Mol. Wt.: 280 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1%(E),(E) by TLC, GC Appearance: tinted oil Solubility: ethanol, ethyl ether, hexane Storage: -20°C	
1258	Methyl 9(Z),11(E)-octadecadienoate Methyl ester of CLA (9-cis, 11-trans) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1% (E),(E) by TLC, GC Appearance: clear oil Solubility: chloroform, ethyl ether, hexane Storage: -20°C	

Other CLA products and derivatives




Catalog number 1793

- | | | |
|------|---|----------------|
| 1793 | 8-(5-Hexyl-2-furyl)-octanoic acid
Furan fatty acid; 9,12-epoxy-9,11-octadecadienoic acid $C_{18}H_{30}O_3$
CAS#: 4179-44-6 | 25 mg |
| | Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: oil
Solubility: chloroform, ethanol, ethyl ether Storage: $-20^{\circ}C$ | |
| 1409 | 1-Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine
$C_{44}H_{84}NO_8P$ | 25 mg/ml, 1 ml |
| | Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid
Solvent: chloroform Solubility: chloroform, ethanol Storage: $-20^{\circ}C$ | |
| 1410 | 1-Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine
$C_{44}H_{84}NO_8P$ | 25 mg/ml, 1 ml |
| | Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid
Solvent: chloroform Solubility: chloroform, ethanol Storage: $-20^{\circ}C$ | |
| 1411 | 1-Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine
$C_{44}H_{84}NO_8P$ | 25 mg/ml, 1 ml |
| | Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid
Solvent: chloroform Solubility: chloroform, ethanol Storage: $-20^{\circ}C$ | |
| 1794 | Methyl 8-(5-hexyl-2-furyl)-octanoate
Methyl ester of furan fatty acid $C_{19}H_{32}O_3$ CAS#: 10038-16-1 | 25 mg |
| | Source: synthetic Mol. Wt.: 308 Purity: 98+% by TLC, GC Appearance: oil
Solubility: chloroform, ethanol, ethyl ether Storage: $-20^{\circ}C$ | |

Hydroxy fatty acids

2-Hydroxy fatty acids and methyl esters

These products are racemic and 98+% pure by GC and TLC. The 2-hydroxy fatty acids are components of glycosphingolipids and are involved in fatty acid degradation. They are stable and are supplied neat in vials.




Catalog number 1758

1758 1758-1	2-Hydroxydecanoic acid 2-Hydroxy C10:0 acid C ₁₀ H ₂₀ O ₃ CAS#: 5393-81-7	50 mg 1 g
	Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1759 1759-1	Methyl 2-hydroxydecanoate 2-Hydroxy C10:0 methyl ester C ₁₁ H ₂₂ O ₃ CAS#: 71271-24-4	50 mg 1 g
	Source: synthetic Mol. Wt.: 202 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1701 1701-1	2-Hydroxydodecanoic acid 2-Hydroxy C12:0 acid C ₁₂ H ₂₄ O ₃ CAS#: 2984-55-6	50 mg 1 g
	Source: synthetic Mol. Wt.: 216 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1702 1702-1	Methyl 2-hydroxydodecanoate 2-Hydroxy C12:0 methyl ester C ₁₃ H ₂₆ O ₃ CAS#: 51067-85-7	50 mg 1 g
	Source: synthetic Mol. Wt.: 230 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1703 1703-1	2-Hydroxytetradecanoic acid 2-Hydroxy C14:0 acid C ₁₄ H ₂₈ O ₃ CAS#: 2507-55-3	50 mg 1 g
	Source: synthetic Mol. Wt.: 244 Melting Point (°C): 81-82 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1704 1704-1	Methyl 2-hydroxytetradecanoate 2-Hydroxy C14:0 methyl ester C ₁₅ H ₃₀ O ₃ CAS#: 56009-40-6	50 mg 1 g
	Source: synthetic Mol. Wt.: 258 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1705 1705-1	2-Hydroxyhexadecanoic acid 2-Hydroxy C16:0 acid C ₁₆ H ₃₂ O ₃ CAS#: 764-67-0	50 mg 1 g
	Source: synthetic Mol. Wt.: 272 Melting Point (°C): 86-87 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1706 1706-1	Methyl 2-hydroxyhexadecanoate 2-Hydroxy C16:0 methyl ester C ₁₇ H ₃₄ O ₃ CAS#: 16742-51-1	50 mg 1 g
	Source: synthetic Mol. Wt.: 286 Melting Point (°C): 59-60 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1707 1707-1	2-Hydroxyoctadecanoic acid 2-Hydroxy C18:0 acid C ₁₈ H ₃₆ O ₃ CAS#: 629-22-1	50 mg 1 g
	Source: synthetic Mol. Wt.: 300 Melting Point (°C): 92-93 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1708 1708-1	Methyl 2-hydroxyoctadecanoate 2-Hydroxy C18:0 methyl ester C ₁₉ H ₃₈ O ₃ CAS#: 2420-35-1	50 mg 1 g
	Source: synthetic Mol. Wt.: 315 Melting Point (°C): 64-66 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	

1709	2-Hydroxyeicosanoic acid	25 mg
1709-0.5	2-Hydroxy C20:0 acid C ₂₀ H ₄₀ O ₃ CAS#: 16742-48-6	0.5 g
	Source: synthetic Mol. Wt.: 329 Melting Point (°C): 91-92 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1710	Methyl 2-hydroxyeicosanoate	25 mg
1710-0.5	2-Hydroxy C20:0 methyl ester C ₂₁ H ₄₂ O ₃ CAS#: 16742-49-7	0.5 g
	Source: synthetic Mol. Wt.: 343 Melting Point (°C): 62-64 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C	
1711	2-Hydroxydocosanoic acid	25 mg
1711-0.5	2-Hydroxy C22:0 acid C ₂₂ H ₄₄ O ₃ CAS#: 13980-14-8	0.5 g
	Source: synthetic Mol. Wt.: 366 Melting Point (°C): 96-97 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1712	Methyl 2-hydroxydocosanoate	25 mg
1712-0.5	2-Hydroxy C22:0 methyl ester C ₂₃ H ₄₆ O ₃ CAS#: 13980-17-1	0.5 g
	Source: synthetic Mol. Wt.: 371 Melting Point (°C): 72-73 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C	
1713	2-Hydroxytricosanoic acid	10 mg
	2-Hydroxy C23:0 acid C ₂₃ H ₄₆ O ₃ CAS#: 2718-37-8	
	Source: synthetic Mol. Wt.: 371 Melting Point (°C): 98-99 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1714	Methyl 2-hydroxytricosanoate	10 mg
	2-Hydroxy C23:0 methyl ester C ₂₄ H ₄₈ O ₃ CAS#: 118745-41-8	
	Source: synthetic Mol. Wt.: 385 Melting Point (°C): 68-70 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C	
1715	2-Hydroxytetracosanoic acid	5 mg
	2-Hydroxy C24:0 acid; cerebronic acid C ₂₄ H ₄₈ O ₃ CAS#: 544-57-0	
	Source: synthetic Mol. Wt.: 385 Melting Point (°C): 101-104 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1716	Methyl 2-hydroxytetracosanoate	5 mg
	2-Hydroxy C24:0 methyl ester C ₂₅ H ₅₀ O ₃ CAS#: 2433-95-6	
	Source: synthetic Mol. Wt.: 399 Melting Point (°C): 64-65 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C	
1722	2-Hydroxy methyl ester mix	10 mg/ml, 1 ml
	Quantitative mixture	
	Source: synthetic Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
	Contains: 2-OH C14:0, 20%; 2-OH C16:0, 20%; 2-OH C18:0, 15%; 2-OH C20:0, 15.0%; 2-OH C22:0, 10%; 2-OH C23:0, 10%; 2-OH C24:0, 10%	

3-Hydroxy fatty acids and methyl esters

These products are racemic and 98+% pure by GC and TLC. 3-Hydroxy fatty acids occur in the lipid fraction of many microorganisms and are useful in the typing of microbial isolates. They are stable and are supplied neat in vials.



Catalog number 1747

1747 1747-0.5	3-Hydroxyhexanoic acid 3-Hydroxy C6:0 acid C ₆ H ₁₂ O ₃ CAS#: 10191-24-9	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 132 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1748 1748-0.5	Methyl 3-hydroxyhexanoate 3-Hydroxy C6:0 methyl ester C ₇ H ₁₄ O ₃ CAS#: 21188-58-9	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 146 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1745 1745-0.5	3-Hydroxyoctanoic acid 3-Hydroxy C8:0 acid C ₈ H ₁₆ O ₃ CAS#: 88930-08-9	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 160 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1746 1746-0.5	Methyl 3-hydroxyoctanoate 3-Hydroxy C8:0 methyl ester C ₉ H ₁₈ O ₃ CAS#: 85549-54-8	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 174 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1725 1725-0.5	3-Hydroxynonanoic acid 3-Hydroxy C9:0 acid C ₉ H ₁₈ O ₃ CAS#: 88930-09-0	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 174 Melting Point (°C): 60-62 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1726 1726-0.5	Methyl 3-hydroxynonanoate 3-Hydroxy C9:0 methyl ester C ₁₀ H ₂₀ O ₃ CAS#: 83968-06-3	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1727 1727-0.5	3-Hydroxydecanoic acid 3-Hydroxy C10:0 acid C ₁₀ H ₂₀ O ₃ CAS#: 5561-87-5	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 188 Melting Point (°C): 57-60 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1728 1728-0.5	Methyl 3-hydroxydecanoate 3-Hydroxy C10:0 methyl ester C ₁₁ H ₂₂ O ₃ CAS#: 62675-82-5	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 202 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	

1729 1729-0.5	3-Hydroxyundecanoic acid 3-Hydroxy C11:0 acid C ₁₁ H ₂₂ O ₃ CAS#: 40165-88-6	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 202 Melting Point (°C): 74-76 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1730 1730-0.5	Methyl 3-hydroxyundecanoate 3-Hydroxy C11:0 methyl ester C ₁₂ H ₂₄ O ₃ CAS#: 127593-21-9	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 216 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1731 1731-0.5	3-Hydroxydodecanoic acid 3-Hydroxy C12:0 acid C ₁₂ H ₂₄ O ₃ CAS#: 8355-89-3	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 216 Melting Point (°C): 71-72 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	
1732 1732-0.5	Methyl 3-hydroxydodecanoate 3-Hydroxy C12:0 methyl ester C ₁₃ H ₂₆ O ₃ CAS#: 85464-97-7	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 230 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1733 1733-0.5	3-Hydroxytridecanoic acid 3-Hydroxy C13:0 acid C ₁₃ H ₂₆ O ₃ CAS#: 32602-69-0	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 230 Melting Point (°C): 80-83 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1734 1734-0.5	Methyl 3-hydroxytridecanoate 3-Hydroxy C13:0 methyl ester C ₁₄ H ₂₈ O ₃	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 244 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether Storage: -20°C	
1735 1735-0.5	3-Hydroxytetradecanoic acid 3-Hydroxy C14:0 acid C ₁₄ H ₂₈ O ₃ CAS#: 3422-31-9	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 244 Melting Point (°C): 80-81 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1736 1736-0.5	Methyl 3-hydroxytetradecanoate 3-Hydroxy C14:0 methyl ester C ₁₅ H ₃₀ O ₃ CAS#: 55682-83-2	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 258 Melting Point (°C): 36-37 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1739 1739-0.5	3-Hydroxyhexadecanoic acid 3-Hydroxy C16:0 acid C ₁₆ H ₃₂ O ₃ CAS#: 928-17-6	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 272 Melting Point (°C): 85-86 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1740 1740-0.5	Methyl 3-hydroxyhexadecanoate 3-Hydroxy C16:0 methyl ester C ₁₇ H ₃₄ O ₃ CAS#: 51883-36-4	25 mg 0.5 g
	Source: synthetic Mol. Wt.: 286 Melting Point (°C): 43-45 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	

1741 **3-Hydroxyheptadecanoic acid** **25 mg**
1741-0.5 3-Hydroxy C17:0 acid C₁₇H₃₄O₃ **0.5 g**

Source: synthetic **Mol. Wt.:** 286 **Melting Point (°C):** 93-95 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1742 **Methyl 3-hydroxyheptadecanoate** **25 mg**
1742-0.5 3-Hydroxy C17:0 methyl ester C₁₈H₃₆O₃ **0.5 g**

Source: synthetic **Mol. Wt.:** 300 **Melting Point (°C):** 53-55 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C


1743 **3-Hydroxyoctadecanoic acid** **25 mg**
1743-0.5 3-Hydroxy C18:0 acid C₁₈H₃₆O₃ **CAS#:** 45261-96-9 **0.5 g**

Source: synthetic **Mol. Wt.:** 300 **Melting Point (°C):** 52-54 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1744 **Methyl 3-hydroxyoctadecanoate** **25 mg**
1744-0.5 3-Hydroxy C18:0 methyl ester C₁₉H₃₈O₃ **CAS#:** 14531-40-9 **0.5 g**

Source: synthetic **Mol. Wt.:** 315 **Melting Point (°C):** 52-54 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C

Omega hydroxy fatty acids



Catalog number 1754

1754 **Royal Jelly acid** **50 mg**
1754-0.5 10-Hydroxy-2-(E)-decanoic acid; omega-hydroxy C10:1 (2-trans) C₁₀H₁₈O₃ **0.5 g**
CAS#: 14113-05-4

Source: synthetic **Mol. Wt.:** 186 **Melting Point (°C):** 63-65 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1881 **15-Hydroxypentadecanoic acid** **25 mg**
omega-Hydroxy C15:0 C₁₅H₃₀O₃ **CAS#:** 4617-33-8

Source: synthetic **Mol. Wt.:** 258 **Melting Point (°C):** 84-86 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** room temperature

1882 **Methyl 15-hydroxypentadecanoate** **25 mg**
omega-Hydroxy C15:0 fatty acid methyl ester C₁₆H₃₂O₃ **CAS#:** 76529-42-5

Source: synthetic **Mol. Wt.:** 272 **Melting Point (°C):** 50-52 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, warm ethanol, ethyl ether **Storage:** room temperature

1760 **17-Hydroxyheptadecanoic acid** **25 mg**
omega-Hydroxy C17:0 fatty acid C₁₇H₃₄O₃ **CAS#:** 13099-34-8

Source: synthetic **Mol. Wt.:** 286 **Melting Point (°C):** 93-95 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, warm ethanol, ethyl ether **Storage:** room temperature

1761	<p>Methyl 17-hydroxyheptadecanoate omega-Hydroxy C17:0 fatty acid methyl ester $C_{18}H_{36}O_3$ CAS#: 94036-00-7</p> <p>Source: synthetic Mol. Wt.: 300 Melting Point (°C): 59-63 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature</p>	25 mg
1877	<p>20-Hydroxyeicosanoic acid omega-Hydroxy C20:0 fatty acid $C_{20}H_{40}O_3$</p> <p>Source: synthetic Mol. Wt.: 328 Melting Point (°C): 96-98 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature</p>	25 mg
1878	<p>Methyl 20-hydroxyeicosanoate omega-Hydroxy C20:0 fatty acid methyl ester $C_{21}H_{42}O_3$</p> <p>Source: synthetic Mol. Wt.: 342 Melting Point (°C): 69-71 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature</p>	25 mg
1879	<p>21-Hydroxyheneicosanoic acid omega-Hydroxy C21:0 fatty acid $C_{21}H_{42}O_3$</p> <p>Source: synthetic Mol. Wt.: 342 Melting Point (°C): 72-75 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature</p>	25 mg
1880	<p>Methyl 21-hydroxyheneicosanoate omega-Hydroxy C21:0 fatty acid methyl ester $C_{22}H_{44}O_3$</p> <p>Source: synthetic Mol. Wt.: 356 Melting Point (°C): 73-76 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature</p>	25 mg
1818	<p>22-Hydroxydocosanoic acid Phellonic acid; omega-hydroxy C22:0 fatty acid $C_{22}H_{44}O_3$</p> <p>Source: synthetic Mol. Wt.: 356 Melting Point (°C): 100-102 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature</p>	25 mg
1819	<p>Methyl 22-hydroxydocosanoate omega-Hydroxy C22:0 fatty acid methyl ester $C_{23}H_{46}O_3$</p> <p>Source: synthetic Mol. Wt.: 370 Melting Point (°C): 73-75 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature</p>	25 mg
1883	<p>Methyl 27-hydroxyheptacosanoate omega-Hydroxy C27:0 fatty acid methyl ester $C_{28}H_{56}O_3$</p> <p>Source: synthetic Mol. Wt.: 440 Melting Point (°C): 85-89 Purity: 97+% by TLC, GC Appearance: white solid Solubility: chloroform Storage: room temperature</p>	25 mg
1884	<p>Methyl 30-hydroxytriacontanoate omega-Hydroxy C30:0 fatty acid methyl ester $C_{31}H_{62}O_3$</p> <p>Source: synthetic Mol. Wt.: 482 Melting Point (°C): 88-91 Purity: 97+% by TLC, GC Appearance: white solid Solubility: chloroform Storage: room temperature</p>	25 mg

Other hydroxy fatty acids

1815 **Methyl threo-2,3-dihydroxypalmitate** **10 mg**
2,3-Dihydroxy C16:0 fatty acid methyl ester $C_{17}H_{34}O_4$

Source: synthetic **Mol. Wt.:** 302 **Melting Point (°C):** 77-79 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** room temperature

1182 **Ricinelaidic acid** **100 mg**
12-Hydroxy C18:1 (9-trans) fatty acid $C_{18}H_{34}O_3$ **CAS#:** 82188-83-8



Source: synthetic **Mol. Wt.:** 298 **Melting Point (°C):** 50-53 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1183 **Methyl ricinelaidate** **100 mg**
12-Hydroxy C18:1 (9-trans) methyl ester $C_{19}H_{36}O_3$ **CAS#:** 7706-01-6

Source: synthetic **Mol. Wt.:** 312 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1766 **6-Hydroxyoctadecanoic acid** **10 mg**
6-Hydroxy C18:0 fatty acid $C_{18}H_{36}O_3$

Source: synthetic **Mol. Wt.:** 300 **Melting Point (°C):** 80-82 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol **Storage:** room temperature



1898 **Methyl D, L-threo-corynomycolate** **25 mg**
Hydroxy fatty acid with long branched chain $C_{33}H_{66}O_3$


Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 70 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform **Storage:** room temperature

1899 **Methyl D, L-erythro-corynomycolate** **25 mg**
Hydroxy fatty acid with long branched chain $C_{33}H_{66}O_3$

Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 58 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform **Storage:** room temperature

Branched and cyclic fatty acids

iso-Fatty acids and esters



Catalog number 1656

- | | | |
|-------------|---|--------------|
| 1656 | Methyl 11-methyldodecanoate
iso-Tridecanoic methyl ester; iso C13 methyl ester $C_{14}H_{28}O_2$
CAS#: 5129-57-7

Source: synthetic Mol. Wt.: 228 Purity: 98+% by GC Appearance: liquid
Solubility: hexane, ethyl ether, methylene chloride Storage: $-20^{\circ}C$ | 20 mg |
| 1657 | Methyl 12-methyltridecanoate
iso-Tetradecanoic methyl ester; iso C14 methyl ester $C_{15}H_{30}O_2$
CAS#: 5129-58-8

Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: liquid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1605 | 13-Methyltetradecanoic acid
iso-Pentadecanoic acid; iso C15 acid $C_{15}H_{30}O_2$

Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: white solid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1600 | Methyl 13-methyltetradecanoate
iso-Pentadecanoic methyl ester; iso C15 methyl ester $C_{16}H_{32}O_2$
CAS#: 5129-59-9

Source: synthetic Mol. Wt.: 256 Purity: 98+% by GC Appearance: clear liquid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1601 | Methyl 14-methylpentadecanoate
iso-Palmitic methyl ester; iso C16 methyl ester $C_{17}H_{34}O_2$ CAS#: 5129-60-2


Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: clear liquid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1606 | 15-Methylhexadecanoic acid
iso-Heptadecanoic acid; iso C17 acid $C_{17}H_{34}O_2$

Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: white solid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1602 | Methyl 15-methylhexadecanoate
iso-Heptadecanoic methyl ester; iso C17 methyl ester $C_{18}H_{36}O_2$
CAS#: 6929-04-0

Source: synthetic Mol. Wt.: 284 Purity: 98+% by GC Appearance: clear liquid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |
| 1603 | Methyl 17-methyloctadecanoate
iso-Nonadecanoic methyl ester; iso C19 methyl ester $C_{20}H_{40}O_2$

Source: synthetic Mol. Wt.: 312 Purity: 98+% by GC Appearance: clear liquid
Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$ | 20 mg |

anteiso-Fatty acids and esters



1615 **12-Methyltetradecanoic acid** **20 mg**
anteiso-Pentadecanoic acid; anteiso C15 acid $C_{15}H_{30}O_2$ **CAS#:** 5502-94-3

Source: synthetic **Mol. Wt.:** 242 **Purity:** 98+% by GC **Appearance:** white solid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1612 **Methyl 12-methyltetradecanoate** **20 mg**
anteiso-Pentadecanoic methyl ester; anteiso C15 methyl ester $C_{16}H_{32}O_2$
CAS#: 5129-66-8

Source: synthetic **Mol. Wt.:** 256 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1613 **Methyl 13-methylpentadecanoate** **20 mg**
anteiso-Palmitic methyl ester; anteiso C16 methyl ester $C_{17}H_{34}O_2$
CAS#: 5487-50-3

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1616 **14-Methylhexadecanoic acid** **20 mg**
anteiso-Heptadecanoic acid; anteiso C17 acid $C_{17}H_{34}O_2$ **CAS#:** 5918-29-6

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by GC **Appearance:** white solid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1614 **Methyl 14-methylhexadecanoate** **20 mg**
anteiso-Heptadecanoic methyl ester; anteiso C17 methyl ester $C_{18}H_{36}O_2$
CAS#: 2490-49-5

Source: synthetic **Mol. Wt.:** 284 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

Methylated fatty acids

1207 **D,L-2,6-Dimethylheptanoic acid** **50 mg**
2,6-Dimethyl C7:0 fatty acid $C_9H_{18}O_2$

Source: synthetic **Mol. Wt.:** 158 **Purity:** 98+% by TLC, GC **Appearance:** clear oil
Solubility: chloroform **Storage:** room temperature

1791 **10-Methylhexadecanoic acid** **25 mg**
10-Methyl C16:0 fatty acid $C_{17}H_{34}O_2$

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by TLC, GC **Appearance:** clear oil
Solubility: chloroform **Storage:** room temperature

1792 **Methyl 10-methylhexadecanoate** **25 mg**
10-Methyl C16:0 fatty acid methyl ester $C_{18}H_{36}O_2$

Source: synthetic **Mol. Wt.:** 284 **Purity:** 98+% by TLC, GC **Appearance:** clear oil
Solubility: chloroform **Storage:** room temperature

1195 **Phytanic acid** **25 mg**
3,7,11,15-Tetramethylhexadecanoic acid $C_{20}H_{40}O_2$ **CAS#:** 14721-66-5

Source: semi-synthetic **Mol. Wt.:** 312 **Purity:** 97+% by GC **Appearance:** white solid **Solubility:** chloroform, methanol **Storage:** -20°C

Cyclopropyl fatty acids and esters

1822 **Dihydrosterculic acid** **25 mg**
cis-9,10-Methyleneoctadecanoic acid $C_{19}H_{36}O_2$ **CAS#:** 4675-61-0

Source: synthetic **Mol. Wt.:** 296 **Melting Point (°C):** 38-42 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, hexane **Storage:** -20°C

1823 **Methyl dihydrosterculate** **25 mg**
Methyl cis-9,10-methyleneoctadecanoate $C_{20}H_{38}O_2$ **CAS#:** 3971-54-8

Source: synthetic **Mol. Wt.:** 310 **Purity:** 98+% by TLC, GC **Appearance:** oil **Solubility:** chloroform, ethanol, methanol, hexane **Storage:** -20°C

Unusual fatty acids and derivatives

1751 **N-Oleoylethanolamine** **100 mg**
NOE $C_{20}H_{39}NO_2$ **CAS#:** 111-58-0

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, ethyl ether, DMSO **Storage:** -20°C

Activity: acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. *J. Lipid. Res.*, **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. *BBA* **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. *BBRC*, **255**: 456-459, 1999

1786 **N-Hexadecanoylethanolamine** **100 mg**
 $C_{18}H_{37}NO_2$ **CAS#** 544-31-0

Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, **Storage:** -20°C

Activity: inactive as acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. *J. Lipid. Res.*, **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. *BBA* **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. *BBRC*, **255**: 456-459, 1999

1757 **Anandamide** **10 mg/ml, 1 ml**

Arachidonylethanolamide; 5,8,11,14(Z,Z,Z,Z)-eicosatetraenoyl 2-hydroxyethyl-amide $C_{22}H_{37}NO_2$ **CAS#:** 94421-68-8

Source: synthetic **Mol. Wt.:** 347 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform, ethanol **Storage:** $-20^{\circ}C$

Induces apoptosis, endocannabinoid


References:

Wasilewski M., Wieckowski M.R., Dymmowska D, Wojtczak L. *BBA* **1657**: 151-163, 2004
M. van der Stelt and V. Di Marzo; *Prostaglandins Other Lipid Mediat.* **77**, 2005
C. Grimaldi, et al.; *Exp. Cell Res.* **312**, 363, 2006

Other lipids

Tocopherols

Catalog number 1072




1072 **rac-alpha-Tocopherol** **50 mg/ml, 1 ml**

5,7,8-Trimethyltocol $C_{29}H_{50}O_2$ **CAS#:** 59-02-9

Source: synthetic **Mol. Wt.:** 431 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** $-20^{\circ}C$

Catalog number 1071




1071 **rac-beta-Tocopherol** **50 mg/ml, 1 ml**

5,8-Dimethyltocol $C_{28}H_{48}O_2$ **CAS#:** 148-03-8

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** $-20^{\circ}C$

Catalog number 1073




1073 **rac-gamma-Tocopherol** **50 mg/ml, 1 ml**

7,8-Dimethyltocol $C_{28}H_{48}O_2$ **CAS#:** 73980-80-0

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 97% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** $-20^{\circ}C$

Catalog number 1074




1074

rac-5,7-Dimethyltocol
C₂₈H₄₈O₂ CAS#: 493-35-6

50 mg/ml, 1 ml

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, ethyl ether, chloroform, alcohols **Storage:** -20°C

Catalog number 1790




1790

(+)-delta-Tocopherol
8-Methyltocol C₂₇H₄₆O₂ CAS#: 119-13-1

50 mg/ml, 1 ml

Source: natural, plant **Mol. Wt.:** 403 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C

Catalog number 1797




1797

Tocol
rac-Tocol C₂₆H₄₄O₂

50 mg/ml, 1 ml

Source: synthetic **Mol. Wt.:** 389 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, methanol, ethanol
Storage: -20°C

Cholestane derivatives



Catalog number 1006

1006

Cholesterol
C₂₇H₄₆O CAS#: 57-88-5

500 mg

Source: natural, ovine **Mol. Wt.:** 386 **Melting Point (°C):** 147-148 **Purity:** 98+%
by GC **Appearance:** white solid **Solubility:** chloroform, ethanol **Storage:** -20°C

1115 **5-alpha-Cholestane** **100 mg**
C₂₇H₄₈O CAS#: 481-21-0

Source: synthetic **Mol. Wt.:** 373 **Purity:** 98+% by GC **Appearance:** white solid
Solubility: chloroform, ethyl ether, hexane **Storage:** -20°C

1116 **Coprostanol** **25 mg**
5-beta-Cholestane-3-beta-ol C₂₇H₄₈O CAS#: 360-68-9

Source: semi-synthetic **Mol. Wt.:** 390 **Melting Point (°C):** 101-103 **Purity:** 98+%
by GC **Appearance:** white solid **Solubility:** chloroform, ethyl ether, warm methanol
Storage: -20°C

Plant sterols and steryl glucosides

1119 **Plant sterol mix** **25 mg/ml, 1 ml**
Sterol mixture, qualitative

Source: natural, plant **Appearance:** liquid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains: Brassicasterol (13%), campesterol (26%), stigmasterol (7%), β-sitosterol (53%) in order of elution

1123 **Plant sterols kit** **1 kit**
Sterols kit

Source: synthetic or plant **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

Contains in individual packages: steryl glucoside 25 mg, esterified steryl glucoside 10 mg, plant sterol mixture 25 mg, β-sitosterol (55%) 100 mg, desmosterol (85%) 2 mg, lanosterol (55%) 100 mg, stigmasterol 25 mg, ergosterol 25 mg, coprostanol 5 mg, cholestanol 100 mg

1113 **β-Sitostanol** **50 mg**
Stigmastanol C₂₉H₅₂O CAS#: 19466-47-8

Source: synthetic **Mol. Wt.:** 417 **Melting Point (°C):** 127-132 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1120 **Lanosterol** **500 mg**
C₃₀H₅₀O CAS#: 79-63-0

Source: synthetic or plant **Mol. Wt.:** 427 **Purity:** 55% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1121 **Stigmasterol** **100 mg**
5,22-cholestadien-24-beta-ethyl-3-beta-ol C₂₉H₄₈O CAS#: 83-48-7

Source: synthetic **Mol. Wt.:** 413 **Melting Point (°C):** 170 **Purity:** 95% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1122 **Ergosterol** **100 mg**
C₂₈H₄₄O CAS#: 57-87-4

Source: synthetic or plant **Mol. Wt.:** 397 **Melting Point (°C):** 156 **Purity:** 95% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1117 Steryl glucoside **25 mg**
C35H60O6

Source: natural, plant **Mol. Wt.:** 576 **Melting Point (°C):** 283-287 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1
Storage: -20°C

1118 Esterified steryl glucoside **10 mg**
1:1:1, sterol:glucose:fatty acid C51H90O7

Source: natural, plant **Mol. Wt.:** 814 **Purity:** 98+% by TLC **Appearance:** solid film
Solubility: chloroform, ethyl ether, pyridine **Storage:** -20°C

Sterol, glucose and fatty acid in a molar ratio 1:1:1. Mol. Wt. based on β -sitosterol glucoside palmitate.

Propyleneglycol Monoesters

1862 2-Hydroxypropyl hexadecanoate **100 mg**
Propyleneglycol monopalmitate C19H38O3

Source: synthetic **Mol. Wt.:** 314 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** hexane, ethyl ether, alcohol, chloroform, **Storage:** -20°C

1863 2-Hydroxypropyl octadecanoate **100 mg**
Propyleneglycol monostearate C21H42O3

Source: synthetic **Mol. Wt.:** 342 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** hexane, ethyl ether, alcohol, chloroform, **Storage:** -20°C

Standards and reference compounds

Food industry mixes

Each methyl ester mix is carefully prepared by weight.

4210 KEL-FIM-FAME-5 mix **15.5 mg/ml 1 ml**
Methyl ester mix

Source: synthetic or plant **Appearance:** liquid **Solvent:** heptane **Solubility:** heptane
Storage: -20°C

Contains the methyl esters of the following fatty acids (mg/ml in brackets): C8:0 (0.3), C10:0 (0.5), C12:0 (1.0), C13:0 (0.5), C14:0 (0.5), C14:1 (0.3), C15:0 (0.3), C16:0 (2.0), C16:1 (1.0), C17:0 (0.5), C18:0 (1.0), C18:1tr (0.4), C18:1c (3.0), C18:2 (2.0), C20:0 (0.3), C18:3 (1.0), C20:1 (0.3), C22:0 (0.3), C22:1 (0.3), listed in order of their elution.

2009 FIM-FAME-6 mix **33 mg/ml, 1 ml**
Methyl ester mix

Source: synthetic or plant **Appearance:** liquid **Solvent:** heptane **Solubility:** heptane
Storage: -20°C

Contains the methyl esters of these fatty acids. Each methyl ester is 3.03% of the mixture except C16:0 which is 6.06%. C4:0, C6:0, C8:0, C10:0, C11:0, C12:0, C13:0, C14:0, C14:1(cis-9), C15:0, C15:1(cis-10), C16:0, C16:1(cis-9), C17:0, C17:1(cis-10), C18:0, C18:1(trans-9), C18:1(cis-9), C18:2(all-cis-9,12), C20:0, C18:3(all-cis 6,9,12), C20:1(cis-11), C18:3(all-cis 9,12,15), C20:2(all-cis 11,14), C22:0, C20:3(all-cis 8,11,14), C22:1(cis 13), C20:3(all-cis 11,14,17), C20:4(all-cis 5,8,11,14), C22:2(all-cis 13,16), C24:1(cis-15), C22:6(all-cis 4,7,10,13,16,19), listed in order of their elution.

1125 Alditol acetate mix-2 **50 mg/ml, 1 ml**
Quantitative carbohydrate mix

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform
Storage: -20°C

Contains: mannitol, galactitol, glucitol and inositol hexaacetates, 12.5 mg/ml each

Other fatty acid methyl ester mixes

1722 2-Hydroxy methyl ester mix **10 mg/ml, 1 ml**

Source: synthetic **Appearance:** liquid **Solvent:** methylene chloride **Solubility:** methylene chloride **Storage:** -20°C

Quantitative mix contains: C14:0, 20%; C16:0, 20%; C18:0, 15%; C20:0, 15%; C22:0, 10%; C23:0, 10%; C24:0, 10%

1131 Cis-trans isomer standard **5 mg/ml, 5 ml**

Source: margarine **Appearance:** liquid **Solvent:** 5ml methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Analysis of positional cis-trans fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of cis-trans fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 trans isomers (4 peaks), C18:1 cis & trans isomers (2 peaks), C18:1 cis isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

AOCS animal and vegetable oil reference mixes (RM mixes)

By studying problems with the quantitative analysis of animal and vegetable oils and fats, the American Oil Chemists' Society has found certain mixtures to be useful as reference standards. The composition of each mixture (see Table I below) is similar to the fatty acid distribution of certain oils. All mixtures are in methyl ester form and ready for GC analysis

Table I. AOCS Oil Reference Mixes

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated in the Table.

Mix No. Catalog No.	RM-1 1084	RM-2 1085	RM-3 1086	Rapeseed 1083	RM-4 1087	RM-5 1088	RM-6 1089
C8:0 Caprylate						7.0	
C10:0 Caprate						5.0	
C12:0 Laurate						48.0	
C14:0 Myristate			1.0	1.0		15.0	2.0
C16:0 Palmitate	6.0	7.0	4.0	4.0	11.0	7.0	30.0
C16:1 Palmitoleate							3.0
C18:0 Stearate	3.0	5.0	3.0	3.0	3.0	3.0	14.0
C18:1 Oleate	35.0	18.0	45.0	60.0	80.0	12.0	41.0
C18:2 Linoleate	50.0	36.0	15.0	12.0	6.0	3.0	7.0
C18:3 Linolenate	3.0	34.0	3.0	5.0			3.0
C20:0 Arachidate	3.0		3.0	3.0			
C20:1 Eicosenoate				1.0			
C22:0 Behenate			3.0	3.0			
C22:1 Erucate			20.0	5.0			
C24:0 Lignocerate			3.0	3.0			

1083 Rapeseed oil reference mixture

25 mg/ml, 1 ml

AOCS rapeseed oil reference mix

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: ethyl ether, methylene chloride **Storage:** -20°C

Suitable standard for low erucic acid oil

1084 RM-1 mix

50 mg

AOCS reference mix RM-1


Source: synthetic or plant **Appearance:** clear oil **Solubility:** chloroform, ethyl ether
Storage: -20°C

Suitable standard for corn, cottonseed, soybean, safflower, sesame, poppy seed, walnut kapok, and rice oils

1085	<p>RM-2 mix AOCS reference mix RM-2</p> <p>Source: synthetic or plant Appearance: clear oil Solubility: chloroform, ethanol, ethyl ether Storage: -20°C</p> <p>Suitable standard for linseed, perilla, hempseed, and rubberseed oils</p>	50 mg
1086	<p>RM-3 mix AOCS reference mix RM-3</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: ethyl ether, methylene chloride Storage: -20°C</p> <p>Suitable standards for peanut, rapeseed, and mustard seed oils</p>	50 mg/ml, 1 ml
1087	<p>RM-4 mix AOCS reference mix RM-4</p> <p>Source: synthetic or plant Appearance: clear oil Solubility: chloroform, ethyl ether Storage: -20°C</p> <p>Suitable standard for olive, teaseed, and neatsfoot oils</p>	50 mg
1088	<p>RM-5 mix AOCS reference mix RM-5</p> <p>Source: synthetic or plant Appearance: clear oil Solubility: chloroform Storage: -20°C</p> <p>Suitable standard for coconut, palm kernel, babassu and ouri-ouri oils</p>	50 mg
1089	<p>RM-6 mix AOCS reference mix RM-6</p> <p>Source: synthetic or plant Appearance: clear oil Solubility: ethyl ether, methylene chloride Storage: -20°C</p> <p>Suitable standard for lard, beef tallow, mutton tallow, and palm oil</p>	50 mg
1082	<p>RM-7 kit AOCS reference mix RM-7 kit</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C</p> <p>50 mg ampules of RM-1, RM-2, RM-3, RM-4, RM-5, RM-6, and 25 mg of Rapeseed oil reference mixture</p>	50 mg ampules

Custom mixes

Custom fatty acid methyl ester mixes can be prepared to your specification. Minimum quantity requirements apply to these orders.



Cat# 4210 spiked with 0.4 mg/ml C18:2t ester (methyl linoleidate) and chromatographed on a Supelco SP 2330 fused silica column.

Peak number	FAME
1	C8:0
2	C10:0
3	C12:0
4	C13:0
5	C14:0
6	C14:1
7	C15:0
8	C16:0
9	C16:1
10	C17:0
11	C18:0
12	C18:1t-9
13	C18:1c-9
14	C18:2t,t-9,12
15	C18:2c,c-9,12
16	C20:0
17	C18:3
18	C20:1
19	C22:0
20	C22:1

Table II. Standards for GC analysis**GLC Standard mixes**

GLC-10 through GLC-100 standards are **equal weight measures** of fatty acid methyl esters. They are quantitative standards, useful for determining relative **retention times and response factors**.

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated below. All double bonds are cis.

Mixture Number Catalog Number	GLC-10 1095	GLC-30 1097	GLC-40 1098	GLC-50 1099	GLC-60 1100	GLC-70 1101	GLC-80 1102	GLC-90 1103	GLC-100 1104
C8:0 Caprylate		20.0				20.0			
C9:0 Nonanoate						20.0			
C10:0 Caprate		20.0				20.0			
C11:0 Undecanoate						20.0			
C12:0 Laurate		20.0				20.0			
C13:0 Tridecanoate							20.0	20.0	
C14:0 Myristate		20.0					20.0		
C15:0 Pentadecanoate							20.0	20.0	
C16:0 Palmitate	20.0	20.0	25.0				20.0		
C16:1 Palmitoleate-9 cis				20.0					
C17:0 Heptadecanoate							20.0	20.0	
C18:0 Stearate	20.0		25.0						20.0
C18:1 Oleate-9 cis	20.0			20.0					
C18:2 Linoleate-9,12 all cis	20.0								
C18:3 Linolenate-9,12,15 all cis	20.0								
C19:0 Nonadecanoate								20.0	20.0
C20:0 Arachidate			25.0		25.0				20.0
C20:1 Eicosenoate-11 cis				20.0	25.0				
C20:2 Eicosadienoate-11,14 all cis					25.0				
C20:3 Eicosatrienoate-11,14,17 all cis					25.0				
C21:0 Heneicosanoate								20.0	20.0
C22:0 Behenate			25.0						20.0
C22:1 Erucate-13				20.0					
C24:1 Nervonate				20.0					

1095 **GLC-10 mix** **50 mg**
Quantitative GC mix

Source: synthetic or plant **Appearance:** clear oil **Solubility:** methylene chloride
Storage: -20°C

1097 **GLC-30 mix** **50 mg**
Quantitative GC mix

Source: synthetic or plant **Appearance:** clear oil **Solubility:** methylene chloride
Storage: -20°C

1098	<p>GLC-40 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml
1099	<p>GLC-50 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml
1100	<p>GLC-60 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml
1101	<p>GLC-70 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: clear oil Solubility: methylene chloride Storage: –20°C</p>	50 mg
1102	<p>GLC-80 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml
1103	<p>GLC-90 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml
1104	<p>GLC-100 mix Quantitative GC mix</p> <p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: –20°C</p>	50 mg/ml, 1 ml

Water soluble fatty acid mixes

1106	<p>WSFA-2 mix Water soluble fatty acid qualitative mix</p> <p>Appearance: liquid Solvent: water Solubility: water Storage: Room Temp</p> <p>Contains: acetic, propionic, isobutyric, n-butyric, isovaleric and n-valeric acids</p>	5 ml
1108	<p>WSFA-4 mix Water soluble fatty acid qualitative mix</p> <p>Appearance: liquid Solvent: water Solubility: water Storage: Room Temp</p> <p>Contains: acetic, propionic, isobutyric, n-butyric, 2-methyl butyric, isovaleric and n-valeric acids</p>	5 ml

Microbiology standard mixes

1105 **GLC-110 mix** **10 mg/ml, 1 ml**
Bacterial lipid standard, qualitative mix

Source: various **Appearance:** liquid **Solvent:** chloroform **Solubility:** methylene chloride, chloroform **Storage:** -20°C

Contains:

methyl 12-methyltridecanoate	(iso-C14:0)	methyl 14-methylpentadecanoate	(iso-C16:0)
methyl myristate	(C14:0)	methyl palmitate	(C16:0)
methyl 12-methyltetradecanoate	(anteiso-C15:0)	methyl 14-methylhexadecanoate	(anteiso-C17:0)
methyl pentadecanoate	(C15:0)		

1114 **Bacterial acid methyl esters CP mix** **10 mg/ml, 1 ml**
Qualitative mix

Source: various **Appearance:** liquid **Solvent:** methyl caproate **Solubility:** hexane, ethanol, methanol **Storage:** -20°C

A qualitative standard. Mixture consists of equal amounts of the compounds listed.

methyl undecanoate	C11:0	methyl cis-9-hexadecenoate	C16:1 ⁹
methyl 2-hydroxydodecanoate	2-OH C10:0	methyl hexadecanoate	C16:0
methyl dodecanoate	C12:0	methyl 15-methylhexadecanoate	iso-C17:0
methyl tridecanoate	C13:0	methyl cis-9,10-methylenehexadecanoate	C17:0Δ ^{9,10}
methyl 2-hydroxydodecanoate	2-OH C12:0	methyl heptadecanoate	C17:0
methyl 3-hydroxydodecanoate	3-OH C12:0	methyl 2-hydroxyhexadecanoate	2-OH C16:0
methyl tetradecanoate	C14:0	methyl cis, cis-9,12-octadecadienoate	C18:2 ^{9,12}
methyl 13-methyltetradecanoate	iso-C15:0	methyl cis-9-octadecenoate	C18:1 ⁹
methyl 12-methyltetradecanoate	anteiso-C15:0	methyl trans-9-octadecenoate	C18:1 ⁹
methyl pentadecanoate	C15:0	methyl octadecanoate	C18:0
methyl 2-hydroxytetradecanoate	2-OH C14:0	methyl cis-9,10-methyleneoctadecanoate	C19:0Δ ^{9,10}
methyl 3-hydroxytetradecanoate	3-OH C14:0	methyl nonadecanoate	C19:0
methyl 14-methylpentadecanoate	iso-C16:0	methyl eicosanoate	C20:0

1075 **Volatile acid mix** **100 ml**
Qualitative mix

Appearance: liquid **Solvent:** water **Solubility:** water
Storage: 4-8°C

Contains: formic, acetic, propionic, isobutyric, n-butyric, isovaleric, n-valeric, isocaproic, n-caproic, and heptanoic acids

1077 **Non-volatile acid mix** **100 ml**
Qualitative mix

Appearance: liquid **Solvent:** water **Solubility:** water
Storage: 4-8°C

Contains: pyruvic, lactic, oxalacetic, oxalic, methyl malonic, malonic, fumaric and succinic acids.

Biochemical research standard mixes

These mixtures are prepared by precise gravimetric technique. All mixes contain equal amounts of listed components. A data sheet is supplied with each mixture.

1127 **Polar lipid mix** **25 mg/ml, 1 ml**
TLC standards mix

Source: natural, egg, ovine **Appearance:** liquid **Solvent:** chloroform/methanol 2:1
Solubility: chloroform/methanol 2:1 **Storage:** -20°C

Contains: cholesterol, phosphatidylethanolamine, lecithin, and lyso-lecithin

1128	Sphingolipid mix TLC standards mix	25 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol 2:1 Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Contains: cerebrosides, sulfatides, and sphingomyelin	
1129	Non-polar lipid mix A TLC standards mix	25 mg/ml, 1 ml
	Source: natural, plant, ovine Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	Contains: cholesteryl palmitate, tripalmitin, palmitic acid, and cholesterol	
1130	Non-polar lipid mix B TLC standards mix	25 mg/ml, 1 ml
	Source: natural, plant, ovine Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	Contains: cholesteryl oleate, methyl oleate, triolein, oleic acid, and cholesterol	

Glycosphingolipid mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505	Neutral glycosphingolipid qualmix, Glycosylceramides, qualitative mix	1 mg/ml, 1 ml
	Source: natural, bovine and porcine Appearance: liquid Solvent: chloroform/methanol 2:1 Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Contains: cerebrosides, lactosylceramide, ceramide trihexoside, globoside	
1508	Monosialoganglioside mix GM ₃ , GM ₂ , GM ₁ qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: GM ₃ , GM ₂ , GM ₁	
1509	Disialoganglioside mix GD ₃ , GD _{1a} , GD _{1b} , qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: GD ₃ , GD _{1a} , GD _{1b}	
1510	Lactosylceramide and sialosyl derivatives mix LC, GM ₃ , GD ₃ qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: LC, GM ₃ , GD ₃	

1511 **Gangliotetraosylceramide and sialosyl derivatives mix** **0.5 mg/ml, 1 ml**
 asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b} qualitative mix

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water 2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b}

Biochemicals and reagents

Stable isotope labeled compounds

1914 **N-Stearoyl-D₃₅-psychosine, perdeuterated** **5 mg**
 Cerebrosides with N-C18:0-D₃₅ fatty acid side chain C₄₂H₄₆D₃₅NO₈

Source: semi-synthetic, bovine **Mol. Wt.:** 762 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, hot ethanol, chloroform/methanol 2:1 **Storage:** -20°C

1533 **N-Palmitoyl-D₃-glucopsychosine, deuterated** **1 mg**
 N-C16:0-D₃-Glucopsychosine; glucocerebroside with C16:0-D₃ fatty acid side chain C₄₀H₇₄D₃NO₈

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 703 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

1534 **N-Palmitoyl-D₃-lactosylceramide, deuterated** **1 mg**
 N-C16:0-D₃-Lactosylceramide; lactosylceramide with C16:0-D₃ fatty acid side chain C₄₆H₈₄D₃NO₁₃

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 864 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol/water 5:1:0.1 **Storage:** -20°C

2200 **N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine** **1 mg**
 D-erythro-Sphingomyelin with 1-¹³C-palmitic acid; SPM with ¹³C labeled fatty acid ¹²C₃₈¹³CH₇₉N₂O₆P

Source: semi-synthetic, bovine **Mol. Wt.:** 703 **Purity:** 98+% by TLC **Appearance:** waxy solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

2050 **N-Octadecanoyl-D₃-monosialoganglioside GM₁** **0.5 mg**
 N-D₃-Stearoyl-GM₁ C₇₃H₁₂₈N₃O₃₁D₃


Source: semi-synthetic, bovine brain **Mol. Wt.:** 1548 **Purity:** 98+% by TLC **Appearance:** solid, **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C

1536 **N-Octadecanoyl-D₃-sulfatide** **1 mg**
 N-C18:0-D₃-Sulfatide C₄₂H₇₈D₃NO₁₁S

Source: semi-synthetic, bovine **Mol. Wt.:** 833 **Purity:** 98+% by TLC **Appearance:** off-white solid **Solubility:** chloroform/methanol/DI water 2:1:0.1 **Storage:** -20°C

1537 **N-Octadecanoyl-D₃-ceramide trihexoside** **0.5 mg**
 C18:0-D₃-CTH; C18:0-D₃-Gb3; N-Octadecanoyl-D₃-globotriaosylceramide
 C₅₄H₉₈D₃NO₁₈
Source: semi-synthetic, porcine **Mol. Wt.:** 1055 **Purity:** 98+% by TLC
Appearance: off-white solid **Solubility:** chloroform/methanol 2:1, DMSO
Storage: -20°C

Fluorescent compounds



1841 **N-Hexanoyl-NBD-D-erythro-sphingosine** **100 µg**
1841-001 N-C6:0-NBD-ceramide; N-C6:0-NBD-D-erythro-sphingosine, fluorescent; N-
 (NBD-aminocaproyl)-D-erythro-sphingosine C₃₀H₄₉N₅O₆ **CAS#:** 86701-10-2
1 mg

Source: synthetic **Mol. Wt.:** 575 **Melting Point (°C):** 85-88 **Purity:** 98+% by TLC
Appearance: orange film, vacuum dried **Solubility:** chloroform, ethanol, methanol
Storage: -20°C

Reference:
 J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1618 **N-Dodecanoyl-NBD-D-erythro-sphingosine** **100 µg**
1618-001 N-C12:0-NBD-ceramide; N-C12:0-NBD-D-erythro-sphingosine, fluorescent;
 N-(NBD-aminolauroyl)-D-erythro-sphingosine C₃₆H₆₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1857 **N-Hexanoyl-NBD-L-threo-sphingosine** **100 µg**
1857-001 N-C6:0-NBD-ceramide; N-C6:0-NBD-L-threo-sphingosine, fluorescent; N-
 (NBD-aminocaproyl)-L-threo-sphingosine C₃₀H₄₉N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 575 **Purity:** 98+% by TLC **Appearance:** red-orange
 solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Reference:
 J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1620 **N-Dodecanoyl-NBD-L-threo-sphingosine** **100 µg**
1620-001 N-C12:0-NBD-ceramide; N-C12:0-NBD-L-threo-sphingosine, fluorescent;
 N-(NBD-aminolauroyl)-L-threo-sphingosine C₃₆H₆₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1624 **N-Hexanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
1624-001 N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-L-threo-dihydrosphingosine,
 fluorescent; N-(NBD-aminocaproyl)-L-threo-dihydrosphingosine
 C₃₀H₅₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1623 1623-001	N-Dodecanoyl-NBD-L-threo-dihydrosphingosine N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-L-threo-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-dihydrosphingosine $C_{36}H_{63}N_5O_6$	100 µg 1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1626 1626-001	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-dihydrosphingosine $C_{30}H_{51}N_5O_6$	100 µg 1 mg
	Source: synthetic Mol. Wt.: 578 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1625 1625-001	N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-dihydrosphingosine $C_{36}H_{63}N_5O_6$	100 µg 1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1628 1628-001	N-Hexanoyl-NBD-phytosphingosine N-C6:0-NBD-phytoceramide; N-C6:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminocaproyl)-phytosphingosine $C_{30}H_{51}N_5O_7$	100 µg 1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 594 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1627 1627-001	N-Dodecanoyl-NBD-phytosphingosine N-C12:0-NBD-phytoceramide; N-C12:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminolauroyl)-phytosphingosine $C_{36}H_{63}N_5O_7$	100 µg 1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 678 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1912 1912-001	N-Hexanoyl-NBD-sphingosylphosphorylcholine N-C6:0-NBD-sphingomyelin, fluorescent; N-C6:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminocaproyl)-sphingomyelin $C_{35}H_{61}N_6O_9P$ CAS#: 94885-04-8	100 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 740 Purity: 98+% by TLC Appearance: red-brown solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1619 1619-001	N-Dodecanoyl-NBD-sphingosylphosphorylcholine N-C12:0-NBD-sphingomyelin, fluorescent; N-C12:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminolauroyl)-sphingomyelin $C_{41}H_{73}N_6O_4P$	100 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 825 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	

1621 1621-001	N-Hexanoyl-NBD-galactosylceramide N-C6:0-NBD-beta-D-galactosylsphingosine; N-C6:0-NBD-cerebrosides; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-galactosylsphingosine $C_{36}H_{59}N_5O_{11}$	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1622 1622-001	N-Hexanoyl-NBD-glucosylceramide N-C6:0-NBD-beta-D-glucosylsphingosine; N-C6:0-NBD-glucosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-glucosylsphingosine $C_{36}H_{59}N_5O_{11}$	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1629 1629-001	N-Hexanoyl-NBD-lactosylceramide N-Hexanoyl-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminocaproyl)-beta-D-lactosylsphingosine $C_{42}H_{69}N_5O_{16}$	50 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1630 1630-001	N-Dodecanoyl-NBD-lactosylceramide N-Dodecanoyl-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-beta-D-lactosylsphingosine $C_{48}H_{81}N_5O_{16}$	50 µg 1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1631 1631-001	N-Dodecanoyl-NBD-ceramide trihexoside N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl)-ceramide trihexoside $C_{54}H_{91}N_5O_{21}$	100 µg 1 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1145 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1; DMSO; hot methanol Storage: -20°C	
1632 1632-001	N-Dodecanoyl-NBD-sulfatide N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate; N-(NBD-aminolauroyl)-sulfatide $C_{42}H_{71}N_5O_{14}S$	100 µg 1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 901 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

Appendix

Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC.
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat. # 1044 Lecithin (egg)	Cat. # 1070 Lecithin (bovine)	Cat. # 1302 Lecithin (plant)	Cat. # 1045 Phosphatidyl- ethanolamine (egg)	Cat. # 1301 Phosphatidyl- ethanolamine (plant)	Cat. # 1046 lyso-Lecithin (egg)	Cat. # 1047 Phosphatidylserine (bovine)	Cat. # 1048 Phosphatidylinositol (plant)
Fatty Acids								
C14:0		trace						
C16:0	31	35	14	19	22	72	1	36
C16:1		1		1				
C18:0	16	14	4	26	3	24	42	
C18:1	31	33	11	23	7	3	27	7
C18:2	16		66	14	60			50
C18:3			6		8			7
C20:0							1	
C20:1							4	
C20:4				10			4	
C21:0								
C22:0							1	
C22:1							1	
C22:6							7	
C23:0								
C24:0								
C24:1								
C25:0								
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH								
C16:0 2-OH								
C18:0 2-OH								
C20:0 2-OH								
C22:0 2-OH								
C23:0 2-OH								
C24:0 2-OH								
C24:1 2-OH								
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	6	17	0	7	0	1	12	0
Total	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1051 Sphingomyelin (bovine)	Cat: # 1328 Sphingomyelin (porcine RBC)	Cat: # 1053 Phosphatidic acid (semi-synthetic)	Cat: # 1057 Glucocerebrosides (human)	Cat: # 1058 Monogalactosyl-diglycerides (plant)	Cat: # 1059 Digalactosyl diglyceride (plant)	Cat: # 1061 Monosialoganglioside GM ₁	Cat: # 1062 Disialoganglioside GD _{1a}
Fatty Acids								
C14:0							trace	trace
C16:0	3	30	39	14	23	9	1	1
C16:1								
C18:0	45	8	12	3	77	91	86	86
C18:1		2	34				3	3
C18:2			15					
C18:3								
C20:0	1	2		3			4	4
C20:1								
C20:4								
C21:0								
C22:0	4	10		24			2	2
C22:1								
C22:6								
C23:0		1		9			1	1
C24:0	8	30		33			1	1
C24:1	31	14		13			2	2
C25:0								
C25:1								
C26:0		2						
C26:1								
C27:0								
C27:1								
C14:0 2-OH								
C16:0 2-OH								
C18:0 2-OH								
C20:0 2-OH								
C22:0 2-OH								
C23:0 2-OH								
C24:0 2-OH								
C24:1 2-OH								
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	8	1	0	1	0	0	0	0
Total	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Cat. # 1063 Trisialoganglioside GT _{1b}	Cat. # 1064 Gangliotetraosylceramide	Cat. # 1065 Purified mixed gangliosides	Cat. # 1501 Disialoganglioside GD _{1b}	Cat. # 1050 Cerebrosides (bovine)	Cat. # 1066 Cerebrosides Keratin (bovine)	Cat. # 1138 Cerebrosides Phrenosin (bovine)	Cat. # 1049 Sulfatides (bovine)
Fatty Acids								
C14:0	trace	trace	trace	trace				
C16:0	1	1	1	1	trace	trace		trace
C16:1								
C18:0	86	86	86	86	4	5		5
C18:1	3	3	3	3				trace
C18:2								
C18:3								
C20:0	4	4	4	4	1	1		1
C20:1								
C20:4								
C21:0								
C22:0	2	2	2	2	4	9		7
C22:1						trace		trace
C22:6								
C23:0	1	1	1	1	2	5		
C24:0	1	1	1	1	10	25		18
C24:1	2	2	2	2	15	43		29
C25:0					3	3		2
C25:1					1	3		2
C26:0					2	2		1
C26:1					1	4		3
C27:0					2			1
C27:1					2			
C14:0 2-OH								
C16:0 2-OH								
C18:0 2-OH					15		36	5
C20:0 2-OH					1		1	trace
C22:0 2-OH					6		8	3
C23:0 2-OH					5		6	
C24:0 2-OH					17		25	10
C24:1 2-OH					6		9	6
C25:0 2-OH					3		4	2
C25:1 2-OH							2	
C26:0 2-OH							2	
C26:1 2-OH							2	
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	0	0	0	0	0	0	5	5
Total	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1067 Ceramide trihexoside (porcine)	Cat: # 1068 Globosides (porcine)	Cat: # 1118 Esterified steryl glucoside	Cat: # 1056 Ceramides (bovine)	Cat: # 1322 Ceramides (non-hydroxy)	Cat: # 1323 Ceramides (hydroxy)	Cat: # 1535 Monosialoganglioside GM ₄	Cat: # 1332 Sphingomyelin, (egg, chicken)	Cat: # 1516 Tetrasialoganglioside GQ1 _b
Fatty Acids									
C14:0								trace	
C16:0	3	2	34	trace			4	72	5
C16:1									1
C18:0	2	1	8	4	11		2	8	80
C18:1	2		8					3	2
C18:2			36						3
C18:3			4						
C20:0	2	1	1	1	2		trace	2	4
C20:1							trace		
C20:4									
C21:0									
C22:0	17	15	4	4	10		3	5	2
C22:1							4		
C22:6									
C23:0	1	1	2	2	6		4	1	
C24:0	29	23	2	10	24		6	2	
C24:1	5	15		15	31		4	4	
C25:0				9	3				
C25:1				1	3				
C26:0		1		2	2				
C26:1		1		1	3				
C27:0				2					
C27:1				2					
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH		trace		15		24	1		
C20:0 2-OH		trace		1		1	3		
C22:0 2-OH	3	3		6		8	25		
C23:0 2-OH	1			5		6	17		
C24:0 2-OH	19	11		17		35	18		
C24:1 2-OH	10	23				17	7		
C25:0 2-OH				3		4			
C25:1 2-OH									
C26:0 2-OH									
C26:1 2-OH						2			
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	6	3	1	0	5	3	2	3	3
Total	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1329 Sphingomyelin (buttermilk)	Cat: # 1500 Lactosyl ceramide (porcine)	Cat: # 1507 Lactosyl ceramide (buttermilk)	Cat: # 1502 Monosialoganglioside GM2	Cat: # 1503 Monosialoganglioside GM3 (buttermilk)	Cat: # 1504 Disialoganglioside GD3 (buttermilk)	Cat: # 1521 Glucocerebrosides (buttermilk)	Cat: # 1522 Glucocerebrosides (plant)
Fatty Acids								
C14:0	1						trace	
C16:0	21	14	8	2	6	8	15	
C16:1								
C18:0	3	6	3	82	1	1	3	
C18:1		4	trace					
C18:2								
C18:3								
C20:0	1	1	2	7	1	1	2	
C20:1								
C20:4								
C21:0					1	2		
C22:0	23	9	28	4	23	24	31	
C22:1								
C22:6								
C23:0	29	1	33	trace	36	35	28	
C24:0	20	15	20	1	22	21	17	
C24:1	1	5	3	2	3	3		
C25:0								
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH								trace
C16:0 2-OH								79
C18:0 2-OH		trace						trace
C20:0 2-OH								
C22:0 2-OH		8						8
C23:0 2-OH								1
C24:0 2-OH		24						9
C24:1 2-OH		13						
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	1	0	3	2	7	5	4	3
Total	100	100	100	100	100	100	100	100

Spingolipid Metabolism Pathways


Matrea
Lipids for Research
168 Treveler Street
Pleasant Gap, PA 16823 USA
RE: 814 359 5060 Fax: 814 359 5062
e-mail: customer-service@matrea.com Web: www.matrea.com

Ceramide: Its Cycles and Physiological Actions
Ceramide is a key and stable component of the lipid bilayer of cell membranes. It is formed predominantly by the condensation of sphingosine, and by the addition of fatty acids. In addition to its role in membrane structure, ceramide is involved in cell signaling and apoptosis. It is synthesized in the endoplasmic reticulum and Golgi apparatus. Ceramide is a key component of the lipid bilayer of cell membranes. It is formed predominantly by the condensation of sphingosine, and by the addition of fatty acids. In addition to its role in membrane structure, ceramide is involved in cell signaling and apoptosis. It is synthesized in the endoplasmic reticulum and Golgi apparatus.

Synthetic Ceramides
Ceramide with short side chains has been shown to enter readily into cells where they are hydrolyzed to sphingosine and fatty acids. These synthetic ceramides are used in the study of the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide.

Enzyme Inhibitors
Enzyme inhibitors are used to study the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide.

Sphingolipids Sold by Matrea
Matrea offers a wide variety of sphingolipids for research. These include sphingosine, ceramide, sphingomyelin, and ganglioside. They are used in the study of the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide.



Glycosphingolipids
Glycosphingolipids are a class of sphingolipids that have one or more sugar units attached to the sphingosine backbone. They are found in the cell membrane and play a role in cell signaling and recognition. They are used in the study of the function and metabolism of ceramide. They are used in the study of the function and metabolism of ceramide.

Spingolipid Structures and Pathways Wall Chart

In a clear and straightforward manner, this wall chart indicates the structures and relationships between most commonly discussed sphingolipids. Full size copies (approximately 35 x 26 inches) are available on request to customer service.

Literature References

1. B.A. Fenderson, E.M. Eddy, S.Hakomori, *BioEssays* **12**, 173, 1990.
2. A. Gorio et al., *Exp. Brain Res.*, Suppl. **13**, 283, 1984.
3. F. Di Gregorio et al., *Neuropediatrics*, Suppl. **15**, 93, 1984.
4. J.S. Schneider et al., *Science*, **256**, 843, 1992.
5. R.W. Leeden, R. K. Yu, *Methods Enzymol.*, **83**, 139, 1982.
6. M. Faucher et al., *J. Biol. Chem.* **263**, 5319, 1988.
7. Y. Hannun, *ibid.*, **269**, 3125, 1994.
8. R. Kolesnick, D.W. Golde, *Cell*, **77**, 325, 1984.
9. J. M. L. Hauser et al., *J. Biol. Chem.* **269**, 6803, 1994.
10. A. Gómez-Muñoz et al., *ibid.*, **270**, 26318, 1995.
11. B. M. Buehrer, R.M. Bell, *Adv. in Lipid Res.*, **26**, 59, 1993.
12. C. W. Sachs et al., *J. Biol. Chem.*, **270**, 26639, 1995.
13. R.R, Vunnam, N.S. Radin, *Chem. Phys. Lipids*, **265**, 1980
14. J. Inokuchi and N.S. Radin, *J. Lipid Res.*, **28**, 565, 1987.
15. N.S. Radin et al. *J. Biochem*, **111**, 191, 1992.
16. J. Inokuchi et al., *Cancer letters*, **38**, 23, 1987.
17. Y. Lavie et al., *J. Biol. Chem.*, **271**, 19530, 1996.
18. Y. Lavie et al., *J. Biol. Chem.*, **271**, in press, 1996.
19. Y. Hannun et al., *Science*, **235**, 670, 1987.
20. S. Spiegel et al., *Proc. Intern. Conf. Biol. Function Glycosphingolipids*, Santa Barbara, CA, 1990.
21. M. Sugita et al. *Biochim. Biophys. Acta*, **398**, 125, 1975.
22. A. Bielawska et al. *J. Biol. Chem.*, **271**, 12646, 1996.
23. A. Bielawska et al. *ibid.* **267**, 18493, 1992.
24. M.W. Pariza et al., *Cancer Res.* **43**, 2444s, 1983.
25. Y. L Ha et al., *J. Agr. Food Chem.*, **37**, 75, 1989.
26. S. Banni et al. *Abstr. 87th AOCS Mtg. 1996*, p.28.
27. C. Ip et al., *Cancer Res.* **51**, 6118, 1991.
28. M. A. Belury, *Nutr. Rev.* **53** (4 Pt. 1), 83, 1995.
29. C. Liew et al. *Carcinogenesis*, **16**, 3037, 1995.
30. T. D. Shultz et al. *Cancer Let*, **63**, 125, 1992.
31. R.J. Nicolosi, *Abstr. 87th AOCS Mtg.*, 1996.
32. K.N. Lee et al., *Atheroscl.*, **108**(1), 19, 1994.
33. B.F. Haumann, *Inform*, **7**(2), 152, 1996.
34. M.W. Pariza et al., *Abstr.*, 87th AOCS Mtg., 1996.
35. P.W. Parodi, *Austral. J. Dairy Tech.*, **49**, 93, 1994.
36. M.A. Belury et al., *Lipids*, **32**, 199, 1997.
37. W.W. Christie et al. *JAOCS*, **74**, 1231, 1997.
38. Sehat, N. et al., *Lipids*, **33**, 217, 1998.
39. Sébédio, J. L. et al., *Biochim. Biophys. Acta*, 1345,5, 1997.
40. Houseknecht, K.L. et al., *Biochem. Biophys. Res. Comm.*, **224**, 678, 1998.
41. M. Mattie, et. al., *J. Biol. Chem.* **269**:3181, 1994
42. Shayman, J. A., *Kidney Int.* **58**, 11-26, 2000
43. Kolesnick, R. N. et al. *J. Cell Physiol.* **184**, 285-300, 2000
44. Smith, W. L. and Merrill, A. H., Jr. *J. Biol. Chem.* **277**, 25841-25842, 2002
45. Pyne, S. And Pyne, N. J. *Biochem. J.* **349**, 385-402, 2000
46. Merrill, A. H. Jr. *J. Biol. Chem.* **277**, 25843-25846, 2002
47. Hannun, Y. A. and Obeid, L., *J. Biol. Chem.* **277**, 25847-25850, 2002
48. van Meer, G. And Lisman, Q. *J. Biol. Chem.* **277**, 25855-25858, 2002
49. Kolter, T., Proia, R. L. and Sandhoff, K. *J. Biol. Chem.* **277**, 25859-25862, 2002
50. Radin, N. S., *Cancer Invest.* **20**, 779-786, 2002
51. Tettamanti, G., Bassi R., Viani, P., Riboni, L., *Biochimie* **85**(3-4), 423-437, 2003
52. Fredman, P., Hedberg, K., Brezicka, T., *Biodrugs* **17**(3) 155-167, 2003
53. Hoeckstra, D., et al. *J. Lipid Res.* **44**, 869-877, 2003
54. Radin, N. S., *Biochem. J.* **371**:243, 2003
55. Radin, N. S., *Biorg. Med. Chem.* **11**:2123, 2003
56. Radin, N. S., *Cancer Investigation* **20**:1, 2002
57. Borman, S., *Chemical and Engineering News* **82**:32, 31-35, 2004
58. Jacqueline M. Kravcka, Li Li, Zdzislaw M. Szulc, Jacek Bielawski, Besium Ogretmen, Yusuf A. Hannun, Lina M. Obeid, and Alicja Bielawska. *J. Biol. Chem.*, **10**, 1074/jbc. M700647200, February 5, 2007

Cross reference for product numbers and catalog pages

1006	73	1082	79	1194	56
1008	49	1083	78	1195	71
1009	49	1084	78	1196	48
1010	50	1085	78	1197	48
1011	50	1086	79	1198	48
1012	50	1087	79	1199	49
1013	50	1088	79	1200	48
1014	50	1089	79	1203	53
1015	50	1093	76	1204	53
1016	52	1095	81	1205	55
1017	52	1097	81	1206	55
1018	50	1098	82	1207	70
1019	50	1099	82	1208	53
1020	50	1100	82	1241	51
1021	50	1101	82	1242	51
1022	53	1102	82	1243	52
1023	53	1103	82	1244	56
1024	54	1104	82	1245	59
1025	54	1105	83	1245-1	59
1026	54	1106	82	1245-10	59
1027	54	1108	82	1247-1	60
1028	51	1113	74	1247-10	60
1029	51	1114	83	1248	60
1030	51	1115	74	1248-1	60
1031	51	1116	74	1249	60
1032	55	1117	75	1249-1	60
1033	55	1118	75, 93	1249-10	60
1034	55	1119	74	1251	52
1035	51	1120	74	1252	52
1036	51	1121	74	1254	60
1037	52	1122	74	1255	59
1038	52	1123	74	1256	60
1040	52	1124	76	1257	59
1041	56	1125	77	1258	60
1042	55	1127	83	1259	60
1044	40, 90	1128	84	1261	49
1045	41, 90	1129	84	1262	53, 57
1046	40, 90	1130	84	1263	53, 57
1047	40, 90	1131	58, 77	1264	56
1048	41, 90	1136	56	1265	56
1049	22, 92	1138	20, 92	1266	54
1050	20, 92	1147	52, 57	1267	54
1051	15, 91	1148	53, 57	1269	55
1051-1	15	1149	53, 57	1301	41, 90
1052	41	1150	53, 57	1302	40, 90
1053	41, 91	1151	54, 57	1303	47
1056	12, 93	1152	54, 57	1303-2	47
1057	21, 91	1153	54	1304	2
1058	48, 91	1154	54	1305	20
1059	48, 91	1155	56	1306	21
1061	29, 91	1156	57	1310	22
1061-50	29	1157	52	1318	17
1062	30, 91	1161	49	1319	17
1063	30, 92	1162	49	1320	6
1064	28, 92	1163	49	1321	17
1065	31, 92	1164	49	1321-05	17
1066	20, 92	1165	49	1322	12, 93
1067	24, 93	1166	49	1322-05	12
1067-10	24	1167	56	1323	12, 93
1068	25, 93	1175	56	1323-05	12
1069	41	1177	76	1324	4
1070	40, 90	1179	55	1325	20
1071	72	1181	58, 59	1326	4
1072	72	1182	68	1327	17
1073	72	1183	68	1328	16, 91
1074	73	1186	51	1329	16, 94
1075	83	1187	51	1330	5
1077	83	1192	55	1330-1	5
1081	76	1193	55	1332	16, 93

1332-1	16	1615	70	1727-0.5	64
1333	9	1616	70	1728	64
1334	20	1618	14, 86	1728-0.5	64
1334-50	20	1618-001	14, 86	1729	65
1335	20	1619	19, 87	1729-0.5	65
1336	41	1619-001	19, 87	1730	65
1400	43	1620	14, 86	1730-0.5	65
1409	61	1620-001	14, 86	1731	65
1410	61	1621	21, 26, 88	1731-0.5	65
1411	61	1621-001	21, 26, 88	1732	65
1425	42	1622	21, 26, 88	1732-0.5	65
1426	43	1622-001	21, 26, 88	1733	65
1427	43	1623	14, 87	1733-0.5	65
1428	42	1623-001	14, 87	1734	65
1429	42	1624	14, 86	1734-0.5	65
1430	42	1624-001	14, 86	1735	65
1431	43	1625	15, 87	1735-0.5	65
1432	43	1625-001	15, 87	1736	65
1433	44	1626	15, 87	1736-0.5	65
1434	44	1626-001	15, 87	1739	65
1435	44	1627	15, 87	1739-0.5	65
1436	44	1627-001	15, 87	1740	65
1437	43	1628	15, 87	1740-0.5	65
1438	44	1628-001	15, 87	1741	66
1439	44	1629	24, 26, 88	1741-0.5	66
1442	42	1629-001	24, 26, 88	1742	66
1443	43	1630	24, 26, 88	1742-0.5	66
1444	44	1630-001	24, 26, 88	1743	66
1445	43	1631	25, 27, 88	1743-0.5	66
1500	23, 94	1631-001	25, 27, 88	1744	66
1501	30, 92	1632	23, 27, 88	1744-0.5	66
1502	29, 94	1632-001	23, 27, 88	1745	64
1503	29, 94	1656	69	1745-0.5	64
1504	30, 94	1657	69	1746	64
1505	31, 84	1701	62	1746-0.5	64
1507	23, 94	1701-1	62	1747	64
1507-50	23	1702	62	1747-0.5	64
1508	31, 84	1702-1	62	1748	64
1509	31, 84	1703	62	1748-0.5	64
1510	31, 84	1703-1	62	1749	37
1511	32, 85	1704	62	1750	39
1512	28	1704-1	62	1751	34, 71
1513	24	1705	62	1752	39
1514	24	1705-1	62	1753	37
1516	30, 93	1706	62	1754	66
1517	23	1706-1	62	1754-0.5	66
1518	31	1707	62	1755	37
1520	24	1707-1	62	1756	38
1521	21, 94	1708	62	1757	35, 72
1521-50	21	1708-1	62	1758	62
1522	21, 94	1709	63	1758-1	62
1522-100	21	1709-0.5	63	1759	62
1523	24	1710	63	1759-1	62
1524	25	1710-0.5	63	1760	66
1525	31	1711	63	1761	66
1526	30	1711-0.5	63	1766	68
1531	22	1712	63	1773	45
1532	23	1712-0.5	63	1773-1	45
1533	25, 85	1713	63	1773-5	45
1534	26, 85	1714	63	1774	46
1535	29, 93	1715	63	1774-1	46
1536	23, 26, 85	1716	63	1774-5	46
1537	25, 26, 86	1717	39	1775	47
1600	69	1718	39	1775-1	47
1601	69	1719	37	1775-5	47
1602	69	1720	37	1777	46
1603	69	1722	63, 77	1777-1	46
1605	69	1725	64	1777-5	46
1606	69	1725-0.5	64	1778	46
1612	70	1726	64	1778-1	46
1613	70	1726-0.5	64	1778-5	46
1614	70	1727	64	1779	45

1779-1	45	1863	75	2036	13
1780	45	1865	38	2037	9
1780-1	45	1868	38	2038	9
1780-5	45	1875	22	2039	9
1781	46	1876	5	2041	11
1781-1	46	1877	68	2042	10
1781-5	46	1878	68	2043	11
1782	47	1879	68	2044	10
1782-1	47	1880	68	2045	11
1782-5	47	1881	66	2046	18
1783	47	1882	66	2047	12
1783-1	47	1883	67	2050	29, 85
1783-5	47	1884	67	2076	22
1784	46	1886	36	2200	17, 85
1784-1	46	1886-005	36	4210	75
1784-5	46	1887	36		
1786	34, 71	1887-005	36		
1790	73	1888	23		
1791	70	1889	36		
1792	70	1890	16		
1793	61	1891	5		
1794	61	1892	5		
1797	73	1893	5		
1800	38	1894	13		
1802	2	1895	13		
1803	18	1896	6		
1805	6	1897	13		
1806	2	1898	68		
1807	3, 35	1899	68		
1807-025	3, 35	1900	8		
1809	8	1901	7		
1810	9	1903	8		
1815	68	1904	22		
1818	67	1907	16		
1819	67	1909	16		
1822	71	1910	11		
1823	71	1911	16		
1826	2	1912	18, 87		
1827	2	1912-001	18, 87		
1828	8	1913	18		
1829	8	1914	25, 85		
1830	9	1915	9		
1831	4	1916	10		
1831-1	4	1917	16		
1832	9	1918	17		
1833	2	1919	46		
1834	11	1919-1	46		
1835	3	1919-5	46		
1837	3	1920	46		
1838	3	1920-1	46		
1839	5	1920-5	46		
1840	3	1921	47		
1841	14, 86	1921-1	47		
1840-001	14, 86	1921-5	47		
1842	10	1922	46		
1843	10	1922-1	46		
1845	4	1922-5	46		
1846	4	1923	46		
1847	8	1923-1	46		
1848	8	1923-5	46		
1850	10	1950	32		
1851	4	1951	32		
1852	18	1954	32		
1854	11	1957	33		
1855	10	1960	33		
1856	10	1961	33		
1857	14, 86	1962	33		
1857-001	14, 86	1977	32		
1858	39	2009	75		
1859	35	2010	76		
1860	35	2034	13		
1862	75	2035	13		

Product name index

(+)-delta-Tocopherol	73	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol	44	3-Hydroxy C18:0 methyl ester	66
β-Sitostanol	74	1-Palmitoyl-sn-glycero-3-phosphorylcholine	43	3-Hydroxy C6:0 acid	64
1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine	43	1-Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine	61	3-Hydroxy C6:0 methyl ester	64
1,2-Dilauroyl-sn-glycero-3-phosphorylcholine	42	1-Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine	61	3-Hydroxy C8:0 acid	64
1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine	44	1-Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine	61	3-Hydroxy C8:0 methyl ester	64
1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol	43	2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole	39	3-Hydroxy C9:0 acid	64
1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid	42	2,2-Difluoropalmitic acid	39	3-Hydroxy C9:0 methyl ester	64
1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine	42	2,3-Dihydroxy C16:0 fatty acid methyl ester	68	3-Hydroxydecanoic acid	64
1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine	44	2,6-Dimethyl C7:0 fatty acid	70	3-Hydroxydodecanoic acid	65
1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol	43	20-Hydroxyeicosanoic acid	67	3-Hydroxyheptadecanoic acid	66
1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid	42	21-Hydroxyheneicosanoic acid	67	3-Hydroxyhexadecanoic acid	65
1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine	43	22-Hydroxydocosanoic acid	67	3-Hydroxyhexanoic acid	64
1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine	44	2-Fluoropalmitic acid	39	3-Hydroxynonanoic acid	64
1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol	43	2-Hydroxy C10:0 acid	62	3-Hydroxyoctadecanoic acid	66
1,2-Distearoyl-phosphatidyl ethanolamine-methyl-polyethylene-glycol conjugate-2000	44	2-Hydroxy C10:0 methyl ester	62	3-Hydroxyoctanoic acid	64
1,2-Distearoyl-sn-glycero-3-phosphatidic acid	42	2-Hydroxy C12:0 acid	62	3-Hydroxytetradecanoic acid	65
1,2-Distearoyl-sn-glycero-3-phosphorylcholine	43	2-Hydroxy C12:0 methyl ester	62	3-Hydroxytridecanoic acid	65
1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine	44	2-Hydroxy C14:0 acid	62	3-Hydroxyundecanoic acid	65
1,2-Distearoyl-sn-glycero-3-phosphorylglycerol	44	2-Hydroxy C14:0 methyl ester	62	3-keto-C6-Dihydrosphingosine•HCl	5
1,6,7,8-tetrahydroxyoctahydroindolizine	38	2-Hydroxy C16:0 acid	62	3-keto-C8-Dihydrosphingosine•HCl	5
1:1:1 sterol:glucose:fatty acid	75	2-Hydroxy C16:0 methyl ester	62	3-keto-C12-Dihydrosphingosine•HCl	5
10(E),12(Z)-Octadecadienoic acid	59, 60	2-Hydroxy C18:0 acid	62	3-keto-Dihydrosphingosine•HCl	5
10-Hydroxy-2-(E)-decanoic acid	66	2-Hydroxy C18:0 methyl ester	62	3-keto-Sphinganine hydrochloride	5
10-Methyl C16:0 fatty acid	70	2-Hydroxy C20:0 acid	63	4-Hydroxysphinganine	5
10-Methyl C16:0 fatty acid methyl ester	70	2-Hydroxy C20:0 methyl ester	63	5,7,8-Trimethyltocol	72
10-Methylhexadecanoic acid	70	2-Hydroxy C22:0 acid	63	5,8,11,14(Z,Z,Z,Z)-Eicosatetraenoyl-2'-hydroxy-ethyl-amide	35, 72
10-trans, 12-cis CLA	59, 60	2-Hydroxy C22:0 methyl ester	63	5,8-Dimethyltocol	72
11(Z), 13(E)-Octadecadienoic acid	59, 60	2-Hydroxy C23:0 acid	63	5-alpha-Cholestane	74
11-cis, 13-trans CLA	59, 60	2-Hydroxy C23:0 methyl ester	63	5-beta-Cholestane-3-beta-ol	74
11-Hexadecenoic acid (92% cis, 8% trans)	53	2-Hydroxy C24:0 acid	63	6-Hydroxy C18:0 fatty acid	68
12-Hydroxy C18:1 (9-trans) fatty acid	68	2-Hydroxy C24:0 methyl ester	63	6-Hydroxyoctadecanoic acid	68
12-Hydroxy C18:1 (9-trans) methyl ester	68	2-Hydroxy methyl ester mix	63, 77	7,8-Dimethyltocol	72
12-Methyltetradecanoic acid	70	2-Hydroxydecanoic acid	62	8-(5-Hexyl-2-furyl)-octanoic acid	61
13-Methyltetradecanoic acid	69	2-Hydroxydodecanoic acid	62	8-Methyltocol	73
14-Methylhexadecanoic acid	70	2-Hydroxyeicosanoic acid	63	9(E),11(E)-Octadecadienoic acid	58, 59
15-Hydroxypentadecanoic acid	6866	2-Hydroxyhexadecanoic acid	62	9(Z),11(E)-Octadecadienoic acid	59, 60
15-Methylhexadecanoic acid	69	2-Hydroxyoctadecanoic acid	62	9(Z),11(Z)-Octadecadienoic acid	60
17-Hydroxyheptadecanoic acid	66	2-Hydroxytricosanoic acid	63	9,12-epoxy-9,11-octadecadienoic acid	61
1-beta-D-galactosylsphingosine	20	3,7,11,15-Tetramethylhexadecanoic acid	71	9-cis, 11-cis CLA	60
1-beta-D-glucosylsphingosine	21, 22	3-Hydroxy C10:0 acid	64	9-cis, 11-trans CLA	59, 60
1-Hydroxy-2-amino-3-keto-dodecane • HCl	5	3-Hydroxy C10:0 methyl ester	64	9-trans, 11-trans CLA	58, 59
1-Hydroxy-2-amino-3-keto-hexane • HCl	5	3-Hydroxy C11:0 acid	65		
1-Hydroxy-2-amino-3-keto-octane • HCl	5	3-Hydroxy C11:0 methyl ester	65	A	
1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine	43	3-Hydroxy C12:0 acid	65	Alditol acetate mix-1	76
		3-Hydroxy C12:0 methyl ester	65	Alditol acetate mix-2	77
		3-Hydroxy C13:0 acid	65	Anandamide	35, 72
		3-Hydroxy C13:0 methyl ester	65	Anteiso-C15 acid	70
		3-Hydroxy C14:0 acid	65	Anteiso-C15 methyl ester	70
		3-Hydroxy C14:0 methyl ester	65	Anteiso-C16 methyl ester	70
		3-Hydroxy C16:0 acid	65	Anteiso-C17 acid	70
		3-Hydroxy C16:0 methyl ester	65	Anteiso-C17 methyl ester	70
		3-Hydroxy C17:0 acid	66	Anteiso-Heptadecanoic acid	70
		3-Hydroxy C17:0 methyl ester	66	Anteiso-Heptadecanoic methyl ester	70
		3-Hydroxy C18:0 acid	66	Anteiso-Palmitic methyl ester	70
				Anteiso-Pentadecanoic acid	70
				Anteiso-Pentadecanoic methyl ester	70
				Anti-ganglioside asialo GM ₁	32
				Anti-ganglioside asialo GM ₂	32
				Anti-ganglioside GD ₃	32
				Anti-ganglioside GM ₁	32
				Anti-ganglioside GM ₂ (NANA)	33
				Anti-ganglioside GM ₂ (NGNA)	33
				Anti-ganglioside GM ₄	33
				Anti-globoside GL-4	33
				AOCS rapeseed oil reference mix	78
				AOCS reference mix RM-1	78, 80
				AOCS reference mix RM-2	79, 80
				AOCS reference mix RM-3	79, 80

AOCS reference mix RM-4	79, 80	C20:0 methyl ester	51	Coprostanol	74
AOCS reference mix RM-5	79, 80	C20:1 (cis-11) acid	55	CTH	24, 93
AOCS reference mix RM-6	79, 80	C20:1 (cis-11) methyl ester	55	CTH with hydroxy fatty acid side chain	24
AOCS reference mix RM-7 kit	79	C20:2 (cis, cis-11, 14) acid	55	CTH with non-hydroxy fatty acid side chain	24
Arachidic acid	51	C20:2 (cis, cis-11, 14) methyl ester	55	Custom FAME mixes	80
Arachidonic acid	55	C20:3 (all cis-5,8,11) methyl ester	55	D	
Arachidonylethanolamide	35, 72	C20:4 (all cis-5,8,11,14) acid	55	D,L-2,6-Dimethylheptanoic acid	70
Asialo GM ₁	28	C20:4 (all cis-5,8,11,14) methyl ester	55	D,L-C16-Dihydrosphingosine	4
Asialo-GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b} qualitative mix	32, 86	C20:5 (all cis-5,8,11,14,17) acid	56	D,L-erythro-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol.HCl	37
Asialo-GM ₂	28	C20:5 (all cis-5,8,11,14,17) methyl ester	56	D,L-erythro-1-Phenyl-2-hexadecanoyl-amino-3-morpholino-1-propanol.HCl	37
B		C21:0 fatty acid	51	D,L-erythro-C20-Dihydrosphingosine	5
Bacterial acid methyl esters CP mix	83	C21:0 methyl ester	51	D,L-erythro-Dihydrosphingosine	4
Bacterial lipid standard	83	C22:0 fatty acid	51	D,L-erythro-PDMP	37
Behenic acid	51	C22:0 methyl ester	51	D,L-erythro-PPMP	37
beta-Sitostanol	74	C22:1 (cis-13), erucic acid	56	D,L-erythro-Sphinganine, C18 chain	4
		C22:1 (cis-13) methyl ester	56	D,L-erythro-Sphinganine, C20 chain	5
		C22:5 (all cis-7,10,13,16,19) acid	56	D,L-Sphinganine	5
		C22:5 (all cis-7,10,13,16,196 methyl ester	56	D,L-Sphinganine with C16 chain	4
		C22:6 (all cis-4,7,10,13,16,19) omega-3 fatty acid	56	D,L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol.HCl	37
		C22:6 (all cis-4,7,10,13,16,19) methyl ester	56	D,L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol.HCl	37
C		C23:0 fatty acid	51	D,L-threo-PDMP	37
C10:0 methyl ester	49	C23:0 methyl ester	51	D,L-threo-PPMP	37
C11:0 fatty acid	49	C24:0 fatty acid	52	D-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol	35
C11:0 methyl ester	49	C24:0 methyl ester	52	D-erythro-C10-Sphingosine	3
C12:0 acid	49	C24:1 (cis-15) acid	56	D-erythro-C12-Sphingosine	3
C12:0 methyl ester	49	C24:1 (cis-15) methyl ester	57	D-erythro-C14-Sphingosine	2
C13:0 fatty acid	49	C26:0 acid	52	D-erythro-C16-Sphingosine	3
C13:0 methyl ester	49	C26:0 methyl ester	52	D-erythro-C20-Dihydrosphingosine	4
C14:0 acid	50	C6:0 methyl ester	48	D-erythro-C20-Sphingosine	3
C14:0 methyl ester	50	C7:0 fatty acid	48	D-erythro-Dihydrosphingosine	4
C14:1 (cis-9) fatty acid	52	C7:0 fatty acid methyl ester	48	D-erythro-Dihydrosphingosine-1-phosphate	18
C14:1 (cis-9) methyl ester	52	C8:0 acid	48	D-erythro-SPC	17
C15:0 fatty acid	50	C8:0 methyl ester	49	D-erythro-Sphinganine, C18 chain	4
C15:0 methyl ester	50	C9:0 fatty acid	49	D-erythro-Sphinganine, C20 chain	4
C16:0 fatty acid	50	C9:0 methyl ester	49	D-erythro-Sphingomyelin with 1- ¹³ C-palmitic acid	17, 85
C16:0 methyl ester	50	Caprylic acid	48	D-erythro-Sphingosine	2
C16:1 (cis-11) acid	53	Castanospermine	38	D-erythro-Sphingosine-1-phosphate	18
C16:1 (cis-9) acid	52	CDH, ceramide beta-lactoside	23, 94	D-erythro-Sphingosyl-phosphorylcholine	17
C16:1 (cis-9) methyl ester	52	Ceramide beta-D-glucoside	21, 94	DGDG (hydrogenated, distearoyl)	48
C16:1 (trans-9) acid	52, 57	Ceramide-galactoside-3-sulfate	22, 92	DHA	56
C16:1 (trans-9) methyl ester	53, 57	Ceramide trihexosides	24, 93	DHDPC	43
C17:0 fatty acid	50	Ceramide trihexosides (bottom spot)	24	Digalactosyldiglyceride	48, 91
C17:0 methyl ester	50	Ceramide trihexosides (top spot)	24	Dihydrosphingosylphosphorylcholine	18
C17:1 (cis-10) acid	53	Ceramide-1-phosphorylcholine	15, 16	Dihydrosterculic acid	71
C17:1 (cis-10) methyl ester	53	Ceramides	12, 24, 93, 94	Disialoganglioside GD _{1a}	30, 91
C18:0-D ₃ -Gb3	25, 26, 87	Ceramides with hydroxy and nonhydroxy acyl groups	12	Disialoganglioside GD _{1b}	30, 92
C18:0-D ₃ -CTH	25, 26, 86	Ceramides with mostly hydroxy acyl groups	12	Disialoganglioside GD ₃	30, 94
C18:0 fatty acid	50	Ceramides with mostly non-hydroxy acyl groups	12	Disialoganglioside mix	31, 84
C18:0 methyl ester	50	Cerebronic acid	63	DLPC	42
C18:1 (cis-9) acid	53	Cerebroside sulfate	22	DLPE	44
C18:1 (cis-9) methyl ester	53	Cerebroside, Kerasin	20, 92	DLPG	43
C18:1 (cis-11) acid	54	Cerebroside, Phrenosin	20, 92	D-MAPP	35
C18:1 (cis-11) methyl ester	54	Cerebrosides	20, 92	DMPA	42
C18:1 (trans-9) acid	53, 57	Cerebrosides with C2:0 fatty acid	20	DMPC	42
C18:1 (trans-9) methyl ester	53, 57	Cerebrosides with N-C18:0-D ₃₅ fatty acid side chain	25, 85	DMPE	44
C18:1 (trans-11) acid	53, 57	Cerotic acid	52	DMPG	43
C18:1 (trans-11) methyl ester	53, 57	Cholestane	74	Docosahexaenoic acid	56
C18:2 (cis,cis-9,12) acid	54	Cholesterol	73	Docosanoic acid	51
C18:2 (cis,cis-9,12) methyl ester	54	cis-11-Octadecenoic acid	54	Docosapentaenoic acid	56
C18:2 (trans, trans-9, 12) acid	54, 57	cis-6-Hexadecenoic acid	52		
C18:2 (trans, trans-9,12) methyl ester	54, 57	cis-9,10-Methyleneoctadecanoic acid	71		
C18:3 (all cis-6,9,12) acid	54	cis-trans isomer standard	58, 77		
C18:3 (all cis-6,9,12) methyl ester	54	cis-vaccenic acid	54		
C18:3 (all cis-9,12,15) acid	54	Conduritol B epoxide	36		
C18:3 (all cis-9,12,15) methyl ester	54				
C19:0 fatty acid	51				
C19:0 methyl ester	51				
C19:1 (cis-10) acid	55				
C19:1 (cis-10) methyl ester	55				
C20:0 fatty acid	51				

Docosenoic acid	56	GD _{1a}	30, 91	Lactosylceramide	23, 94
Dodecanoic acid	49	GD _{1b}	30, 92	Lactosylceramide and sialosyl	
DOPI-3,4,5-P3	47	GD ₃	30, 94	derivatives mix	31, 84
DOPI-4,5-P2	46	GD ₃ , GD _{1a} , GD _{1b} , qualitative mix	31, 84	Lactosylceramide with C16:0-D ₃ fatty	
DOPI-4-P	46	Gg3	28	acid side chain	26, 85
DOPI-5-P	46	Gg4	28	Lactosylceramide with C16:0 fatty acid	
DPPA	42	GLC-10 mix	81	side chain	23
DPPC	43	GLC-100 mix	81, 82	Lactosylsphingosine	23
DPPE	44	GLC-110 mix	83	Lanosterol	74
DPPG	43	GLC-30 mix	81	Lauric acid	49
DPPI	45	GLC-40 mix	81, 82	LC, GM ₃ , GD ₃ qualitative mix	31, 84
DPPI-3,4,5-P3	47	GLC-50 mix	81, 82	LC	23
DPPI-3,4-P2	46	GLC-60 mix	81, 82	Lecithin	40, 90
DPPI-3P	45	GLC-70 mix	81, 82	L-erythro-2-Tetradecanoylamino-1-	
DPPI-4,5-P2	47	GLC-80 mix	81, 82	phenyl-1-propanol	35
DPPI-4-P	46	GLC-90 mix	81, 82	L-erythro-Dihydrosphingosine	4
DPPI-5-P	46	GLC-10 mix	81, 82	L-erythro-Sphinganine, C18 chain	4
DSPA	42	Globosides	25, 93	L-erythro-Sphingosine	2
DSPA	42	Globotetrahexosylceramide	25	L-erythro-Sphingosine, C18 chain	2
DSPC	43	Globotriaosylceramide	24	Lignoceric acid	52
DSPE	44	Glucocerebroside with C16:0-D ₃ fatty		Linoelaidic acid	54, 57
DSPE-MPEG-2000	44	acid side chain	25, 85	Linoleic acid	54
DSPG	44	Glucocerebroside with C22:0 fatty acid		Linolenic acid	54
D-threo-1-Phenyl-2-decanoylamino-3-		side chain	22	L-MAPP	35
morpholino-1-propanol•HCl	38	Glucocerebrosides	21, 91, 94	Loxastatin	39
D-threo-1-Phenyl-2-		Glucocerebrosides, plant	21, 94	L-threo-1-Phenyl-2-decanoylamino-3-	
hexadecanoylamino-3-morpholino-		Glucopsychosine	21, 22	morpholino-1-propanol•HCl	37
1-propanol•HCl	38	Glucosylceramide	21, 22	L-threo-1-Phenyl-2-hexadecanoylamino-	
D-threo-Dihydrosphingosine	4	Glucosylsphingosine	21, 22	3-morpholino-1-propanol•HCl	38
D-threo-PDMP	38	Glycosylceramides, qualitative mix		L-threo-Dihydrosphingosine	3, 35
D-threo-PPMP	38		31, 84	L-threo-PDMP	37
D-threo-Sphinganine, C18 chain	4	GM ₁	29, 91	L-threo-PPMP	38
D-threo-Sphingosine	2	GM ₂	29, 94	L-threo-SPC	17
D-threo-Sphingosine, C18 chain	2	GM ₃	29, 94	L-threo-Sphinganine, C18 chain	3, 35
		GM ₃ , GM ₂ , GM ₁ qualitative mix	31, 84	L-threo-Sphingosine	2
		GM ₄	29	L-threo-Sphingosine, C18 chain	2
		GQ _{1b}	30	L-threo-Sphingosylphosphorylcholine	17
		GT _{1b}	30, 92	lyso-Ceramide trihexoside	24
				lyso-Cerebroside	20
E		H		lyso-CTH	24
E-64-d	39	Heneicosanoic acid	51	lyso-Dihydrosphingomyelin	18
Eicosadienoic acid	55	Heptadecanoic acid	50	lyso-globotriosylsphingosine	24
Eicosanoic acid	51	Heptadecenoic acid	53	lyso-Glucocerebroside	21, 22
Eicosapentaenoic acid	56	Heptanoic acid	48	lyso-GM ₁	31
Eicosenoic acid	55	Hexacosanoic acid	52	lyso-Lactosylceramide	23
Elaidic acid	53, 57	Hexadecanoic acid	50	lyso-LC	23
EPA	56	Hydroxy fatty acid with long branched		lyso-Lecithin	40, 90
Ergosterol	74	chain	68, 71	lyso-Phosphatidylcholine	40, 90
EST	39			lyso-Monosialoganglioside GM ₁	31
Esterified steryl glucosides	75, 93			lyso-Phosphatidylcholine	43
				lyso-Sphingomyelin	17
				lyso-Sulfatide (NH ₄ ⁺ salt)	22
F				M	
FAME mixes, custom	80	I		Mead acid methyl ester	55
FIM-FAME-6 mix	75	iso-C13 methyl ester	69	Main phospholipid (MPL) of	
FIM-FAME-7 mix	76	iso-C14 methyl ester	69	<i>Thermoplasma acidophilum</i>	
Fluorescent sphingomyelin	18, 19	iso-C15 acid	69	(>95% pure)	47
Fluorescent LC	24, 26, 89	iso-C15 methyl ester	69	Main phospholipid (MPL) of	
Fucosylated mono-		iso-C16 methyl ester	69	<i>Thermoplasma acidophilum</i>	
sialoganglioside GM ₁	30	iso-C17 acid	69	(>50% pure)	47
Fucosyl-GM ₁	30	iso-C17 methyl ester	69	Margaric acid	50
Furan fatty acid	61	iso-C19 methyl ester	69	Methyl 10(E), 12(Z)-octadecadienoate	60
		iso-Heptadecanoic acid	69	Methyl 10-methylhexadecanoate	70
G		iso-Heptadecanoic methyl ester	69	Methyl 11-methyldodecanoate	69
Galactosylceramide, ceramide beta-D-		iso-Nonadecanoic methyl ester	69	Methyl 12-methyltetradecanoate	70
galactoside	20	iso-Palmitic methyl ester	69	Methyl 12-methyltridecanoate	69
Galactosylceramide with mostly 2-		iso-Pentadecanoic acid	69	Methyl 13-methylpentadecanoate	70
hydroxy fatty acid side chains	20	iso-Pentadecanoic methyl ester	69	Methyl 13-methyltetradecanoate	69
Galactosylceramide with mostly non-		iso-Tetradecanoic methyl ester	69	Methyl 14-methylhexadecanoate	70
hydroxy fatty acid side chain	20	iso-Tridecanoic methyl ester	69	Methyl 14-methylpentadecanoate	69
gamma-Linolenic acid	54				
Gangliotetraosylceramide	28, 92	K			
Gangliotetraosylceramide and sialosyl		KEL-FIM-FAME-5 mix	75		
derivatives mix	32, 85				
Gangliotriosylceramide	28	L			
Gb3	24	Lactocerebrosides	23		
Gb4	25				

Methyl 15-hydroxypentadecanoate	66	Methyl heptadecenoate	53	N-Acetyl-sphingosyl-	
Methyl 15-methylhexadecanoate	69	Methyl heptanoate	48	phosphorylethanolamine	17
Methyl 17-hydroxyheptadecanoate	67	Methyl hexacosanoate	52	N-Acetyl-sulfatide	22
Methyl 17-methyloctadecanoate	69	Methyl hexadecanoate	50	N-C2:0 Ceramide of D-erythro-C14-	
Methyl 20-hydroxyeicosanoate	67	Methyl hexanoate	48	sphingosine	10
Methyl 21-hydroxyheneicosanoate	67	Methyl homogamma linolenate	55	N-C2:0-Cerebroside	20
Methyl 22-hydroxydocosanoate	67	Methyl laurate	49	N-C2:0-D-erythro-Ceramide	7
Methyl 27-hydroxyheptacosanoate	67	Methyl lignocerate	52	N-C2:0-D-erythro-Dihydroceramide	11
Methyl 2-fluoropalmitate	39	Methyl linoelaidate	54, 57	N-C2:0-L-erythro-Ceramide	8
Methyl 2-hydroxydecanoate	62	Methyl linoleate	54	N-C2:0-L-threo-Ceramide	8
Methyl 2-hydroxydocosanoate	63	Methyl linolenate	54	N-C2:0-Phytoceramide	13
Methyl 2-hydroxydodecanoate	62	Methyl margarate	50	N-C2:0-sulfatide	22
Methyl 2-hydroxyeicosanoate	63	Methyl myristate	50	N-C6:0 Ceramide of D-erythro-C8-	
Methyl 2-hydroxyhexadecanoate	62	Methyl myristoleate	52	sphingosine	10
Methyl 2-hydroxyoctadecanoate	62	Methyl nervonate	57	N-C6:0-D-erythro-Ceramide	8
Methyl 2-hydroxytetraacosanoate	63	Methyl nonadecanoate	51	N-C6:0-D-erythro-Dihydroceramide	11
Methyl 2-hydroxytetradecanoate	62	Methyl nonadecenoate	55	N-C6:0-D-threo-Ceramide	8
Methyl 2-hydroxytricosanoate	63	Methyl nonanoate	49	N-C6:0-L-erythro-Ceramide	8
Methyl 30-hydroxytriacontanoate	67	Methyl octadecanoate	50	N-C6:0-L-threo-Ceramide	8
Methyl 3-hydroxydecanoate	64	Methyl octanoate	49	N-C6:0-NBD-beta-D-galactosyl	
Methyl 3-hydroxydodecanoate	65	Methyl oleate	53	sphingosine-	21, 26, 88
Methyl 3-hydroxyheptadecanoate	66	Methyl palmitate	50	N-C6:0-NBD-beta-D-glucosyl-	
Methyl 3-hydroxyhexadecanoate	65	Methyl palmitelaidate	53, 57	sphingosine	21, 26, 88
Methyl 3-hydroxyhexanoate	64	Methyl palmitoleate	52	N-C6:0-NBD-beta-D-lactosyl-	
Methyl 3-hydroxynonanoate	64	Methyl pentadecanoate	50	sphingosine	24, 26, 88
Methyl 3-hydroxyoctadecanoate	66	Methyl ricinelaidate	68	N-C6:0-NBD-ceramide	14, 86
Methyl 3-hydroxyoctanoate	64	Methyl stearate	50	N-C6:0-NBD-cerebroside	21, 26, 88
Methyl 3-hydroxytetradecanoate	65	Methyl tetraacosanoate	52	N-C6:0-NBD-D-erythro-	
Methyl 3-hydroxytridecanoate	65	Methyl tetradecanoate	50	dihydrosphingosine	15, 87
Methyl 3-hydroxyundecanoate	65	Methyl threo-2,3-dihydroxypalmitate	68	N-C6:0-NBD-D-erythro-sphingosine	
Methyl 5,8,11-eicosatrienoate	55	Methyl trans 11-octadecenoate	53, 57		14, 86
Methyl 8-(5-hexyl-2-furyl)-octanoate	61	Methyl trans vaccenate	53, 57	N-C6:0-NBD-dihydroceramide	14, 87
Methyl 9(E),11(E)-octadecadienoate	59	Methyl tricosanoate	51	N-C6:0-NBD-galactosylceramide	
Methyl 9(Z), 11(E)-octadecadienoate	59, 60	Methyl tridecanoate	49		21, 26, 88
Methyl 9(Z), 11(Z)-octadecadienoate	60	Methyl undecanoate	49	N-C6:0-NBD-glucosylceramide	21, 26, 88
Methyl arachidate	51	MGDG (hydrogenated, distearoyl)	48, 91	N-C6:0-NBD-L-threo-	
Methyl arachidonate	55	Mixed gangliosides	31, 92	dihydrosphingosine	14, 86
Methyl behenate	51	Monoclonal antibody to GD ₃	32	N-C6:0-NBD-L-threo-sphingosine	14, 86
Methyl caprate	49	Monogalactosyldiglyceride	48, 91	N-C6:0-NBD-lactosylceramide	24, 26, 88
Methyl caproate	48	Monosialoganglioside GM ₁	29, 91	N-C6:0-NBD-phytoceramide	15, 87
Methyl caprylate	49	Monosialoganglioside GM ₂	29, 94	N-C6:0-NBD-phytosphingosine	15, 87
Methyl cerotate	52	Monosialoganglioside GM ₃	29, 94	N-C6:0-NBD-sphingomyelin	18, 87
Methyl cis-11-octadecenoate	54	Monosialoganglioside GM ₄	29, 93	N-C6:0-NBD-sphingosyl-	
Methyl cis-9,10-		Monosialoganglioside mix	31, 84	phosphorylcholine	18, 87
Methyleneoctadecanoate	71	MPL of <i>Thermoplasma acidophilum</i>		N-C6:0-Phytoceramide	13
Methyl cis-vaccenate	54	(>50% pure)	47	N-C8:0-CPPC	36
Methyl D,L-erythro-corynomycolate	68	Myristic acid	50	N-C8:0-Cyclopropenylceramide	36
Methyl D,L-threo-corynomycolate	68	Myristoleic acid	52	N-C8:0-D-erythro-Ceramide	8
Methyl decanoate	49	N		N-C8:0-D-erythro-Dihydroceramide	11
Methyl dihydrosterculate	71	N-[(1R, 2S)-2-hydroxy-1-		N-C8:0-D-threo-Ceramide	9
Methyl docosahexaenoate	56	hydroxymethyl-2-(2-tridecyl-1-		N-C8:0-Galactosylceramide	20
Methyl docosanoate	51	cylopropenyl) ethyl] hexadecamide	36	N-C8:0-L-threo-Ceramide	9
Methyl docosenoate	56	N-[(1R, 2S)-2-hydroxy-1-		N-C8:0-Phytoceramide	13
Methyl docosapentaenoate	56	hydroxymethyl-2-(2-tridecyl-1-		N-C10:0-D-erythro-Ceramide	9
Methyl dodecanoate	49	cylopropenyl) ethyl] octanamide	36	N-C12:0-NBD-beta-D-lactosyl-	
Methyl eicosadienoate	55	N-1- ¹³ C-Palmitoyl-		sphingosine	24, 26, 88
Methyl eicosanoate	51	spingosylphosphorylcholine	17, 85	N-C12:0-NBD-ceramide	14, 86, 88
Methyl eicosapentaenoate	56	N,N-Dihexyl-D-erythro-sphingosine	6	N-C12:0-NBD-CTH	25, 27, 88
Methyl eicosenoate	55	N,N-Dimethyl-D-erythro-sphingosine	6	N-C12:0-NBD-D-erythro-	
Methyl elaidate	53, 57	N-Acetyl-D-erythro-dihydrosphingosine	11	dihydrosphingosine	15, 87
Methyl erucate	56	N-Acetyl-D-erythro-sphinganine	11	N-C12:0-NBD-dihydroceramide	14, 15, 87
Methyl ester of CLA (10-trans, 12-cis)	60	N-Acetyl-D-erythro-sphingosine	7	N-C12:0-NBD-D-erythro-sphingosine	
Methyl ester of CLA (9-cis, 11-cis)	60	N-Acetyl-D-erythro-sphingosine (C14			14, 86, 87
Methyl ester of CLA (9-cis, 11-trans)	59, 60	sphingoid base)	10	N-C12:0-NBD-globotriaosylceramide	
Methyl ester of CLA(9-trans, 11-trans)	59	N-Acetyl-L-erythro-sphingosine	8		25, 27, 88
Methyl ester of furan fatty acid	61	N-Acetyl-L-threo-sphingosine	8	N-C12:0-NBD-L-threo-	
Methyl ester of omega-3 fatty acid	56, 57	N-Acetyl-phytosphingosine	13	dihydrosphingosine	14, 87
Methyl gamma-linolenate	54	N-Acetyl psychosine	20	N-C12:0-NBD-L-threo-sphingosine	14, 86
Methyl heneicosanoate	51	N-Acetyl sphingosine	20	N-C12:0-NBD-lactosylceramide	24, 26, 88
Methyl heptadecanoate	50	N-Acetyl-sphingosyl-beta-D-		N-C12:0-NBD-phytoceramide	15, 87
		galactoside-3-sulfatide	22	N-C12:0-NBD-phytosphingosine	15, 87
		N-Acetyl-sphingosylphosphorylcholine	16	N-C12:0-NBD-sphingomyelin	19, 87

N-C12:0-NBD-sphingosyl-phosphorylcholine	19, 87	N-Hexanoyl-D-erythro-dihydrosphingosine	11	N-(NBD-aminolauroyl) sulfatide	27, 88
N-C12:0-NBD-sulfatide	23, 27, 88	N-Hexanoyl-D-erythro-sphinganine	11	N-Nonadecanoyl-D-erythro-sphingosine	9
N-C15:0-Cerebroside	20	N-Hexanoyl-D-erythro-sphingosine	8	N-Octadecanoyl-D-erythro-dihydrosphingosine	11
N-C15:0-D-erythro-Ceramide	9	N-Hexanoyl-D-erythro-sphingosine (C8 sphingoid base)	10	N-Octadecanoyl-D-erythro-sphinganine	11
N-C16:0-D-erythro-Ceramide	9	N-Hexanoyl-D-threo-sphingosine	8	N-Octadecanoyl-D-erythro-sphingosine	9
N-C16:0-Ceramide-1-phosphate	18	N-Hexanoyl-L-erythro-sphingosine	8	N-Octadecanoyl-D-threo-sphingosine	10
N-C16:0-CPPC	36	N-Hexanoyl-L-threo-sphingosine	8	N-Octadecanoyl-D ₃ -ceramide trihexoside	25, 26, 86
N-C16:0-Cyclopropenylceramide	36	N-Hexanoyl-NBD-beta-D-lactosyl-sphingosine	24, 26, 88	N-Octadecanoyl-D ₃ -globotriaosylceramide	25, 26, 86
N-C16:0-D ₃ -Glucosylpsychosine	25, 85	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine	15, 87	N-Octadecanoyl-D ₃ -monosialoganglioside GM ₁	29, 85
N-C16:0-D ₃ -Lactosylceramide	26, 85	N-Hexanoyl-NBD-D-erythro-sphingosine	14, 86	N-Octadecanoyl-D ₃ -sulfatide	23, 26, 85
N-C16:0-D-erythro-Ceramide	9	N-Hexanoyl-NBD-galactosylceramide	21, 26, 88	N-Octadecanoyl-L-erythro-sphingosine	10
N-C16:0-Phytoceramide	13	N-Hexanoyl-NBD-glucoylceramide	21, 26, 88	N-Octadecanoyl-sphingosylphosphorylcholine	16
N-C17:0-Ceramide trihexoside	24	N-Hexanoyl-NBD-L-threo-dihydrosphingosine	14, 86	N-Octanoyl-β-D-galactosylceramide	20
N-C17:0-D-erythro-Ceramide	9	N-Hexanoyl-NBD-L-threo-sphingosine	14, 86	N-Octanoyl-D-erythro-dihydrosphingosine	11
N-C18:0-D-erythro-Ceramide	9	N-Hexanoyl-NBD-lactosylceramide	24, 26, 88	N-Octanoyl-D-erythro-sphinganine	11
N-C18:0-D-erythro-Dihydroceramide	11	N-Hexanoyl-NBD-phytosphingosine	15, 87	N-Octanoyl-D-erythro-sphingosine	8
N-C18:0-D-threo-Ceramide	10	N-Hexanoyl-NBD-sphingosyl-phosphorylcholine	18, 87	N-Octanoyl-D-threo-sphingosine	9
N-C18:0-D ₃ -Sulfatide	23, 26, 85	N-Hexanoyl-phytosphingosine	13	N-Octanoyl-L-threo-sphingosine	9
N-C18:0-L-erythro-Ceramide	10	N-Hexanoyl-sphingosylphosphorylcholine	16	N-Octanoyl-phytosphingosine	13
N-C18:0-L-threo-Ceramide	10	N-(NBD-aminocaproyl)-beta-D-galactosylsphingosine	26, 88	NOE	34, 71
N-C18:0-Phytoceramide	13	N-(NBD-aminocaproyl)-beta-D-glucoylsphingosine	26, 88	N-Oleylethanolamine	34, 71
N-C19:0-D-erythro-Ceramide	9	N-(NBD-aminocaproyl)-beta-D-lactosylsphingosine	26, 88	Nonadecanoic acid	51
N-C23:0-Ceramide trihexoside	25	N-(NBD-aminocaproyl)-D-erythro-dihydrosphingosine	15, 87	Nonadecanoic acid	55
N-C24:0-D-erythro-Ceramide	10	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	Nonanoic acid	49
N-C24:0-Phytoceramide	13	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	Non-polar lipid mix A	84
N-C24:0-Sulfatide	23	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	Non-polar lipid mix B	84
N-D ₃ -Stearoyl-GM ₁	29, 85	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	Non-volatile acid mix	83
N-Decanoyl-D-erythro-sphingosine	9	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl serinol	6
N-Docosanoyl-β-glucosylsphingosine	22	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl-D ₃ -glucosylpsychosine	25, 85
N-Docosanoyl-D-erythro-sphingosylphosphorylcholine	17	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl-D ₃ -lactosylceramide	26, 85
N-Docosanoyl-glucoylpsychosine	22	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl-lactosylceramide	23
N-Dodecanoyl-NBD-beta-D-lactosyl-sphingosine	24, 26, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl-sphingosyl-beta-D-galatoside-3-sulfate	22
N-Dodecanoyl-NBD-ceramide trihexoside	25, 27, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Palmitoyl-sulfatide	22
N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine	15, 87	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Pentadecanoyl-D-erythro-sphingosine	9
N-Dodecanoyl-NBD-D-erythro-sphingosine	14, 86	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Pentadecanoyl-psychosine	20
N-Dodecanoyl-NBD-L-threo-dihydrosphingosine	14, 87	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-ceramide	10
N-Dodecanoyl-NBD-L-threo-sphingosine	14, 86	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-dihydroceramide	11
N-Dodecanoyl-NBD-lactosylceramide	24, 26, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxy-C16:0-D-erythro-dihydrosphingosine	12
N-Dodecanoyl-NBD-lyso-sulfatide	23, 27, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-ceramide	10
N-Dodecanoyl-NBD-phytosphingosine	15, 87	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-dihydroceramide	11
N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate	23, 27, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-dihydrosphingosine	11
N-Dodecanoyl-NBD-sphingosyl-phosphorylcholine	19, 87	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-sphingosine	10
N-Dodecanoyl-NBD-sulfatide	23, 27, 88	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxyhexadecanoyl-D-erythro-dihydrosphingosine	12
N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine	16	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-dihydrosphingosine	11
Nervonic acid	56	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-sphingosine	10
Neutral glycosphingolipid qualmix	31, 84	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Stearoyl-D ₃₅ -psychosine, perdeuterated	25, 85
N-Heptadecanoyl ceramide trihexoside	24	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Stearoyl-phytosphingosine	13
N-Heptadecanoyl-D-erythro-sphingosine	9	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Tetracosanoyl-D-erythro-sphingosine	10
N-Heptadecanoyl globotriaosylceramide	24	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Tetracosanoyl-phytosphingosine	13
N-Heptadecanoyl-sphingosylphosphorylcholine	16	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Tetracosanoyl-sphingosyl-beta-D-galactoside-3-sulfate	23
N-Hexadecanoyl-D-erythro-sphingosine	9	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86	N-Tetracosanoyl-sulfatide	23
N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate	18	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86		
N-Hexadecanoyl-phytosphingosine	13	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86		
N-Hexadecanoylethanolamine	34, 71	N-(NBD-aminocaproyl)-D-erythro-sphingosine	14, 86		

N-Tricosanoyl ceramide trihexoside	25
N-Tricosanoyl globotriaosylceramide	25
O	
Octadecadienoic acid-10(E),12(Z)	60
Octadecadienoic acid-11(Z), 13(E)	60
Octadecadienoic acid-9(E),11(E)	59
Octadecadienoic acid-9(Z),11(E)	59
Octadecadienoic acid-9(Z),11(Z)	60
Octadecanoic acid	50
Octanoic acid	48
Oleic acid	53
omega 3 fatty acid	56
omega-hydroxy C10:1 (2-trans)	66
omega-Hydroxy C15:0	66
omega-Hydroxy C15:0 fatty acid methyl ester	66
omega-Hydroxy C17:0 fatty acid	66
omega-Hydroxy C17:0 fatty acid methyl ester	67
omega-Hydroxy C20:0 fatty acid	67
omega-Hydroxy C20:0 fatty acid methyl ester	67
omega-Hydroxy C21:0 fatty acid	67
omega-Hydroxy C21:0 fatty acid methyl ester	67
omega-hydroxy C22:0 fatty acid	67
omega-Hydroxy C22:0 fatty acid methyl ester	67
omega-Hydroxy C27:0 fatty acid methyl ester	67
omega-Hydroxy C30:0 fatty acid methyl ester	67
P	
PA	41
Palmitelaic acid	52, 57
Palmitic acid	50
Palmitoleic acid	52
Palmitoyl sulfatide	22
Palmitoyl serinol	6
Palmitoyl lactosylceramide	23
PC	40
PDMP	37, 38
PE	41
Pelargonic acid	49
Pentadecanoic acid	50
Phellonic acid	67
Phosphatidic acid	41, 91
Phosphatidylcholine	40
Phosphatidylethanolamine	41, 90
Phosphatidylinositol	41, 90
Phosphatidylinositol 3-phosphate dipalmitoyl	45
Phosphatidylinositol 4-phosphate dioctanoyl	46
Phosphatidylinositol 4-phosphate dipalmitoyl	46
Phosphatidylinositol 5-phosphate dioctanoyl	46
Phosphatidylinositol 5-phosphate dipalmitoyl	46
Phosphatidylinositol-bis-3,4-phosphate dipalmitoyl	46
Phosphatidylinositol-bis-4,5-phosphate dioctanoyl	46
Phosphatidylinositol-bis-4,5-phosphate dipalmitoyl	46, 47
Phosphatidylinositol, dipalmitoyl	45

Phosphatidylinositol-tris-3,4,5-phosphate dipalmitoyl	47
Phosphatidylinositol-tris-3,4,5-phosphate dioctanoyl	47
Phosphatidylserine	40, 90
Phosphoglycerides kit	41
Phrenosin	20, 92
Phytanic acid	71
Phytosphingosine	5
PI	41, 45
PI-3,4,5-P3, dioctanoyl	47
PI-3,4,5-P3, dipalmitoyl	47
PI-3,4-P2 dipalmitoyl	46
PI-3-P dipalmitoyl	45
PI-4,5-P2 dioctanoyl	46
PI-4,5-P2 dipalmitoyl	47
PI-4-P dioctanoyl	46
PI-4-P dipalmitoyl	46
PI-5-P dioctanoyl	46
PI-5-P dipalmitoyl	46
Plant sterol mix	74
Plant sterols kit	74
Polar lipid mix	83
Polyclonal antibody to asialo-GM ₁	32
Polyclonal antibody to asialo-GM ₂	32
Polyclonal antibody to GL-4	33
Polyclonal antibody to GM ₁	32
Polyclonal antibody to GM ₂ (NANA)	33
Polyclonal antibody to GM ₂ (NGNA)	33
Polyclonal antibody to GM ₄	33
POPC	43
POPG	44
PPMP	37, 38
PS	40, 90
Psychosine	20
PUFA-1	76
PUFA-2	76
PUFA-3	76
Purified mixed gangliosides	31, 92
Purified MPL of <i>Thermoplasma acidophilum</i> (>95% pure)	47
Q	
Qualitative mix, bacterial lipid standard	83
Qualitative mix, cis-trans isomers	58, 78
Qualitative mix, disialogangliosides	84
Qualitative mix, gangliotetraosyl ceramide and sialosyl derivatives	85
Qualitative mix, glycosyl-ceramides	84
Qualitative mix, lactosylceramide and sialosyl derivatives	84
Qualitative mix, monosialogangliosides	84
Qualitative mix, non-polar lipids	84
Qualitative mix, non-volatile acids	83
Qualitative mix, polar lipids	83
Qualitative mix, PUFA	76
Qualitative mix, sphingolipids	84
Qualitative mix, TLC standards	84
Qualitative mix, volatile acids	83
Qualitative mix, water soluble fatty acids	82
Quantitative mix, bacterial fatty acid methyl esters	83
Quantitative mix, carbohydrates	76, 77
Quantitative mix, GC	81, 82
Quantitative mix, hydroxy methyl esters	63
Quantitative mix, methyl esters	75, 76

R	
rac-5,7-Dimethyltocol	73
rac-alpha-Tocopherol	72
rac-beta-Tocopherol	72
rac-gamma-Tocopherol	72
rac-Tocol	73
Rapeseed oil reference mixture	78
Ricinelaic acid	68
RM-1 mix	78, 80
RM-2 mix	79, 80
RM-3 mix	79, 80
RM-4 mix	79, 80
RM-5 mix	79, 80
RM-6 mix	79, 80
RM-7 kit	79
Royal Jelly acid	66
S	
S-1-P	18
Safingol	3, 35
Sapienic acid	52
S-P-A	18
SPC	17
Sphingolipid mix	84
Sphingomyelin 15, 16, 91, 93, 94	16
Sphingomyelin, C17:0 fatty acid	16
Sphingomyelin, C18:0 fatty acid	16
Sphingomyelin, C2:0 fatty acid	16
Sphingomyelin, C20:0 fatty acid	16
Sphingomyelin, C22:0 fatty acid	17
Sphingomyelin, C6:0 fatty acid	16
Sphingosine	2
Sphingosine-1-galactoside-3-sulfate	22
Sphingosine with C10 chain	3
Sphingosine with C12 chain	3
Sphingosine with C14 chain	2
Sphingosine with C16 chain	3
Sphingosine with C20 chain	3
Sphingosine with C18 chain	2
Sphingosine, with tertiary amine group	6
Sphingosine, D-erythro	2
Sphingosine, D-threo	2
Sphingosine, L-erythro	2
Sphingosine, L-threo	2
Sphingosylphosphorylcholine	17
Sphingosylphosphorylethanolamine with C2 fatty acid side chain	17
SPM	15, 16
SPM with ¹³ C labeled fatty acid	17, 85
Stearic acid	50
Sterol mixture, plant	74
Sterols kit	74
Steryl glucosides	75
Stigmastanol	74
Stigmasterol	74
Sulfatide with C16:0 fatty acid side chain	22
Sulfatides	22, 92
T	
Tetracosanoic acid	52
Tetracosanoyl sulfatide	23
Tetradecanoic acid	50
Tetramethylhexadecanoic acid-3,7,11,15	71
Tetrasialoganglioside GQ _{1b}	30, 93
THI	39
TLC standards mix	83, 84
Tocol	73
trans 11-Octadecenoic acid	53, 57
trans vaccenic acid	53, 57

Tricosanoic acid	51
Tridecanoic acid	49
Trimethyltocol	72
Trisialoganglioside GT _{1b}	30, 92

U

Undecanoic acid	49
-----------------	----

V

Volatile acid mix	83
-------------------	----

W

Water soluble fatty acid qualitative mix	82
WSFA-2 mix	82
WSFA-4 mix	82

MATREYA LLC

International Dealers & Representatives

Australia (South Australia)

Adelab Scientific
36 Holland Street
Thebarton 5031
Tel: 61-88-234-7955
Fax: 61-88-234-7897
Email: info@adelab.com.au
Web: www.adelab.com.au

Canada

MJS Biolynx
PO Box 1150
300 Laurier Blvd
Brockville, Ontario K6V 5W1
Tel: 613-498-2126
Toll free: 888-593-5969
Fax: 613 342 1341
Email: sales@biolynx.ca
Web: www.biolynx.ca

France

Biovalley
18 Route de Tournan
Marne la Vallée Cedex 3
Conches, 77601
Tel: 33-16-007-2020
Fax: 33-16-007-5051
Email: biovalley@biovalley.fr
Web: www.biovalley.mgn.fr

Italy

Superchrom S.r.l.
Via C. Menotti, 11
Milano, 20129
Tel: 39-02-738-6315
Fax: 39-027-010-0100
Email: Superchrom@iol.it

Korea

Kim & Friends, Inc.
3Fl., 1019-16 Sinjeong-dong
Yangcheon-gu
Seoul 158-861
Tel: 82-2-2647-6611
Fax: 82-2-2647-6687
Email: kslee@kimnfriends.co.kr
Web: www.kimnfriends.co.kr

New Zealand

Phenomenex NZ Ltd.
PO Box 31-6-1
Milford, Auckland
Tel: 649-478-0951
Fax: 649-478-0952
Email: info@phenomenex.co.za
Web: www.phenomenex.com

Belgium

SerCoLab BV/BA
West Kaai 7
Merksem, B-2170
Tel: 323-640-3315
Fax: 323-644-0405
Email: info@sercolab.be
Web: www.sercolab.be

China

Shenzhen Bolin Chemical Co. LTD
Rm 1309, No. 1 News Bldg
2 Shennan Zhong Road
Shenzhen, 518027
Tel: 867-556-165-1418 (1428)
Fax: 867-558-209-6552
Email: lh@bolin-lkmlabs.com
Web: www.bolin-lkmlabs.com

Germany

BIOTREND Chemikalien GmbH
Eupenerstr. 157
Köln, 50933
Tel: 49-221-949-8320
Fax: 49-221-949-8325
Email: jaeger@biotrend.com
Web: www.biotrend.com

Japan

Techno Chemical Corp
27-9, Honkomagome 1 Chome
Bunkyo-ku, Tokyo 113
Tel: 81-33-947-7310
Fax: 81-33-947-7306
Email: info@technochemical.com
Web: www.technochemical.com

Netherlands

Distrilab BV
Olmenlaan 6-C
Leusden, 3833 AV
Tel: 31-33-494-7834
Fax: 31-33-432-1441
Email: info@distrilab.nl
Web: www.distrilab.nl

South Africa

Anatech Instruments (PTY) Ltd
PO Box 98485, Sloane Park
Gauteng, 2152
Tel: 27-11-462-6776
Fax: 27-11-704-6490
Email: sales@anatach.co.za

Spain

Teknokroma S.C.C.L.
Camí de Can Calders, 14
Apartado de Correos, 147
Sant Cugat del Vallès
Barcelona 08190
Tel: 34-93-674-8800
Fax: 34-93-675-2405
Email: commercial@teknokroma.es
Web: www.teknokroma.es

Switzerland

VWR International AG-Life Sciences
Lerzenstrasse 16/18
Dietikon, 8953
Tel: 41-44-745-1437
Fax: 41-44-745-1410
Email: Francesca.sagona@ch.vwr.com
Web: www.juro.ch

United Kingdom (England)

Universal Biologicals Ltd.
Passhouse Farmhouse
Papworth St. Agnes
Cambridge CB3 8QU
Tel: 44-148-083-9015
Fax: 44-148-803-1912
Email: info@universalbiologicals.ltd.uk
Web: www.universalbiologicals.ltd.uk

Sweden

Larodan Fine Chemicals AB
Limhamnsgårdens Alle'9
Malmö, S-21616
Tel: 464-016-4155
Fax: 464-015-5498
Email: info@larodan.se
Web: www.larodan.se

Taiwan

Super Chroma Enterprise Ltd
5F.226 Roosevelt Rd, Section 5
Taipei ROC, 116
Tel: 88-622-930-7110
Fax: 88-622-930-7112
Email: sc@superchroma.com.tw
Web: www.superchroma.com.tw

Matreya LLC Ordering Information

Office Hours: Monday - Friday, 8:00am - 4:00pm, Eastern Time

• Ordering and Customer Service

Telephone (Toll Free)
Telephone (worldwide)
Fax (24/7 worldwide)
e-mail
e-commerce website

(800) 342 3595
(814) 359 5060
(814) 359 5062
customerservice@matreya.com
www.matreya.com

• Technical Service

Telephone (worldwide)
Fax (worldwide)
e-mail

(814) 359 5060
(814) 359 5062
techservice@matreya.com

• Terms

Prices and discounts are subject to change without notice. Freight charges are prepaid and added as a separate charge on the invoice. Orders placed are F.O.B. Pleasant Gap, PA. **Net 30 Days.**

• Shipping

Standard orders: FedEx 2nd-Day
Rush Orders: FedEx Priority or Standard overnight
International Orders: FedEx International Priority
Items requiring dry ice for shipping will be charged a \$40 fee.

• Return Policy

All returns must have prior authorization. Items incorrectly ordered by you and returned to Matreya are subject to a 25% (\$25 minimum) restocking charge. The following items may not be returned: frozen products, items which have passed their expiration dates, custom synthesized products and accommodation orders. If a product has been incorrectly sent to you due to an error on our part, a credit will be issued to your account immediately. Please inspect and verify your order upon receipt. No products will be returned after 30 days (assuming proper storage and handling by customer).

• Disclaimer

The information and data included in this catalog are correct and reliable to the best of our knowledge. However, we offer no guarantees and assume no responsibility for this information. No license or immunity under any patent is granted or implied through our sale of any material. Any information sent to us by the customer, relative to his or her interest in Matreya's products, must be submitted voluntarily and without obligation on the part of Matreya.

Matreya LLC
168 Tressler Street
Pleasant Gap, PA 16823 USA
Federal ID No. 20-1237500

