

MATREYA



Lipids, Biochemicals,
and Standards for
Life Science Research

2013 - 2014

About Matreya LLC

- Matreya strives to develop, manufacture and deliver products of the highest value to our customers.
- Quality will always be the best achievable by state-of-the-art techniques, typically greater than 98%
- We strive for rapid delivery. 95+% of our products are shipped within 24 hours of receipt of an order.
- Within the area of sphingolipids and glycolipids, we have earned a reputation as the preferred problem solver and technology leader.
- When you demand quality and consistency, you may rely on Matreya lipids.

Matreya Products for Biochemistry Research.

We offer one of the widest selection of ceramides for intracellular signaling research available. We stock antibodies to glycosphingolipids as well as inhibitors of enzymes involved in glycosphingolipid metabolism.

Our products provide the valuable tools for the study of cell membrane and its structure, growth regulators in the cellular metabolism, and intracellular mediators.

We are able to make our products better and better with the latest technology in Chromatography, Mass Spectrometry, and NMR techniques.

We are proud to offer our products as a valuable tool for your life science research needs.

Matreya Products for Microbiology Research.

Matreya stocks many unusual fatty acid standards produced by bacteria that are useful for culture characterization.

Matreya Products for the Food and Agriculture Industries.

Many of Matreya's fatty acid products have been industry standards for many years. The acids and their methyl esters are used as standards in analysis and quality control.

Custom Preparations.

Our experience in chemical synthesis and the extraction and purification of natural products allows us to produce custom preparations with the same high quality and purity as the products listed in the catalog. Depending on the complexity of the molecule, delivery will be 4 to 12 weeks after receipt of an order, usually less than 6 weeks.

If you can't find a product in the catalog, please check the INDEX, where we also try to list common synonyms for our products.

TABLE OF CONTENTS

TABLE OF CONTENTS	i
Technical Service.....	iii
Natural Products.....	iii
Storage.....	iii
Sphingolipid Structures and Pathways.....	iii
Package Weight.....	iii
Matreya's Mission.....	iii
Sphingoid Bases, Sphingolipids and Glycosphingolipids	1
Sphingosines.....	2
Synthetic Sphingosines with C18 Sphingoid Base	2
Synthetic Sphingosines with Sphingoid Bases other than C18.....	2
Synthetic Dihydrosphingosines.....	3
3-Keto-Dihydrosphingosines.....	4
Phytosphingosines.....	5
Other Sphingosine Derivatives and Precursors.....	5
Ceramides	6
Synthetic Ceramides Derived from C18-Sphingosine	7
2-Hydroxy Ceramides	9
Ceramide Made from Sphingosines with Sphingoid Bases Other Than C18.....	10
Dihydroceramides	10
2-Hydroxy Dihydroceramides	11
Ceramides From Natural Sources	11
Phytoceramides	12
Fluorescent Ceramides.....	13
Phosphosphingolipids	15
Sphingomyelins.....	15
Sphingosylphosphorylcholines (SPC).....	17
Sphingosine Phosphates	17
Fluorescent Sphingomyelins.....	18
Glycosphingolipids	19
Galactosylceramides and Glucosylceramides.....	19
Lactosylceramides	24
Ceramide Trihexosides	25
Globosides	26
Stable Isotopes Labeled Glycolipids.....	27
Fluorescent Compounds.....	27
Gangliosides	29
Glycosphingolipid Reference Mixtures for TLC	32
Antibodies Directed Against Glycolipids	33
Enzyme Inhibitors	35
Glycerolipids	40
Glycerophospholipids.....	40
Natural Phospholipids.....	40
Synthetic Phospholipids.....	42
Phosphatidylcholines	42
Phosphatidylglycerols	44
Phosphatidylethanolamines.....	45
Phosphatidylinositols	46
Bacterial Tetraethers.....	47
Glycosyl Glycerides.....	48
Fatty Acids	48

Simple Fatty Acids.....	48
Saturated Fatty Acids and Methyl Esters.....	48
Unsaturated Fatty Acids and Methyl Esters.....	53
Trans Fatty Acids and Methyl Esters.....	59
Conjugated Linoleic Acid Isomers (CLA).....	61
Other CLA Products and Derivatives.....	63
Hydroxy Fatty Acids.....	63
2-Hydroxy Fatty Acids and Methyl Esters.....	63
3-Hydroxy Fatty Acids and Methyl Esters.....	66
Omega Hydroxy Fatty Acids.....	68
Other Hydroxy Fatty Acids.....	70
Branched and Cyclic Fatty Acids.....	70
Iso-Fatty Acids and Esters.....	70
Anteiso-Fatty Acids and Esters.....	72
Methylated Fatty Acids.....	72
Cyclopropyl Fatty Acids and Esters.....	73
Unusual Fatty Acids and Derivatives.....	73
Other Lipids.....	74
Tocopherols.....	74
Cholestane Derivatives.....	76
Plant Sterols and Steryl Glucosides.....	76
Standards and Reference Compounds.....	78
Food Industry Mixtures.....	78
Each methyl ester mixture is carefully prepared by weight.....	78
Polyunsaturated Fatty Acid Methyl Esters Mixtures.....	79
Carbohydrate Mixtures.....	79
Other Fatty Acid Methyl Ester Mixtures.....	80
AOCS Animal and Vegetable Oil Reference Mixtures (RM Mixtures).....	81
Custom Mixtures.....	83
GLC Standard Mixtures.....	84
Water Soluble Fatty Acid Mixtures.....	85
Microbiology Standard Mixtures.....	86
Biochemical Research Standard Mixtures.....	87
Glycosphingolipid Reference Mixtures for TLC.....	87
Biochemicals and Reagents.....	88
Stable Isotope Labeled Compounds.....	88
Fluorescent Compounds.....	89
Sphingolipid Structures and Pathways Wall Chart.....	98
Literature References.....	99
Cross Reference for Product Numbers and Catalog Pages.....	100
Product Name Index.....	103
Tables	
Table I. AOCS Oil Reference Mixes.....	81
Table II. Standards for GC Analysis.....	84
Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC.....	93

All chemicals listed in this catalog are for investigational use only. Chemicals are not for human consumption or to be used in food or food additives. Matreya assumes no liability for any use of these chemicals by the end user. We believe the information in this catalog, offered in good faith, is accurate.

Limited Warranty: All Matreya Products, except those specifically exempted, are warranted (for 30 days) to be free of defects in materials and workmanship, if properly stored. Any replacements required as a result of such defects will be made without charge provided that such defective products are returned with a written explanation. Please request a Returned Goods Authorization before returning products under this warranty.

Technical Service

Our technical service department may be contacted by telephone at 800.342.3595, or by e-mail at techservice@matreya.com.

Natural Products

Some of our glycolipids are extracted from natural sources. These products have a normal heterogeneity in their lipid components, particularly in the fatty acids. Variations include carbon chain length as well as the presence or absence of 2-hydroxy fatty acids. Products based on sphingosine may contain longer chain sphingoid bases as well as chains with multiple double bonds. This heterogeneity may result in additional spots showing on TLC plates or multiple peaks in LC analyses. We have listed the typical fatty acid compositions of our natural products in the appendix.

Storage

Catalog items in unopened containers are stable for at least one year when stored under the conditions indicated in the catalog listing. Items containing unsaturated fatty acids are subject to oxidation and should be stored in a solution of organic solvents or under argon. Glycolipids and phospholipids should not be stored in aqueous solutions due to potential hydrolysis.

Sphingolipid Structures and Pathways

In a clear and straightforward manner, this wall chart indicates the structures and relationships between most commonly discussed sphingolipids. A one-page thumbnail version of the chart is shown on page 98. Full size copies (approximately 35 x 26 inches) are available on request to customer service.

Package Weight

Unless otherwise specified, the package will contain at least the indicated amount and usually slightly more. The user is cautioned to always measure the required amount from the container.

Matreya's Mission

Matreya is committed to manufacturing high purity lipids to be used as research standards in the biotechnology and pharmaceutical areas. These lipids will be offered world-wide at a fair market price, and at a profit sufficient to assure company growth, for the benefit of its customers, employees, share holders, and community. Matreya will also be committed to fast delivery, excellent technical backup, new product development, safety, and environmentally friendly.


Spingoid Bases, Spingolipids and Glycospingolipids

Spingoid bases such as spingosine are the characteristic structural unit of the spingolipids. The bases are long chain aliphatic amines, containing two or three hydroxyl groups, and typically a *trans*-double bond at C4. In animal tissues, the most abundant base is spingosine with a C18 aliphatic chain containing a double bond in position 4. The saturated analogue is dihydrospingosine or sphinganine. In plants, the common long chain base is the 4 hydroxy saturated base phytospingosine.

Spingolipids are widely distributed in animal tissues, particularly cell membranes. Spingoid bases linked to fatty acids via an amide bond at C2 are ceramides and are present in trace amounts in most tissues. Glycospingolipids (ceramides having various mono- and oligosaccharides on the OH group at C1) are neutral glycospingolipids (i.e., cerebrosides and globosides). Those with sialic acid derivatized sugars are acidic glycolipids (i.e., gangliosides). They are amphiphilic and can be solubilized in buffers via sonication and micelle formation.


Gangliosides are present in substantial amounts in nerve cell membranes, and together with globosides are found in the membranes of white and red blood cells. These plus the glycospingolipids of the lacto- and neolacto-series are involved in cell recognition (e.g. blood group determinants). Glycolipid expression on the surface of cells determines their antigenicity as well as their status, i.e. differentiated vs. undifferentiated (embryonic), normal vs. malignant, etc. (1). The ganglioside GM1 stimulates nerve growth (2,3) and has been reported to have a curative effect on experimental Parkinsonism (4). For an overview, see (5). Gangliosides are also being investigated as potential anti-tumor vaccines (6). Glycospingolipids are also essential for the correct functioning of cell surface receptors (7). Matreya is your best source for many spingolipids. Most of Matreya's spingosines and ceramides are fully synthetic and as such 98%+ pure. Others, particularly the glycospingolipids are highly purified natural products (98%+), and can be used either as standards or biochemical reagents without further purification.

Through total synthesis, all four isomers of spingosine are available as well as a number of spingosines with other than 18 carbons and a number of ceramides (for details in using ceramides in cell culture see Hauser et al. [9]). Fluorescent labeled ceramides, glycospingolipids and spingomyelins are also available for study. D. N. Brindley and his group have been exploring the interaction of ceramides, spingosine and spingosine 1-phosphate in regulating DNA synthesis and phospholipase D activity. **See Literature References on page 99.**



Sphingosines

Synthetic Sphingosines with C18 Sphingoid Base



Catalog number 1802

1802 D-erythro-Sphingosine 25 mg

Sphingosine with C18 chain C₁₈H₃₇NO₂ CAS#: 123-78-4

Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: solid

Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

Selective inhibitor of phosphokinase C

1806 L-threo-Sphingosine 10 mg

L-threo-Sphingosine, C18 chain C₁₈H₃₇NO₂

Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: solid

Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

1826 L-erythro-Sphingosine 5 mg

L-erythro-Sphingosine, C18 chain C₁₈H₃₇NO₂ CAS#: 6036-75-5

Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: solid

Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

1827 D-threo-Sphingosine 5 mg

D-threo-Sphingosine, C18 chain C₁₈H₃₇NO₂ CAS#: 6036-85-7

Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: solid

Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

Synthetic Sphingosines with Sphingoid Bases other than C18

Varying chain lengths allow the study of translocation effects of sphingosines and ceramides into cells.

1833 D-erythro-C14-Sphingosine 5 mg

Sphingosine with C14 chain C₁₄H₂₉NO₂

Source: synthetic Mol. Wt.: 243 Purity: 98+% by TLC, GC Appearance: solid

Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C

1634 omega-N-NBD-D-erythro-C14-Sphingosine 1 mg

omega-N-(7-nitrobenzo-2-oxa-1,3-diazol-4-yl)-(2S)-amino-tetradec-(4E)-ene-(1,3R)-diol C₂₀H₃₁N₅O₅

Source: synthetic Mol. Wt.: 422 Purity: 98+% by TLC Appearance: solid

Solubility: methanol, ethanol, chloroform/methanol, 9:1 Storage: -20°C

1835 **D-erythro-C16-Sphingosine** **5 mg**
Sphingosine with C16 chain $C_{16}H_{33}NO_2$


Source: synthetic **Mol. Wt.:** 271 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

1838 **D-erythro-C12-Sphingosine** **5 mg**
Sphingosine with C12 chain $C_{12}H_{25}NO_2$ CAS#: 6918-49-6

Source: synthetic **Mol. Wt.:** 215 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$


1840 **D-erythro-C20-Sphingosine** **5 mg**
Sphingosine with C20 chain $C_{20}H_{41}NO_2$

Source: synthetic **Mol. Wt.:** 328 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$



Synthetic Dihydrospingosines

D,L-*threo*- Dihydrospingosine has also been found to be a significant inhibitor of sphingosine kinase (8). The D,L-*erythro*-isomer has been used as an inactive control. We offer all four isomers in pure form making detailed studies possible. Safingol, the L-*threo*-isomer is a potent inhibitor of PKC and as such is capable of reversing multi-drug resistance in cancer cells (9). **See Literature References on page 99.**



1807 **L-threo-Dihydrospingosine (Safingol)** **5 mg**
1807-025 L-*threo*-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ CAS#: 15639-50-6 **25 mg**

Source: synthetic **Mol. Wt.:** 301 **Melting Point ($^{\circ}C$):** 103-114 **Purity:** 98+% by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

Inhibitor of PKC

1831 **D-erythro-Dihydrospingosine** **25 mg**
1831-1 D-*erythro*-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ CAS#: 764-22-7 **1 g**

Source: synthetic **Mol. Wt.:** 301 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$

Inhibitor of PLA_2 and PLD

1846 **L-erythro-Dihydrosphingosine** **1 mg**
L-erythro-Sphinganine, C18 chain $C_{18}H_{39}NO_2$

Source: synthetic **Mol. Wt.:** 301 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$

1851 **D-threo-Dihydrosphingosine** **1 mg**
D-threo-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ **CAS#:** 6036-86-8

Source: synthetic **Mol. Wt.:** 301 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** $-20^{\circ}C$

1324 **D,L-erythro-Dihydrosphingosine** **25 mg**
D,L-erythro-Sphinganine, C18 chain $C_{18}H_{39}NO_2$ **CAS#:** 3102-56-5

Source: synthetic **Mol. Wt.:** 301 **Purity:** erythro 77%; threo 23% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$

Inhibitor of sphingosine kinase

1326 **D,L-C16-Dihydrosphingosine (mixed isomers)** **10 mg**
D,L-Sphinganine with C16 chain $C_{16}H_{35}NO_2$

Source: synthetic **Mol. Wt.:** 273 **Purity:** erythro 90%, threo 10% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: $-20^{\circ}C$


1845 **D-erythro-C20-Dihydrosphingosine** **5 mg**
D-erythro-Sphinganine, C20 chain $C_{20}H_{43}NO_2$ **CAS#:** 24006-62-0

Source: synthetic **Mol. Wt.:** 330 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: warm ethanol, chloroform/methanol, 5:1 **Storage:** $-20^{\circ}C$

1839 **D,L-erythro-C20-Dihydrosphingosine** **10 mg**
D,L-erythro-Sphinganine, C20 chain $C_{20}H_{43}NO_2$

Source: synthetic **Mol. Wt.:** 330 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: warm ethanol, chloroform/methanol, 5:1 **Storage:** $-20^{\circ}C$

3-Keto-Dihydrosphingosines



1876 **3-keto-Dihydrosphingosine•HCl** **10 mg**
3-keto-Sphinganine hydrochloride $C_{18}H_{37}NO_2 \cdot HCl$ **CAS#:** 18944-28-0

Source: synthetic **Mol. Wt.:** 299 + HCl **Purity:** 98+% by TLC, GC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

1891 **3-keto-C6-Dihydrospingosine•HCl** **10 mg**
1-Hydroxy-2-amino-3-keto-hexane • HCl $C_6H_{13}NO_2 \cdot HCl$

Source: synthetic **Mol. Wt.:** 168 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol, methanol, DI water **Storage:** -20°C


1892 **3-keto-C8-Dihydrospingosine•HCl** **10 mg**
1-Hydroxy-2-amino-3-keto-octane • HCl $C_8H_{17}NO_2 \cdot HCl$

Source: synthetic **Mol. Wt.:** 196 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DI water **Storage:** -20°C

1893 **3-keto-C12-Dihydrospingosine•HCl** **10 mg**
1-Hydroxy-2-amino-3-keto-dodecane • HCl $C_{12}H_{25}NO_2 \cdot HCl$

Source: synthetic **Mol. Wt.:** 252 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

Phytosphingosines



1330 **Phytosphingosine** **50 mg**
1330-1 4-Hydroxysphinganine $C_{18}H_{39}NO_3$ **CAS# 554-62-1** **1 g**


Source: natural, yeast (*Pichia ciferri*) **Mol. Wt.:** 318 **Purity:** 98+% by TLC, GC
Appearance: solid **Solubility:** ethanol, methanol, chloroform/methanol, 2:1 (warm)
Storage: -20°C

Other Sphingosine Derivatives and Precursors

1320 **N,N-Dimethyl-D-erythro-sphingosine** **5 mg/ml, 1 ml**
 $C_{20}H_{41}NO_2$ **CAS#:** 119567-63-4

Source: synthetic **Mol. Wt.:** 328 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: isopropanol **Solubility:** chloroform, ethanol, isopropanol, methanol
Storage: -20°C

Inhibitor of phosphokinase C



1896 **N,N-Dihexyl-D-erythro-sphingosine** **5 mg/ml, 1 ml**
 Sphingosine with tertiary amine group $C_{30}H_{61}NO_2$

Source: synthetic **Mol. Wt.:** 468 **Purity:** 95% by TLC **Appearance:** liquid
Solvent: ethanol **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

1805 **N-Palmitoyl serinol** **10 mg**
 $C_{19}H_{39}NO_3$ **CAS#:** 126127-31-9

Source: synthetic **Mol. Wt.:** 329 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, methanol, ethanol **Storage:** $-20^{\circ}C$

Sphingosine precursor

Ceramides

Ceramide is a fatty acid amide of sphingosine. It may be formed by dehydrogenation of dihydroceramide; by hydrolysis of sphingomyelin or glycosphingolipids; or by acylation of free sphingosine. Ceramide functions as a precursor in the synthesis of sphingomyelin (by an exchange reaction with phosphatidylcholine and phosphatidylethanolamine); of glycosphingolipids (by glycosylation with UDP-hexose); and of free sphingosine and fatty acid by hydrolysis. The sphingosine can be phosphorylated by a kinase to form sphingosine-1-phosphate, which may undergo further hydrolysis or cleavage.

Control of sphingolipid metabolism maintains vital balance points in cell physiology. Two of ceramide's metabolites, sphingosine-1-phosphate and glucosylceramide, produce cell proliferation. Sphingosine-1-phosphate is also a highly active regulator of angiogenesis, vascular maturation, cardiac development, immunity, and directed cell movement. Sphingosine, the free base, is a potent inhibitor of protein kinase C and is involved in intracellular calcium regulation.

Sphingolipid enzymes seem to be particularly active in cancers, so modifying their activities by exogenous action may provide alternatives to chemical therapies. These enzymes are controlled by many known agents, such as 1,25-dihydroxy-vitamin D_3 , tumor necrosis factor- α , nerve growth factor, interleukin 1, endothelial growth factor, glutathione, arachidonic acid, dexamethasone, many anticancer drugs, therapeutic radiation, and activators of the FAS receptor.

Ceramide exerts numerous biological effects, including induction of cell maturation, cell cycle arrest, terminal cell differentiation, cell senescence, and cell death. Other effects include producing reactive oxygen in mitochondria (followed by apoptosis) and stimulating phosphorylation of certain proteins (especially mitogen activated protein). It also stimulates some protein phosphatases (especially protein phosphatase 2A). Thus ceramide is an important controller of protein activity


It is apparent from these relationships that ceramide exists at the crux of several enzyme reaction cycles and that experiments involving sphingolipid function call for control of all of the cycles and their branch-off points. Matreya is the major supplier of these lipids, which can be used as standards for analysis of tissues (a much needed part of modern research) and identification of major sphingolipids.

Ceramides with short side chains have been shown to enter easily into cells where they are biologically active. Ceramides with longer side chains, however, also enter cells if dissolved in dodecane-isopropanol first. Fluorescent labeled ceramides and sphingomyelin made from fluorescent labeled acids instead of plain fatty acids are also available for the study of the localization and metabolism of sphingolipids in the cell. Matreya now offers all four isomers of C2, C4, C6 and C18 ceramides. The corresponding dihydroceramides are being used as inactive controls

In three major reviews, Radin (10-12) has discussed the biochemistry and chemistry of ceramide and outlined many potential approaches to cancer therapy using ceramides and related compounds as generators of apoptosis.

See Literature References on page 99.

Synthetic Ceramides Derived from C18-Sphingosine



1901 **N-Acetyl-D-erythro-sphingosine** **10 mg**
1901-100 N-C2:0-D-erythro-Ceramide C₂₀H₃₉NO₃ CAS#: 3102-57-6 **100 mg**

Source: synthetic Mol. Wt.: 342 Purity: 98+ by TLC, GC Appearance: solid
Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml) Storage: -20°C

1829 **N-Acetyl-L-threo-sphingosine** **1 mg**
N-C2:0-L-threo-Ceramide C₂₀H₃₉NO₃

Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: solid
Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml)
Storage: -20°C

1847 **N-Acetyl-L-erythro-sphingosine** **1 mg**
N-C2:0-L-erythro-Ceramide C₂₀H₃₉NO₃

Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: solid
Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml)
Storage: -20°C

1900 **N-Hexanoyl-D-erythro-sphingosine** **10 mg**
1900-100 N-C6:0-D-erythro-Ceramide C₂₄H₄₇NO₃ CAS#: 124753-97-5 **100 mg**

Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: solid
Solubility: chloroform, ethanol, DMSO (up to 5 mg/ml) Storage: -20°C

1828 **N-Hexanoyl-L-threo-sphingosine** **1 mg**
N-C6:0-L-threo-Ceramide C₂₄H₄₇NO₃

Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C

1848 **N-Hexanoyl-L-erythro-sphingosine** **1 mg**
N-C6:0-L-erythro-Ceramide C₂₄H₄₇NO₃

Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC; GC Appearance: solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C

1809	N-Hexanoyl-D-threo-sphingosine N-C6:0-D-threo-Ceramide C ₂₄ H ₄₇ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methanol, DMSO (up to 5 mg/ml) Storage: -20°C	
1903 1903-100	N-Octanoyl-D-erythro-sphingosine N-C8:0-D-erythro-Ceramide C ₂₆ H ₅₁ NO ₃ CAS#: 74713-59-0	10 mg 100 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+ by TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml) Storage: -20°C	
1830	N-Octanoyl-L-threo-sphingosine N-C8:0-L-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C	
1810	N-Octanoyl-D-threo-sphingosine N-C8:0-D-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
1333 1333-100	N-Decanoyl-D-erythro-sphingosine N-C10:0-D-erythro-Ceramide C ₂₈ H ₅₅ NO ₃	10 mg 100 mg
	Source: synthetic Mol. Wt.: 454 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol, DMSO (up to 5mg/ml) Storage: -20°C	
2037 2037-100	N-Pentadecanoyl-D-erythro-sphingosine N-C15:0-D-erythro-Ceramide C ₃₃ H ₆₅ NO ₃	10 mg 100 mg
	Source: synthetic Mol. Wt.: 524 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
1915 1915-100	N-Hexadecanoyl-D-erythro-sphingosine N-C16:0-D-erythro-Ceramide; N-Palmitoyl-D-erythro-sphingosine C ₃₄ H ₆₇ NO ₃ CAS#: 24696-26-2	10 mg 100 mg
	Source: synthetic Mol. Wt.: 538 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
2038 2038-100	N-Heptadecanoyl-D-erythro-sphingosine N-C17:0-D-erythro-Ceramide C ₃₅ H ₆₉ NO ₃ CAS#: 67492-16-4	10 mg 100 mg
	Source: synthetic Mol. Wt.: 552 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	

1832 **N-Octadecanoyl-D-erythro-sphingosine** **10 mg**
1832-100 N-C18:0-D-*erythro*-Ceramide; N-Stearoyl-D-*erythro*-sphingosine **100 mg**
C₃₆H₇₁NO₃ **CAS#:** 2304-81-6

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1 (up to 5mg/ml)
Storage: -20°C

2039 **N-Nonadecanoyl-D-erythro-sphingosine** **10 mg**
2039-100 N-C19:0-D-*erythro*-Ceramide C₃₇H₇₃NO₃ **100 mg**

Source: synthetic **Mol. Wt.:** 580 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, warm ethanol, warm methanol **Storage:** -20°C

1843 **N-Octadecanoyl-L-threo-sphingosine** **1 mg**
N-C18:0-L-*threo*-Ceramide; N-Stearoyl-L-*threo*-sphingosine C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

1850 **N-Octadecanoyl-L-erythro-sphingosine** **1 mg**
N-C18:0-L-*erythro*-Ceramide; N-Stearoyl-L-*erythro*-sphingosine C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

1855 **N-Octadecanoyl-D-threo-sphingosine** **1 mg**
N-C18:0-D-*threo*-Ceramide; N-Stearoyl-D-*threo*-sphingosine C₃₆H₇₁NO₃

Source: synthetic **Mol. Wt.:** 566 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) **Storage:** -20°C

1916 **N-Tetracosanoyl-D-erythro-sphingosine** **5 mg**
1916-25 N-C24:0-D-*erythro*-Ceramide; N-Lignoceroyl-D-*erythro*-sphingosine **25 mg**
C₄₂H₈₃NO₃ **CAS#:** 34435-05-7

Source: synthetic **Mol. Wt.:** 650 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform **Storage:** -20°C

1930 **N-Tetracosenoyl-D-erythro-sphingosine** **5 mg**
1930-25 N-*cis*-15-C24:1-D-*erythro*-Ceramide; N-Nervonoyl-D-*erythro*-sphingosine **25 mg**
C₄₂H₈₁NO₃ **CAS#:** 54164-50-0

Source: synthetic **Mol. Wt.:** 648 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO, warm methanol **Storage:** -20°C

2-Hydroxy Ceramides

2042 **N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-sphingosine** **5 mg**
N-(R,S)-*alpha*-Hydroxy-C12:0-D-*erythro*-ceramide C₃₀H₅₉NO₄

Source: synthetic **Mol. Wt.:** 498 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, methanol, ethanol, DMSO **Storage:** -20°C

2044 **N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-sphingosine** **5 mg**
N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-ceramide; N-(R,S)-alpha-Hydroxystearoyl-D-erythro-sphingosine C₃₆H₇₁NO₄

Source: synthetic **Mol. Wt.:** 582 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.5 **Storage:** -20°C

Ceramide Made from Sphingosines with Sphingoid Bases Other Than C18


1842 **N-Acetyl-D-erythro-sphingosine (C14 sphingoid base)** **5 mg**
N-C2:0 Ceramide of D-erythro-C14-sphingosine C₁₆H₃₁NO₃

Source: synthetic **Mol. Wt.:** 285 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml) **Storage:** -20°C

2077 **N-Hexadecanoyl-D-erythro-C16-sphingosine (C16 sphingoidbase)** **1 mg**
N-Palmitoyl-D-erythro-C16-sphingosine; N-C16:0 Ceramide of D-erythro-C16-sphingosine C₃₂H₆₃NO₃

Source: synthetic **Mol. Wt.:** 510 **Purity:** 98% by TLC, MS **Appearance:** solid
Solubility: chloroform, warm ethanol, warm methanol **Storage:** -20°C

Dihydroceramides



1834 **N-Acetyl-D-erythro-dihydrosphingosine** **5 mg**
N-C2:0-D-erythro-Dihydroceramide; N-Acetyl-D-erythro-sphinganine
C₂₀H₄₁NO₃

Source: synthetic **Mol. Wt.:** 344 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

1910 **N-Hexanoyl-D-erythro-dihydrosphingosine** **5 mg**
N-C6:0-D-erythro-Dihydroceramide; N-Hexanoyl-D-erythro-sphinganine
C₂₄H₄₉NO₃

Source: synthetic **Mol. Wt.:** 400 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** -20°C

1854 **N-Octanoyl-D-erythro-dihydrosphingosine** **5 mg**
N-C8:0-D-erythro-Dihydroceramide; N-Octanoyl-D-erythro-sphinganine
C₂₆H₅₃NO₃

Source: synthetic **Mol. Wt.:** 428 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, ethanol, DMSO **Storage:** -20°C

2041 **N-Octadecanoyl-D-erythro-dihydrosphingosine** **10 mg**
 N-C18:0-D-erythro-Dihydroceramide; N-Octadecanoyl-D-erythro-sphinganine; N-Stearoyl-D-erythro-dihydrosphingosine $C_{36}H_{73}NO_3$
Source: synthetic **Mol. Wt.:** 568 **Purity:** 98% by TLC **Appearance:** solid
Solubility: hot ethanol, DMSO, warm chloroform/methanol, 5:1 **Storage:** -20°C


2-Hydroxy Dihydroceramides

2043 **N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-dihydrosphingosine** **5 mg**
 N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-dihydroceramide $C_{30}H_{61}NO_4$
Source: synthetic **Mol. Wt.:** 500 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.5 **Storage:** -20°C

2045 **N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-dihydrosphingosine** **5 mg**
 N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-dihydroceramide; N-(R,S)-alpha-Hydroxystearoyl-D-erythro-dihydrosphingosine $C_{36}H_{73}NO_4$
Source: synthetic **Mol. Wt.:** 584 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.5 **Storage:** -20°C

2047 **N-(R,S)-alpha-Hydroxyhexadecanoyl-D-erythro-dihydrosphingosine** **5 mg**
 N-(R,S)-alpha-Hydroxy-C16:0-D-erythro-dihydroceramide; N-(R,S)-alpha-Hydroxypalmitoyl-D-erythro-dihydrosphingosine $C_{34}H_{69}NO_4$
Source: synthetic **Mol. Wt.:** 556 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.5 **Storage:** -20°C

Ceramides From Natural Sources



General ceramide structure

1056 **Ceramides** **25 mg**
 Ceramides with hydroxy and non-hydroxy acyl groups
 $C_{42}H_{83}NO_4$ **CAS#:** 104404-17-13
Source: natural, bovine **Mol. Wt.:** 666(2-hydroxylignoceroyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C


1322 **Ceramides** **10 mg**
1322-05 Ceramides with mostly non-hydroxy acyl groups $C_{36}H_{71}NO_3$ **50 mg**
Source: natural, bovine **Mol. Wt.:** 566 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, chloroform/methanol, 2:1 **Storage:** -20°C

2036 **N-Tetracosanoyl-phytosphingosine** **5 mg**
N-C24:0-Phytoceramide; N-Lignoceroyl-phytosphingosine C₄₂H₈₅NO₄
Source: semisynthetic, yeast (*Pichia ciferri*) **Mol. Wt.:** 668 **Purity:** 98+% by TLC,
MS **Appearance:** solid **Solubility:** chloroform/methanol, 5:1 **Storage:** -20°C

Fluorescent Ceramides

1841 **N-Hexanoyl-NBD-D-erythro-sphingosine** **100 µg**
1841-001 N-C6:0-NBD-Ceramide; N-C6:0-NBD-D-erythro-Sphingosine C₃₀H₄₉N₅O₆ **1 mg**
CAS#: 86701-10-2

Source: synthetic **Mol. Wt.:** 576 **Melting Point (°C):** 85.7-87.9 **Purity:** 98+% by
TLC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C



Catalog number 1618

Excitation: 460 nm
Emission: 535 nm

1618 **N-Dodecanoyl-NBD-D-erythro-sphingosine** **100 µg**
1618-001 N-C12:0-NBD-Ceramide; N-C12:0-NBD-D-erythro-Sphingosine C₃₆H₆₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1857 **N-Hexanoyl-NBD-L-threo-sphingosine** **100 µg**
1857-001 N-C6:0-NBD-Ceramide; N-C6:0-NBD-L-threo-Sphingosine C₃₀H₄₉N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 576 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

1620 **N-Dodecanoyl-NBD-L-threo-sphingosine** **100 µg**
1620-001 N-C12:0-NBD-Ceramide; N-C12:0-NBD-L-threo-Sphingosine C₃₆H₆₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1624 **N-Hexanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
1624-001 N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-L-threo-Dihydrosphingosine C₃₀H₅₁N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1623 **N-Dodecanoyl-NBD-L-*threo*-dihydrosphingosine** **100 µg**
1623-001 N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-L-*threo*-Dihydrosphingosine **1 mg**
C₃₆H₆₃N₅O₆

Source: synthetic **Mol. Wt.:** 662 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1626 **N-Hexanoyl-NBD-D-*erythro*-dihydrosphingosine** **100 µg**
1626-001 N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-D-*erythro*-Dihydrosphingosine **1 mg**
C₃₀H₅₁N₅O₆

Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1625 **N-Dodecanoyl-NBD-D-*erythro*-dihydrosphingosine** **100 µg**
1625-001 N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-D-*erythro*-
Dihydrosphingosine C₃₆H₆₃N₅O₆ **1 mg**

Source: synthetic **Mol. Wt.:** 662 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1628 **N-Hexanoyl-NBD-phytosphingosine** **100 µg**
1628-001 N-C6:0-NBD-Phytoceramide; N-C6:0-NBD-Phytosphingosine C₃₀H₅₁N₅O₇ **1 mg**

Source: semisynthetic, bacteria **Mol. Wt.:** 594 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1627 **N-Dodecanoyl-NBD-phytosphingosine** **100 µg**
1627-001 N-C12:0-NBD-Phytoceramide; N-C12:0-NBD-Phytosphingosine C₃₆H₆₃N₅O₇ **1 mg**


Source: semisynthetic, bacteria **Mol. Wt.:** 678 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

**See Biochemicals and Reagents section (page 88) for additional
fluorescent labeled products.**

**Compounds with fluorescent labels other than NBD are available
on custom basis. Contact Technical Service for more information.**

Phosphosphingolipids

Sphingomyelins



Catalog number 1051

1051 **Sphingomyelin** **25 mg**
1051-1 SPM; Ceramide-1-phosphorylcholine C₄₁H₈₃N₂O₆P CAS#: 85187-10-6 **1 g**

Source: natural, bovine **Mol. Wt.:** 731 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1328 **Sphingomyelin** **25 mg**
SPM; Ceramide-1-phosphorylcholine C₄₇H₉₅N₂O₆P CAS#: 85187-10-6

Source: natural, porcine RBC **Mol. Wt.:** 815 (lignoceroyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1329 **Sphingomyelin** **25 mg**
1329-1 SPM; Ceramide-1-phosphorylcholine C₄₆H₉₃N₂O₆P CAS#: 85187-10-6 **1 gram**

Source: natural, bovine buttermilk **Mol. Wt.:** 801 (tricosanoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1332 **Sphingomyelin** **25 mg**
1332-1 SPM; Ceramide-1-phosphorylcholine C₃₉H₇₉N₂O₆P **1 gram**

Source: natural, chicken, egg **Mol. Wt.:** 703 (palmitoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, methanol, warm ethanol **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1907 **N-Acetyl-sphingosylphosphorylcholine** **5 mg**
N-C2:0-Sphingomyelin C₂₅H₅₁N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 506 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, chloroform/methanol, 2:1 **Storage:** -20°C

Mixture of D-erythro and L-threo isomers

1909 **N-Hexanoyl-sphingosylphosphorylcholine** **5 mg**
N-C6:0-Sphingomyelin C₂₉H₅₉N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 563 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** ethanol, chloroform/methanol, 2:1 **Storage:** -20°C

Mixture of D-erythro and L-threo isomers

1911 N-Octadecanoyl-sphingosylphosphorylcholine 5 mg

N-C18:0-Sphingomyelin; N-Stearoyl-sphingosylphosphorylcholine
C₄₁H₈₃N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 731 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Mixture of D-erythro and L-threo isomers

1890 N-Heptadecanoyl-sphingosylphosphorylcholine 5 mg

N-C17:0-Sphingomyelin C₄₀H₈₁N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 717 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Mixture of D-erythro and L-threo isomers


1917 N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine 0.5 mg

N-C20:0-Sphingomyelin C₄₃H₈₇N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 759 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** ethanol, methanol, chloroform/methanol, 14:1

Storage: -20°C



Catalog number 1918

1918 N-Docosanoyl-D-erythro-sphingosylphosphorylcholine 0.5 mg

N-C22:0-Sphingomyelin C₄₅H₉₁N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 787 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** ethanol, methanol, chloroform/methanol, 14:1

Storage: -20°C

2200 N-1-¹³C-Hexadecanoyl-sphingosylphosphorylcholine 1mg

D-erythro-Sphingomyelin with 1-¹³C-palmitic acid; N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine ¹²C₃₈¹³CH₇₉N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 703 **Purity:** 98+% by TLC


Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1327 N-Acyl-sphingosylphosphorylethanolamine 5 mg

Ceramide phosphorylethanolamine C₄₃H₈₇N₂O₆P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 773 (tricosanoyl) **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

Sphingosylphosphorylcholines (SPC)



Catalog number 1318

1318 **D-erythro-Sphingosylphosphorylcholine** **5 mg**
D-erythro-SPC C₂₃H₄₉N₂O₅P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 465 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1319 **L-threo-Sphingosylphosphorylcholine** **5 mg**
L-threo-SPC C₂₃H₄₉N₂O₅P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 465 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C


1321 **Sphingosylphosphorylcholine** **10 mg**
1321-05 lyso-Sphingomyelin; SPC (mixture of D-erythro and L-threo isomers) **50 mg**
C₂₃H₄₉N₂O₅P **CAS#:** 82970-80-7

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 465 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1913 **lyso-Dihydro-sphingomyelin** **1 mg**
Dihydro-sphingosylphosphorylcholine (mixture of D-erythro and L-threo isomers) C₂₃H₅₁N₂O₅P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 467 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

Sphingosine Phosphates



Catalog number 1803

1803 **D-erythro-Sphingosine-1-phosphate** **5 mg**
S-1-P C₁₈H₃₈NO₅P **CAS#:** 26993-30-6

Source: synthetic **Mol. Wt.:** 380 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform plus a few drops of TFA, chloroform/methanol/40% dimethylamine, 5:15:3, 1mg/ml **Storage:** -20°C


1852 **D-erythro-Dihydrosphingosine-1-phosphate** **5 mg**
C₁₈H₄₀NO₅P CAS#: 19794-97-9

Source: synthetic **Mol. Wt.:** 382 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform plus a few drops of TFA, chloroform/methanol/40%
dimethylamine, 5:15:3, 1mg/ml **Storage:** -20°C

2046 **N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate, NH₄⁺ salt** **5 mg**
N-C16:0-Ceramide-1-phosphate C₃₄H₆₈NO₆P•NH₃

Source: synthetic **Mol. Wt.:** 618+NH₃ **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/acetic acid, 60:15:25 **Storage:** -20°C

Fluorescent Sphingomyelins



1912 **N-Hexanoyl-NBD-sphingosylphosphorylcholine** **100 µg**
1912-001 N-C6:0-NBD-Sphingomyelin; N-C6:0-NBD-Sphingosylphosphorylcholine **1 mg**
C₃₅H₆₁N₆O₉P CAS#: 94885-04-8

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 740 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Mixture of *D-erythro* and *L-threo* isomers

1619 **N-Dodecanoyl-NBD-sphingosylphosphorylcholine** **100 µg**
1619-001 N-C12:0-NBD-Sphingomyelin; N-C12:0-NBD-Sphingosylphosphorylcholine **1 mg**
C₄₁H₇₃N₆O₉P

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 825 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

Mixture of *D-erythro* and *L-threo* isomers

See Biochemicals and Reagents section (page 88) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information.


Glycosphingolipids

Glycosphingolipids are widely distributed in animal and plant tissues. They consist of a ceramide (Cer) bound in glycosidic linkage through the primary hydroxyl to a mono- or oligosaccharide which may contain substituents such as a sulfate, acetate, or phosphate group. They are amphiphilic and the less glycosylated compounds can be dispersed in buffers by dissolving them in a detergent or organic solvent (EtOH, DMSO, isoPrOH) and mixing by sonication.


Galactosphingolipids, mainly GalCer (cerebrosides) and its sulfate ester, occur in large amounts in the nervous system. Glucosphingolipids, the simplest of which is GlcCer (glucocerebrosides), are very widely distributed, particularly in nerve cell membranes. GlcCer is isolated from a variety of natural sources including human, bovine, and plant. Each of these sources has a heterogeneity in the fatty acid content of the ceramide as well as an occasional variation in the sphingoid chain. Globosides (containing both glucose and galactose) are a prominent group of glycosphingolipids, they contain an α -linked galactose moiety and are typically located in blood cell membranes. Gangliosides are another prominent group of glycosphingolipids; they are acidic because of substitution with sialic (neuraminic) acid. The glycosphingolipids function in a wide range of enzyme and structural interactions, such as immunological or membrane recognition phenomena, binding of microbial pathogens, hormone and growth factor actions, cancer cell growth and malignancy, atherosclerosis, genetic disease errors, blood group determinants, etc. Tissues change in glycosphingolipid composition during embryogenesis, maturation, aging, and other vital physiological processes. Some glycosphingolipids stimulate cell proliferation, others induce apoptosis, effects of great significance to cancer therapy and maturational development. Marked differences in glycosphingolipid composition are seen in normal and cancerous cells. See references (13-25).

See Literature References on page 99.

Galactosylceramides and Glucosylceramides



Galactosylceramide



Glucosylceramide

1050

Cerebrosides

50 mg

Galactosylceramide; Ceramide *beta*-D-galactoside

$C_{48}H_{93}NO_9$ CAS#: 85305-88-0

Source: natural, bovine Mol. Wt.: 828 (2-hydroxytetracosanoyl) Purity: 98+% by

TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: $-20^{\circ}C$

See Table III (pg. 93-97) for fatty acid content

1066

Cerebroside; Kerasin (top spot)

10 mg

Galactosylceramide with mostly non-hydroxy fatty acid side chain

$C_{42}H_{81}NO_8$ CAS#: 536-13-0

Source: natural, bovine Mol. Wt.: 810 (nervonyl, [24:1]) Purity: 98+% by TLC

Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.5

Storage: $-20^{\circ}C$

See Table III (pg. 93-97) for fatty acid content

1138 Cerebroside; Phrenosin (bottom spot) 10 mg

Galactosylceramide with mostly 2-hydroxy fatty acid side chains
 $C_{42}H_{81}NO_9$ CAS#: 37211-11-3

Source: natural, bovine **Mol. Wt.:** 744 (2-hydroxystearoyl) **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.5

Storage: $-20^{\circ}C$

See Table III (pg. 93-97) for fatty acid content

1305 Psychosine (free amine form) 10 mg

lyso-Cerebroside; 1-*beta*-D-Galactosylsphingosine

$C_{24}H_{47}NO_7$ CAS#: 2238-90-6

Source: semisynthetic, bovine **Mol. Wt.:** 461 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** ethanol, chloroform/methanol/DI water, 5:1:0.1

Storage: $-20^{\circ}C$

1914 N-Octadecanoyl-D₃₅-psychosine (perdeuterated C_{18:0} fatty acid) 5 mg


N-C_{18:0}-D₃₅-Cerebroside, perdeuterated; N-Stearoyl-D₃₅-Psychosine, perdeuterated $C_{42}H_{46}D_{35}NO_8$

Source: semisynthetic, bovine **Mol. Wt.:** 762 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform, hot ethanol, chloroform/methanol, 2:1

Storage: $-20^{\circ}C$

Deuterium labeled stearoyl sidechain



Catalog number 1325

1325 N-Acetyl-psychosine 10 mg

N-C_{2:0}-Cerebroside $C_{26}H_{49}NO_8$

Source: semisynthetic, bovine **Mol. Wt.:** 503 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** $-20^{\circ}C$

1335 N-Pentadecanoyl-psychosine 5 mg

N-C_{15:0}-Cerebroside $C_{39}H_{75}NO_8$

Source: semisynthetic, bovine **Mol. Wt.:** 686 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform/ methanol, 2:1 **Storage:** $-20^{\circ}C$

1334 N-Octanoyl-*beta*-D-galactosylceramide 10 mg
1334-50 50 mg

N-C_{8:0}-Galactosylceramide $C_{32}H_{61}NO_8$

Source: semisynthetic, bovine **Mol. Wt.:** 588 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** ethanol, methanol, chloroform/methanol, 9:1

Storage: $-20^{\circ}C$

1621 **N-Hexanoyl-NBD-galactosylceramide** **100 µg**
1621-001 N-C6:0-NBD-*beta*-D-Galactosylsphingosine; N-C6:0-NBD-Cerebroside **1 mg**
C₃₆H₅₉N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1 **Storage:** -20°C

1633 **N-Dodecanoyl-NBD-galactosylceramide** **100 µg**
1633-001 N-C12:0-NBD-*beta*-D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside **1 mg**
C₄₂H₇₁N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 822 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, DMSO, chloroform/methanol, 2:1 **Storage:** -20°C

1057 **Glucocerebrosides, Gaucher's spleen** **5 mg**
1057-25 Glucosylceramide; Ceramide *beta*-D-glucoside C₄₈H₉₃NO₈ **25 mg**
CAS# 85305-87-9


Source: natural, human **Mol. Wt.:** 812 (lignoceryl) **Purity:** 98+% by TLC, GC, MS
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1521 **Glucocerebrosides** **5 mg**
1521-50 Glucosylceramide; Ceramide *beta*-D-glucoside C₄₆H₈₉NO₈ **50 mg**

Source: natural, bovine buttermilk **Mol. Wt.:** 784 (docosanoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

See Table III (pg. 93-97) for side chain variants



Catalog number 1522

1522 **Glucocerebrosides, plant** **5 mg**
1522-100 Glucosylceramide; Ceramide *beta*-D-glucoside C₄₀H₇₅NO₉ **100 mg**

Source: natural, plant **Mol. Wt.:** 714 (2-hydroxyhexadecanoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4,8) and most of the fatty acids are of the 2-hydroxy type. See Table III (pg. 93-97) for fatty acid content

1622 **N-Hexanoyl-NBD-glucosylceramide** **100 µg**
1622-001 N-C6:0-NBD-*beta*-D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide **1 mg**
C₃₆H₅₉N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1 **Storage:** -20°C


1306 **Glucopsychosine** **5 mg**

Glucosylsphingosine; *lyso*-Glucocerebroside; 1-*beta*-D-Glucosylsphingosine
C₂₄H₄₇NO₇ CAS#: 52050-17-6

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 461 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** ethanol, methanol, chloroform/methanol, 2:1

Storage: -20°C



Catalog number 1310

1310 **Glucopsychosine** **5 mg**

Glucosylsphingosine; *lyso*-Glucocerebroside; 1-*beta*-D-Glucosylsphingadienine C₂₄H₄₅NO₇ CAS#: 52050-17-6

Source: natural, plant **Mol. Wt.:** 460 (based on 1-*beta*-D-glucosylsphinga-4,8-dienine)

Purity: 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol, 4:1

Storage: -20°C

Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4, 8)

1531 **N-Docosanoyl-glucopsychosine** **1 mg**

N-C22:0-Glucocerebroside; N-Docosanoyl-*beta*-glucosylsphingosine
C₄₆H₈₉NO₈

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 784 **Purity:** 98+% by TLC


Appearance: solid **Solubility:** chloroform **Storage:** -20°C

1533 **N-Hexadecanoyl-D₃-glucopsychosine, deuterated** **1 mg**

N-C16:0-D₃-Glucopsychosine, deuterated; N-C16:0-D₃-Glucocerebroside, deuterated; N-Palmitoyl-D₃-glucopsychosine, deuterated C₄₀H₇₄D₃NO₈

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 703 **Purity:** 98+% by TLC

Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C



Catalog number 1049

1049 **Sulfatides** **50 mg**

Ceramide-galactoside-3-sulfate; Cerebroside sulfate C₄₂H₈₀NNaO₁₁S
CAS#: 85496-63-5

Source: natural, bovine **Mol. Wt.:** 830 (stearoyl) Na⁺ Salt **Purity:** 98+% by TLC

Appearance: solid **Solubility:** DMSO, chloroform/methanol/DI water, 2:1:0.1 (if needed, a few drops of acetic acid) **Storage:** -20°C


See Table III (pg. 93-97) for fatty acid content

1904	lyso-Sulfatide (NH₄⁺ salt) Sphingosine-1-galactoside-3-sulfate C ₂₄ H ₄₇ NO ₁₀ S•NH ₃ CAS#: 38621-58-8	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 542 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C	
2076	N-Acetyl-sulfatide N-C2:0-Sulfatide; N-Acetyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfatide C ₂₆ H ₄₉ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 584 Purity: 98+% by TLC Appearance: solid Solubility: ethanol, methanol, chloroform/methanol, 1:1 Storage: -20°C	
1875	N-Hexadecanoyl-sulfatide N-C16:0-Sulfatide; N-Palmitoyl-sulfatide; N-Palmitoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate C ₄₀ H ₇₇ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 780 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C	
1932	N-Octadecanoyl-sulfatide N-C18:0-Sulfatide; N-Octadecanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate C ₄₂ H ₈₁ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 808 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1933	N-Octadecenoyl-sulfatide N-C18:1-Sulfatide; N-Octadecenoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate C ₄₂ H ₇₉ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 806 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1888	N-Tetracosanoyl sulfatide N-C24:0-Sulfatide; N-Tetracosanoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate; N-Lignoceroyl-sulfatide C ₄₈ H ₉₃ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 892 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1931	N-Tetracosenoyl-sulfatide N-Nervonyl-sulfatide; N-C24:1-Sulfatide; N-Tetracosenoyl-sphingosyl- <i>beta</i> -D-galactoside-3-sulfate C ₄₈ H ₉₁ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 890 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide, deuterated N-C18:0-D ₃ -Sulfatide, deuterated; N-Stearoyl-D ₃ -sulfatide, deuterated C ₄₂ H ₇₈ D ₃ NO ₁₁ S	1 mg
	Source: semisynthetic, bovine Mol. Wt.: 811 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C	

1632 **N-Dodecanoyl-NBD-sulfatide** **100 µg**
1632-001 N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD-*lyso*-sulfatide; N-Dodecanoyl-
NBD-sphingosyl-*beta*-D-galactoside-3-sulfate $C_{42}H_{71}N_5O_{14}S$ **1 mg**

Source: semisynthetic, bovine **Mol. Wt.:** 901 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

Lactosylceramides



1500 **Lactosylceramides** **1 mg**
LC; Lactocerebrosides; CDH; Ceramide *beta*-lactoside
 $C_{48}H_{91}NO_{13}$ **CAS#:** 4682-48-8

Source: natural, porcine RBC **Mol. Wt.:** 890 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, chloroform/methanol/DI water, 5:1:0.1
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1507 **Lactosylceramides** **5 mg**
1507-50 LC; Lactocerebrosides; CDH; Ceramide *beta*-lactoside $C_{53}H_{101}NO_{13}$ **50 mg**
CAS#: 4682-48-8

Source: natural, bovine buttermilk **Mol. Wt.:** 960 (tricosanoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 5:1:0.1
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1517 ***lyso*-Lactosylceramide** **1 mg**
Lactosylsphingosine; *lyso*-LC $C_{30}H_{57}NO_{12}$

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 623 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1
Storage: -20°C

1532 **N-Hexadecanoyl-lactosylceramide** **1 mg**
N-C16:0-Lactosylceramide; N-Palmitoyl-lactosylceramide $C_{46}H_{87}NO_{13}$


Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 862 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1
Storage: -20°C

1534 **N-Hexadecanoyl-D₃-lactosylceramide, deuterated** **1 mg**
N-C16:0-D₃-Lactosylceramide, deuterated; N-Palmitoyl-D₃-lactosylceramide,
deuterated $C_{46}H_{84}D_3NO_{13}$

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 865 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 5:1:0.1
Storage: -20°C

1538	N-Heptadecanoyl-lactosylceramide N-C17:0-Lactosylceramide; Lactosylceramide with C17:0 fatty acid side chain $C_{47}H_{89}NO_{13}$	1 mg
	Source: semisynthetic, bovine buttermilk Mol. Wt.: 876 Purity: 98+% by TLC, MS Appearance: solid Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C	
1629 1629-001	N-Hexanoyl-NBD-lactosylceramide N-Hexanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C6:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide $C_{42}H_{69}N_5O_{16}$	50 ug 1 mg
	Source: semisynthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C	
1630 1630-001	N-Dodecanoyl-NBD-lactosylceramide N-Dodecanoyl-NBD- <i>beta</i> -D-lactosylsphingosine; N-C12:0-NBD- <i>beta</i> -D-Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide $C_{48}H_{81}N_5O_{16}$	50 µg 1 mg
	Source: semisynthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C	

Ceramide Trihexosides



Catalog number 1067

1067 1067-10	Ceramide trihexosides CTH; Gb3; Globotriaosylceramide $C_{60}H_{113}NO_{18}$ CAS#: 71965-57-6	1 mg 10 mg
	Source: natural, porcine RBC Mol. Wt.: 1137 (tetracosanoyl) Purity: 98+% by TLC Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 2:1 Storage: -20°C	
	See Table III (pg. 93-97) for fatty acid content	
1513	Ceramide trihexosides (top spot) CTH with non-hydroxy fatty acid side chain $C_{54}H_{101}NO_{18}$	0.5 mg
	Source: natural, porcine RBC Mol. Wt.: 1052 (stearoyl) Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C	
1514	Ceramide trihexosides (bottom spot) CTH with hydroxy fatty acid side chain $C_{54}H_{101}NO_{19}$	0.5 mg
	Source: natural, porcine RBC Mol. Wt.: 1068 (2-hydroxystearoyl) Purity: 98+% by TLC Appearance: solid Solubility: DMSO, hot methanol, chloroform/methanol, 1:1 Storage: -20°C	

1520 **lyso-Ceramide trihexoside** **1 mg**
lyso-CTH; *lyso*-Globotriaosylsphingosine C₃₆H₆₇NO₁₇ CAS# 126550-86-5

Source: semisynthetic, porcine RBC **Mol. Wt.:** 786 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1
Storage: -20°C

1523 **N-Heptadecanoyl-ceramide trihexoside** **0.5 mg**
N-C17:0-Ceramide trihexoside; N-Heptadecanoyl globotriaosylceramide
C₅₃H₉₉NO₁₈

Source: semisynthetic, porcine RBC **Mol. Wt.:** 1038 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, hot methanol, chloroform/methanol, 2:1
Storage: -20°C

1524 **N-Tricosanoyl-ceramide trihexoside** **0.5 mg**
N-C23:0-Ceramide trihexoside; N-Tricosanoyl globotriaosylceramide
C₅₉H₁₁₁NO₁₈

Source: semisynthetic, porcine RBC **Mol. Wt.:** 1122 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, hot methanol, chloroform/methanol, 2:1
Storage: -20°C


1631 **N-Dodecanoyl-NBD-ceramide trihexoside** **100 µg**
1631-001 N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide C₅₄H₉₁N₅O₂₁ **1 mg**

Source: semisynthetic, porcine RBC **Mol. Wt.:** 1145 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, hot methanol, chloroform/methanol, 2:1
Storage: -20°C

1537 **N-Octadecanoyl-D₃-ceramide trihexoside, deuterated** **0.5 mg**
N-C18:0-D₃-CTH, deuterated; N-C18:0-D₃-Gb3, deuterated; N-Octadecanoyl-D₃-globotriaosylceramide, deuterated; N-Stearoyl-D₃-ceramide trihexoside, deuterated C₅₄H₉₈D₃NO₁₈

Source: semisynthetic, porcine RBC **Mol. Wt.:** 1055 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, chloroform/methanol, 2:1 **Storage:** -20°C

Globosides



Catalog number 1068

1068 **Globosides** **5 mg**
Gb4; Globotetrahexosylceramide C₆₈H₁₂₆N₂O₂₃ CAS#: 11034-93-8

Source: natural, porcine RBC **Mol. Wt.:** 1340 (tetracosanoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, hot methanol, chloroform/methanol, 2:1
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

Stable Isotopes Labeled Glycolipids

1914 **N-Octadecanoyl-D₃₅-psychosine, perdeuterated** **5 mg**

N-C18:0-D₃₅-Cerebrosides, perdeuterated; N-Stearoyl-D₃₅-psychosine, perdeuterated C₄₂H₄₆D₃₅NO₈

Source: semisynthetic, bovine **Mol. Wt.:** 762 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, hot ethanol, chloroform/methanol, 2:1
Storage: -20°C

Deuterium labeled stearoyl sidechain

1533 **N-Hexadecanoyl-D₃-glucopsychosine, deuterated** **1 mg**

N-C16:0-D₃-Glucopsychosine, deuterated; N-C16:0-D₃-Glucocerebroside, deuterated; N-Palmitoyl-D₃-glucopsychosine, deuterated C₄₀H₇₄D₃NO₈

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 703 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1534 **N-Hexadecanoyl-D₃-lactosylceramide, deuterated** **1 mg**

N-C16:0-D₃-Lactosylceramide, deuterated; N-Palmitoyl-D₃-lactosylceramide, deuterated C₄₆H₈₄D₃NO₁₃

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 865 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 5:1:0.1
Storage: -20°C

1536 **N-Octadecanoyl-D₃-sulfatide, deuterated** **1 mg**

N-C18:0-D₃-Sulfatide, deuterated; N-Stearoyl-D₃-sulfatide, deuterated C₄₂H₇₈D₃NO₁₁S

Source: semisynthetic, bovine **Mol. Wt.:** 811 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1
Storage: -20°C

1537 **N-Octadecanoyl-D₃-ceramide trihexoside, deuterated** **0.5 mg**

N-C18:0-D₃-CTH, deuterated; N-C18:0-D₃-Gb3, deuterated; N-Octadecanoyl-D₃-globotriaosylceramide, deuterated; N-Stearoyl-D₃-ceramide trihexoside, deuterated C₅₄H₉₈D₃NO₁₈

Source: semisynthetic, porcine **Mol. Wt.:** 1055 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, chloroform/methanol, 2:1 **Storage:** -20°C

Fluorescent Compounds

1621 **N-Hexanoyl-NBD-galactosylceramide** **100 µg**

1621-001 N-C6:0-NBD-*beta*-D-Galactosylsphingosine; N-C6:0-NBD-Cerebroside
C₃₆H₅₉N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1
Storage: -20°C

1633 **N-Dodecanoyl-NBD-galactosylceramide** **100 µg**
1633-001 N-C12:0-NBD-*beta*-D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside **1 mg**
C₄₂H₇₁N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 822 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, DMSO, chloroform/methanol, 2:1
Storage: -20°C

1622 **N-Hexanoyl-NBD-glucosylceramide** **100 µg**
1622-001 N-C6:0-NBD-*beta*-D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide **1 mg**
C₃₆H₅₉N₅O₁₁

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1 **Storage:** -20°C

1629 **N-Hexanoyl-NBD-lactosylceramide** **50 µg**
1629-001 N-Hexanoyl-NBD-*beta*-D-lactosylsphingosine; N-C6:0-NBD-*beta*-D-
Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide C₄₂H₆₉N₅O₁₆ **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 900 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1630 **N-Dodecanoyl-NBD-lactosylceramide** **50 µg**
1630-001 N-Dodecanoyl-NBD-*beta*-D-lactosylsphingosine; N-C12:0-NBD-*beta*-D-
Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide C₄₈H₈₁N₅O₁₆ **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 984 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1631 **N-Dodecanoyl-NBD-ceramide trihexoside** **100 µg**
1631-001 N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide C₅₄H₉₁N₅O₂₁ **1 mg**

Source: semisynthetic, porcine RBC **Mol. Wt.:** 1145 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, hot methanol, chloroform/methanol, 2:1
Storage: -20°C

1632 **N-Dodecanoyl-NBD-sulfatide** **100 µg**
1632-001 N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD-*lyso*-sulfatide; N-Dodecanoyl-
NBD-sphingosyl-*beta*-D-galactoside-3-sulfate C₄₂H₇₁N₅O₁₄S **1 mg**


Source: semisynthetic, bovine **Mol. Wt.:** 901 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

**See Biochemicals and Reagents section (page 88) for additional
fluorescent labeled products.**

**Compounds with fluorescent labels other than NBD are available on
custom basis. Contact Technical service for more information.**

Gangliosides

The diagram below can be used with the general formulas given in the ganglioside descriptions to construct the individual structures.



1064 **Gangliotetraosylceramide** **1 mg**
 Asialo GM₁; Gg4 C₆₂H₁₁₄N₂O₂₃ CAS#: 71012-19-6

Source: semisynthetic, bovine **Mol. Wt.:** 1256 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,4
 See Table III (pg. 93-97) for fatty acid content

1512 **Gangliotriosylceramide** **100 µg**
 Asialo GM₂; Gg3 C₅₆H₁₀₄N₂O₁₈

Source: semisynthetic, human **Mol. Wt.:** 1093 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3

1061 **Monosialoganglioside GM₁ (NH₄⁺ salt)** **5 mg**
1061-50 GM₁ C₇₃H₁₃₁N₃O₃₁•NH₃ CAS#: 37758-47-7 **50 mg**

Source: natural, bovine **Mol. Wt.:** 1547 + NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,4,5
 See Table III (pg. 93-97) for fatty acid content

1518 ***lyso*-Monosialoganglioside GM₁ (NH₄⁺ salt)** **500 µg**
lyso-GM₁ C₅₅H₉₇N₃O₃₀•NH₃ CAS#: 171483-40-2

Source: semisynthetic, bovine **Mol. Wt.:** 1280 +NH₃ **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.2
Storage: -20°C

1526 **Fucosylated monosialoganglioside GM₁ (NH₄⁺ salt)** **500 µg**
Fucosyl-GM₁ C₇₉H₁₄₁N₃O₃₅•NH₃ CAS#: 71812-11-8

Source: natural, porcine **Mol. Wt.:** 1693 + NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

2050 ***N-omega*-CD₃-Octadecanoyl monosialoganglioside GM₁ (NH₄⁺ salt)** **0.5 mg**
N-CD₃-Stearoyl GM₁, C₇₃H₁₂₈N₃O₃₁D₃•NH₃

Source: semisynthetic, bovine **Mol. Wt.:** 1550 + NH₃ **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

1502 **Monosialoganglioside GM₂ (NH₄⁺ salt)** **500 µg**
GM₂ C₆₇H₁₂₁N₃O₂₆•NH₃ CAS#: 19600-01-2

Source: natural, human Tay-Sachs **Mol. Wt.:** 1385+ NH₃ (stearoyl) **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,5
See Table III (pg. 93-97) for fatty acid content

2051 ***N-omega*-CD₃-Octadecanoyl monosialoganglioside GM₂ (NH₄⁺ salt)** **250 µg**
N-CD₃-Stearoyl GM₂ C₆₇H₁₁₈D₃N₃O₂₆•NH₃

Source: semisynthetic, human Tay-Sachs **Mol. Wt.:** 1388 + NH₃ **Purity:** 98+% by TLC, MS **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

1503 **Monosialoganglioside GM₃ (NH₄⁺ salt)** **1 mg**
GM₃ C₆₄H₁₁₈N₂O₂₁•NH₃ CAS#: 54827-14-4

Source: natural, bovine buttermilk **Mol. Wt.:** 1252+ NH₃ (tricosanoyl) **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol, 2:1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,5
See Table III (pg. 93-97) for fatty acid content

2052 ***N-omega*-CD₃-Octadecanoyl monosialoganglioside GM₃ (NH₄⁺ salt)** **250 µg**
N-CD₃-Stearoyl GM₃ C₅₉H₁₀₅D₃N₂O₂₁•NH₃

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 1185 + NH₃ **Purity:** 98+% by TLC, MS **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.2; forms micellar solution in water **Storage:** -20°C

1535 **Monosialoganglioside GM₄, egg (NH₄⁺ salt)** **0.5 mg**
GM₄ C₅₇H₁₀₆N₂O₁₇•NH₃ CAS#: 66456-69-7

Source: natural, egg, chicken **Mol. Wt.:** 1091+NH₃ (2-hydroxydocosanoyl)
Purity: 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol, 2:1;
forms micellar solution in water **Storage:** -20°C

General formula: 2,5
See Table III (pg. 93-97) for fatty acid content

1062 **Disialoganglioside GD_{1a} (NH₄⁺ salt)** **5 mg**
GD_{1a} C₈₄H₁₄₈N₄O₃₉•2NH₃ CAS#: 12707-58-3

Source: natural, bovine **Mol. Wt.:** 1838 + 2NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar
solution in water **Storage:** -20°C

General formula: 1,2,3,4,5,7
See Table III (pg. 93-97) for fatty acid content

1501 **Disialoganglioside GD_{1b} (NH₄⁺ salt)** **1 mg**
GD_{1b} C₈₄H₁₄₈N₄O₃₉•2NH₃ CAS#: 19553-76-5

Source: natural, bovine **Mol. Wt.:** 1838+ 2NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar
solution in water **Storage:** -20°C

General formula: 1,2,3,4,5,6
See Table III (pg. 93-97) for fatty acid content

1527 **Disialoganglioside GD₂ (NH₄⁺ salt)** **0.5 mg**
GD₂ C₇₈H₁₃₈N₄O₃₄•2NH₃ CAS#: 65988-71-8

Source: semisynthetic, rabbit **Mol. Wt.:** 1676 + 2NH₃ (stearoyl) **Purity:** 98+% by
TLC, MS **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1;
forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,5,6

1504 **Disialoganglioside GD₃ (NH₄⁺ salt)** **5 mg**
GD₃ C₇₅H₁₃₅N₃O₂₉•2NH₃ CAS#: 62010-37-1

Source: natural, bovine buttermilk **Mol. Wt.:** 1543+2NH₃ (tricosanoyl) **Purity:**
98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol, 2:1; forms
micellar solution in water **Storage:** -20°C

General formula: 1,2,5,6
See Table III (pg. 93-97) for fatty acid content

1063 **Trisialoganglioside GT_{1b} (NH₄⁺ salt)** **5 mg**
GT_{1b} C₉₅H₁₆₅N₅O₄₇•3NH₃ CAS#: 59247-13-1

Source: natural, bovine **Mol. Wt.:** 2129 + 3NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar
solution in water **Storage:** -20°C

General Formula: 1,2,3,4,5,6,7
See Table III (pg. 93-97) for fatty acid content

1516 **Tetrasialoganglioside GQ_{1b} (NH₄⁺ salt)** **100 µg**
GQ_{1b} C₁₀₆H₁₈₂N₆O₅₅•4NH₃ CAS#: 68652-37-9

Source: natural, bovine **Mol. Wt.:** 2421+4NH₃ (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,4,5,6,7,8
See Table III (pg. 93-97) for fatty acid content

1065 **Mixed Gangliosides, purified, bovine (NH₄⁺ salt)** **25 mg**
Mixed Gangliosides

Source: natural, bovine **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water
Storage: -20°C

Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides
See Table III (pg. 93-97) for fatty acid content

1525 **Mixed Gangliosides, purified, porcine, (NH₄⁺ salt)** **25 mg**

Source: natural, porcine **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water
Storage: -20°C

Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides

Glycosphingolipid Reference Mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505 **Neutral Glycosphingolipid Mixture** **1 mg/ml, 1 ml**

Source: natural, bovine and porcine **Appearance:** liquid
Solvent: chloroform/methanol, 2:1 **Solubility:** chloroform/methanol, 2:1
Storage: -20°C

Contains: cerebrosides, lactosylceramides, ceramide trihexosides, globosides

1508 **Monosialoganglioside Mixture** **0.5 mg/ml, 1 ml**

Source: natural, bovine, human **Appearance:** liquid
Solvent: chloroform/methanol/DI water, 2:1:0.1 **Solubility:** chloroform/methanol/DI water, 2:1:0.1 **Storage:** -20°C

Contains: GM₃, GM₂, GM₁

1509 **Disialoganglioside Mixture** **0.5 mg/ml, 1 ml**

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/DI water, 2:1:0.1 **Solubility:** chloroform/methanol/DI water, 2:1:0.1 **Storage:** -20°C

Contains: GD₃, GD_{1a}, GD_{1b}

1510 **Lactosylceramide and Sialosyl Derivatives Mixture** **0.5 mg/ml, 1 ml**

Source: natural, bovine buttermilk **Appearance:** liquid
Solvent: chloroform/methanol/DI water, 2:1:0.1 **Solubility:** chloroform/methanol/DI water, 2:1:0.1 **Storage:** -20°C

Contains: LC, GM₃, GD₃

1511 **Gangliotetraosylceramide and Sialosyl Derivatives Mixture** **0.5 mg/ml, 1 ml**

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/DI water, 2:1:0.1 **Solubility:** chloroform/methanol/DI water, 2:1:0.1 **Storage:** -20°C

Contains: asialo GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b}

Antibodies Directed Against Glycolipids

These monoclonal and polyclonal antibodies are directed against the carbohydrate chains of Matreya's glycolipids. The same carbohydrate moieties are found on many glycoproteins. The antibodies are for use in ELISA or TLC immunoblotting applications (9). All antibodies are quality tested by actual performance in ELISA and TLC immunoblotting. The antibodies contain no preservatives and are shipped on dry ice.

See Literature References on page 99.

1977 **Anti-ganglioside GD₃** **50 µl**

Monoclonal antibody to GD₃, isotype IgG/IgM

Source: natural, mouse hybridoma R-24 cell line **Appearance:** liquid
Solubility: DI water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for TLC immunoblotting, ELISA

1950 **Anti-ganglioside asialo GM₁** **100 µl**

Polyclonal antibody to asialo GM₁, isotype IgG/IgM

Source: natural, rabbit **Appearance:** liquid **Solubility:** DI water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to GM₁

1951 **Anti-ganglioside asialo GM₂** **50 µl**

Polyclonal antibody to asialo GM₂, isotype IgG/IgM

Source: natural, rabbit **Appearance:** liquid **Solubility:** DI water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting

1954 **Anti-ganglioside GM₁** **100 µl**

Polyclonal antibody to GM₁, isotype IgG/IgM

Source: natural, rabbit **Appearance:** liquid **Solubility:** DI water **Storage:** -20°C
Dry Ice Charge Applies

Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to asialo-GM₁

1961	Anti-ganglioside GM₂ (NANA) Polyclonal antibody to GM ₂ (NANA), isotype IgG/IgM	50 µl
	Source: natural, rabbit Appearance: liquid Solubility: DI water Storage: -20°C Dry Ice Charge Applies	
	Suitable for ELISA, TLC-immunoblotting	
1957	Anti-ganglioside GM₄ Polyclonal antibody to GM ₄ , isotype IgG/IgM	50 µl
	Source: natural, rabbit Appearance: liquid Solubility: DI water Storage: -20°C Dry Ice Charge Applies	
	Suitable for ELISA, TLC-immunoblotting	
1963	Anti-ganglioside GD₂ Polyclonal antibody to GD ₂ , isotype IgG/IgM	50 µl
	Source: natural, rabbit Appearance: liquid Solubility: DI water Storage: -20°C Dry Ice Charge Applies	
	Suitable for ELISA, TLC-immunoblotting	
1964	Anti-ganglioside GD_{1b} Polyclonal antibody to GD _{1b} , isotype IgG/IgM	50 µl
	Source: natural, rabbit Appearance: liquid Solubility: DI water Storage: -20°C Dry Ice Charge Applies	
	Suitable for ELISA, TLC-immunoblotting	
1960	Anti-globoside GL-4 Polyclonal antibody to GL-4, isotype IgG/IgM	50 µl
	Source: natural, rabbit Appearance: liquid Solubility: DI water Storage: -20°C Dry Ice Charge Applies	
	Suitable for ELISA, TLC-immunoblotting	

Enzyme Inhibitors


Ceramide: UDPglucose Transferase. PDMP (D,L-*threo*-1-phenyl-2-decanoylamino-3-morpholino-1-propanol-HCl) closely resembles the natural sphingolipid substrate of brain glucosyl transferase and is a very potent and competitive inhibitor of the enzyme (26). It has been shown to block outgrowth of neurites in cultured retina and to block glucolipid synthesis in cultured 3T3 cells (27). N.S. Radin and co-workers have shown (28) that PPMP has activity equivalent to that of PDMP when cell homogenates and brain and liver microsomes are used, but it is about 20 times more potent when used with intact cells. In another paper (29), Radin's group has shown that PDMP has substantial activity against Ehrlich ascites tumors in mice. Recent publications from the laboratory of Myles Cabot (30, 31) show that PPMP can reverse multi-drug resistance in cancer cells by causing a build-up of ceramide and preventing the synthesis of glycosylated ceramides. **See Literature References on page 99.**

Matreya also offers the resolved D- and L-*threo*-isomers of PDMP and PPMP.

Protein Kinase C Inhibitor. Sphingosine is a potent and reversible inhibitor of protein kinase C (32); it also has been shown at low concentrations to stimulate DNA synthesis and act synergistically with known growth factors (33). Note that Safingol (our L-*threo*-dihydrosphingosine) has also been shown to partially reverse multi-drug resistance in cancer cells (31) *via* inhibition of protein kinase C.

Dihydroceramide desaturase Inhibitor. Cyclopropenylceramide is the first known inhibitor of this enzyme and may allow significant studies on the role of ceramide in apoptosis. Matreya is the only source for this inhibitor. (34)

Ceramidase Inhibitors. N-Oleylethanolamine has been shown to be an efficacious inhibitor of the ceramidase found in human kidney and cerebellum (35). It is specifically an inhibitor of acid ceramidase (36) with an IC₅₀ of ca. 500 μM. N-Hexadecanoylethanolamine can be used as an inactive control. D-MAPP is a potent (IC₅₀ approximately 5 μM) inhibitor of alkaline ceramidase. Its enantiomer L-MAPP is inactive as an inhibitor and acts as a substrate for this enzyme (36, 37). **See Literature References on page 99.**



Catalog number 1751

1751 N-Oleylethanolamine 100 mg
NOE C₂₀H₃₉NO₂ CAS#: 111-58-0

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, ethyl ether, DMSO
Storage: -20°C

Activity: acid ceramidase inhibitor


1786 N-Hexadecanoylethanolamine 100 mg
C₁₈H₃₇NO₂ CAS# 544-31-0

Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, DMSO
Storage: -20°C

Activity: inactive as acid ceramidase inhibitor

1807 L-threo-Dihydrosphingosine (Safingol) 5 mg
1807-025 L-threo-Sphinganine, C18 chain C₁₈H₃₉NO₂ CAS#: 15639-50-6 25 mg

Source: synthetic **Mol. Wt.:** 301 **Melting Point (°C):** 103-114 **Purity:** 98+% by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C



Catalog number 1859

1859 **D-MAPP** **100 mg**

D-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol $C_{23}H_{39}NO_2$
 CAS#: 143492-39-1

Source: synthetic **Mol. Wt.:** 361 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol **Storage:** $-20^{\circ}C$


Activity: alkaline ceramidase inhibitor

1860 **L-MAPP** **100 mg**

L-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol $C_{23}H_{39}NO_2$
 CAS#: 143492-38-0

Source: synthetic **Mol. Wt.:** 361 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol **Storage:** $-20^{\circ}C$

Activity: inactive as alkaline ceramidase inhibitor



Catalog number: 1886

1886 **N-C8:0-Cyclopropenylceramide** **1 mg**
1886-005 **5 mg**

N-C8:0-CPPC; N-[(1R, 2S)-2-Hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] octanamide; GT₁₁ $C_{27}H_{51}NO_3$

Source: synthetic **Mol. Wt.:** 438 **Melting Point (°C):** 69-70 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol
Storage: $-20^{\circ}C$


Activity: Dihydroceramide desaturase inhibitor

1887 **N-C16:0-Cyclopropenylceramide** **1 mg**
1887-005 **5 mg**

N-C16:0-CPPC; N-[(1R, 2S)-2-Hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] hexadecanamide $C_{35}H_{67}NO_3$

Source: synthetic **Mol. Wt.:** 550 **Melting Point (°C):** 156-157 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol
Storage: $-20^{\circ}C$

Activity: Dihydroceramide desaturase inhibitor



Catalog number 1889

1889 **Conduritol B Epoxide** **25 mg**

$C_6H_{10}O_5$ CAS#: 6090-95-5

Source: synthetic **Mol. Wt.:** 162 **Melting Point (°C):** 164-166 **Purity:** 98+% by TLC, NMR **Appearance:** solid **Solubility:** DI water, DMSO, methanol (slightly) **Storage:** -20°C

Inhibits α - and β -glucosidase activity; specific inhibitor of glucocerebrosidase in cultured cells.

1719 **D,L-threo-PDMP** **100 mg**

D,L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
 $C_{23}H_{38}N_2O_3 \cdot HCl$ CAS#: 80938-69-8

Source: synthetic **Mol. Wt.:** 427 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** ethanol, methanol, chloroform, DMSO **Storage:** -20°C


Activity: glucosylceramide synthase inhibitor

1720 **D,L-threo-PPMP** **100 mg**

D,L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
 $C_{29}H_{50}N_2O_3 \cdot HCl$ CAS#: 149022-18-4

Source: synthetic **Mol. Wt.:** 511 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** alcohols, chloroform **Storage:** -20°C

Activity: glucosylceramide synthase inhibitor



Catalog number 1749

1749 **L-threo-PDMP** **10 mg**

L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
 $C_{23}H_{38}N_2O_3 \cdot HCl$ CAS#: 109836-81-9

Source: synthetic **Mol. Wt.:** 427 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** ethanol, methanol **Storage:** -20°C

1753 **D,L-erythro-PPMP** **100 mg**
D,L-erythro-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
C₂₉H₅₀N₂O₃•HCl

Source: synthetic **Mol. Wt.:** 511 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** -20°C


1755 **D,L-erythro-PDMP** **100 mg**
D,L-erythro-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
C₂₃H₃₈N₂O₃•HCl **CAS#:** 109760-77-2

Source: synthetic **Mol. Wt.:** 427 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol, DMSO **Storage:** -20°C

1756 **D-threo-PDMP** **10 mg**
D-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl
C₂₃H₃₈N₂O₃•HCl **CAS#:** 109836-82-0

Source: synthetic **Mol. Wt.:** 427 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol, methanol **Storage:** -20°C

Activity: glucosylceramide synthase inhibitor



Catalog number 1865

1865 **D-threo-PPMP** **10 mg**
D-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
C₂₉H₅₀N₂O₃•HCl

Source: synthetic **Mol. Wt.:** 511 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol, methanol **Storage:** -20°C


Activity: glucosyl ceramide synthase inhibitor

1868 **L-threo-PPMP** **10 mg**
L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl
C₂₉H₅₀N₂O₃•HCl

Source: synthetic **Mol. Wt.:** 511 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: ethanol, methanol **Storage:** -20°C

1800 **Castanospermine** **25 mg**
1,6,7,8-Tetrahydroxyoctahydroindolizine C₈H₁₅NO₄ **CAS#:** 79831-76-8

Source: natural, plant **Mol. Wt.:** 189 **Melting Point (°C):** 210-215 **Purity:** 98+%
by TLC, NMR **Appearance:** solid **Solubility:** DI water, methanol/DI water, 90:10
Storage: -20°C



Catalog number 1717

1717 **2-Fluoropalmitic acid** **25 mg**

$C_{16}H_{31}FO_2$ CAS#: 89270-22-4

Source: synthetic Mol. Wt.: 274 Melting Point (°C): 83-85 Purity: 98+% by TLC
 Appearance: solid Solubility: chloroform, ethanol, methanol Storage: -20°C

Activity: Acyl-CoA synthase inhibitor

1718 **Methyl 2-fluoropalmitate** **10 mg**

$C_{17}H_{33}FO_2$

Source: synthetic Mol. Wt.: 288 Melting Point (°C): 36-38 Purity: 98+% by TLC
 Appearance: solid Solubility: chloroform, ethanol, methanol Storage: -20°C

Activity: inactive ester of 2-fluoropalmitic acid

1750 **2,2-Difluoropalmitic acid** **25 mg**


$C_{16}H_{30}F_2O_2$

Source: synthetic Mol. Wt.: 292 Melting Point (°C): 50.8-53 Purity: 98+% by
 TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol
 Storage: -20°C

1858 **2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole** **1 mg**

THI $C_9H_{14}N_2O_5$ CAS#: 94944-70-4

Source: synthetic Mol. Wt.: 230 Melting Point (°C): n/a Purity: 99% by HPLC,
 MS, NMR Appearance: solid Solubility: DI water Storage: -20°C



Catalog number 1752

1752 **EST** **5 mg**

E-64-d; Loxastatin $C_{17}H_{30}N_2O_5$ CAS#: 88321-09-9



Source: synthetic Mol. Wt.: 342 Melting Point (°C): 125-127 Purity: 98+% by
 TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol
 Storage: -20°C

Activity: cystein protease inhibitor

Glycerolipids

Glycerophospholipids

Natural Phospholipids



1044 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC $C_{44}H_{84}NO_8P$ **CAS#:** 8002-43-5

Source: natural, chicken, egg **Mol. Wt.:** 787 (oleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether, ethanol
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1070 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC $C_{44}H_{84}NO_8P$ **CAS#:** 8002-43-5

Source: natural, bovine **Mol. Wt.:** 787 (oleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1302 **Lecithin** **50 mg/ml, 1 ml**
Phosphatidylcholine; PC $C_{44}H_{80}NO_8P$ **CAS#:** 8002-43-5

Source: natural, plant **Mol. Wt.:** 782 (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1046 **lyso-Lecithin** **50 mg**
lyso-Phosphatidylcholine $C_{24}H_{50}NO_7P$ **CAS#:** 9008-30-4

Source: semisynthetic, chicken, egg **Mol. Wt.:** 496 (palmitoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1047 **Phosphatidylserine** **50 mg/ml, 1 ml**
PS $C_{42}H_{78}NO_{10}P$

Source: natural, bovine **Mol. Wt.:** 788 (oleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, toluene
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1048 **Phosphatidylinositol (Na⁺ salt)** **10 mg/ml, 1 ml**
PI C₄₅H₇₈O₁₃P•Na CAS# 383907-36-6

Source: natural, plant **Mol. Wt.:** 858 + Na (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C

See Table III (pg. 93-97) for fatty acid content

1336 **Phosphatidylinositol, plant, soy, (Na salt)** **50 mg/ml, 1 ml**
C₄₆H₈₀O₁₃P•Na CAS# 383907-36-6

Source: natural, plant, soy **Mol. Wt.:** 872 +Na (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform, ethyl ether
Storage: -20°C

1053 **Phosphatidic acid (NH₄⁺ salt)** **50 mg**
PA C₃₉H₇₂O₈P•NH₄⁺

Source: semisynthetic, chicken, egg **Mol. Wt.:** 718 (oleoyl, NH₄⁺) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethyl ether **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1045 **Phosphatidylethanolamine** **50 mg/ml, 1 ml**
PE C₄₁H₇₈NO₈P CAS#: 39382-08-6

Source: natural, chicken, egg **Mol. Wt.:** 744 (oleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1069 **Phosphatidylethanolamine** **50 mg/ml, 1 ml**
PE C₄₁H₇₈NO₈P CAS#: 90989-93-8

Source: natural, bovine **Mol. Wt.:** 744 (oleoyl) **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform **Storage:** -20°C

1301 **Phosphatidylethanolamine** **50 mg/ml, 1 ml**
PE C₄₁H₇₄NO₈P CAS#: 90989-93-8

Source: natural, plant **Mol. Wt.:** 740 (linoleoyl) **Purity:** 98+% by TLC
Appearance: liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

See Table III (pg. 93-97) for fatty acid content

1052 **Phosphoglycerides Kit** **1 each**


Source: natural, chicken egg, bovine, plant **Purity:** 98+% by TLC **Appearance:** liquid/solid
Solvent: various **Storage:** -20°C

Individually packed in ampules and vials (Purity 98+%): Phosphatidic acid NH₄⁺ salt 10mg; Phosphatidylethanolamine, egg (in 1 ml CHCl₃) 10mg; Sphingomyelin, bovine 10mg; Phosphatidylserine, bovine (in 1 ml CHCl₃) 10 mg; Lecithin, egg (in 1 ml CHCl₃) 10 mg; lyso-Lecithin, egg 10 mg; Cerebrosides, bovine 10mg; Sulfatides, bovine 10mg; Phosphatidylinositol, Na⁺ salt, plant (in 1 ml CHCl₃) 3mg

Synthetic Phospholipids

These phospholipids have 98+% chemical purity except where stated and 99% fatty acid chain purity. Store at -20°C. Solubility: see individual entries

Phosphatidic acid derivatives



1428 **1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt)** **100 mg**
DMPA C₃₁H₆₀O₈P•Na CAS#: 80724-31-8

Source: synthetic Mol. Wt.: 615 Purity: 98+% by TLC Appearance: solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C

1429 **1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt)** **100 mg**
DPPA C₃₅H₆₈O₈P•Na CAS#: 71065-87-7

Source: synthetic Mol. Wt.: 671 Purity: 98+% by TLC Appearance: solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C


1430 **1,2-Distearoyl-sn-glycero-3-phosphatidic acid (Na⁺ salt)** **100 mg**
DSPA C₃₉H₇₆O₈P•Na CAS#: 108321-18-2

Source: synthetic Mol. Wt.: 727 Purity: 98+% by TLC Appearance: solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C

Phosphatidylcholines

1442 **1,2-Dilauroyl-sn-glycero-3-phosphorylcholine** **100 mg**
DLPC C₃₂H₆₄NO₈P CAS#: 18194-25-7

Source: synthetic Mol. Wt.: 622 Purity: 98+% by TLC Appearance: solid
Solubility: methylene chloride, methanol Storage: -20°C



1425 **1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine** **100 mg**
 DMPC C₃₆H₇₂NO₈P CAS#: 18194-24-6

Source: synthetic Mol. Wt.: 678 Purity: 98+% by TLC Appearance: solid
 Melting Point: 130-139°C Solubility: methylene chloride, methanol Storage: -20°C

1426 **1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine** **100 mg**
 DPPC C₄₀H₈₀NO₈P CAS#: 63-89-8

Source: synthetic Mol. Wt.: 734 Purity: 98+% by TLC Appearance: solid
 Solubility: methylene chloride, methanol Storage: -20°C

1400 **1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine** **50 mg**
 DHDPC C₄₂H₈₄NO₈P CAS#: 70897-27-7

Source: synthetic Mol. Wt.: 762 Purity: 98+% by TLC Appearance: solid
 Solubility: methylene chloride, methanol Storage: -20°C

1427 **1,2-Distearoyl-sn-glycero-3-phosphorylcholine** **100 mg**
 DSPC C₄₄H₈₈NO₈P CAS#: 816-94-4

Source: synthetic Mol. Wt.: 790 Purity: 98+% by TLC Appearance: solid
 Solubility: methylene chloride, methanol Storage: -20°C

1437 **1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine** **100 mg**
 POPC C₄₂H₈₂NO₈P CAS#: 26853-31-6

Source: synthetic Mol. Wt.: 760 Purity: 98+% by TLC Appearance: solid
 Solubility: methylene chloride, methanol Storage: -20°C

1445 **1-Palmitoyl-sn-glycero-3-phosphorylcholine** **100 mg**
lyso-PPC C₂₄H₅₀NO₇P CAS#: 17364-16-8

Source: synthetic Mol. Wt.: 496 Purity: 98+% by TLC Appearance: solid
 Solubility: methylene chloride, methanol Storage: -20°C


1409 **1-Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine** **25 mg/ml, 1ml**
 C₄₄H₈₄NO₈P

Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid
 Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C

1410 **1-Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine** **25 mg/ml, 1ml**
 $C_{44}H_{84}NO_8P$
Source: synthetic **Mol. Wt.:** 786 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform, ethanol **Storage:** -20°C

1411 **1-Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine** **25 mg/ml, 1ml**
 $C_{44}H_{84}NO_8P$
Source: synthetic **Mol. Wt.:** 786 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: chloroform **Solubility:** chloroform, ethanol **Storage:** -20°C

Phosphatidylglycerols



1443 **1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt)** **100 mg**
DLPG $C_{30}H_{58}O_{10}P \cdot Na$ **CAS#:** 73548-69-3
Source: synthetic **Mol. Wt.:** 632 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol, 5:1 **Storage:** -20°C

1431 **1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt)** **100 mg**
DMPG $C_{34}H_{66}O_{10}P \cdot Na$ **CAS#:** 200880-40-6
Source: synthetic **Mol. Wt.:** 689 **Purity:** 98+% by TLC **Appearance:** solid
Melting Point: 120-129°C **Solubility:** chloroform/methanol, 5:1 **Storage:** -20°C


1432 **1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt)** **100 mg**
DPPG $C_{38}H_{74}O_{10}P \cdot Na$ **CAS#:** 200880-41-7
Source: synthetic **Mol. Wt.:** 745 **Purity:** 98+% by TLC **Appearance:** solid
Melting Point: 122-127°C **Solubility:** chloroform/methanol, 5:1 **Storage:** -20°C

1433 **1,2-Distearoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt)** **100 mg**
DSPG $C_{42}H_{82}O_{10}P \cdot Na$ **CAS#:** 4537-78-4
Source: synthetic **Mol. Wt.:** 801 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol, 5:1 **Storage:** -20°C

1438 **1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol (Na⁺ salt)** **100 mg**
POPG $C_{40}H_{76}O_{10}P \cdot Na$ **CAS#:** 202070-86-8
Source: synthetic **Mol. Wt.:** 771 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol, 5:1 **Storage:** -20°C

Phosphatidylethanolamines

Catalog number 1444




1444	1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine DLPE C ₂₉ H ₅₈ NO ₈ P CAS#: 42436-56-6	100 mg
	Source: synthetic Mol. Wt.: 580 Purity: 98+% by TLC Appearance: solid Solubility: chloroform + methanol mixture Storage: -20°C	
1434	1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine DMPE C ₃₃ H ₆₆ NO ₈ P CAS# 998-07-2	100 mg
	Source: synthetic Mol. Wt.: 636 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5 Storage: -20°C	
1435	1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine DPPE C ₃₇ H ₇₄ NO ₈ P CAS#: 923-61-5	100 mg
	Source: synthetic Mol. Wt.: 692 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5 Storage: -20°C	
1436	1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine DSPE C ₄₁ H ₈₂ NO ₈ P CAS#: 1069-79-0	100 mg
	Source: synthetic Mol. Wt.: 748 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/acetic acid, 95:5; chloroform/methanol/DI water/acetic acid, 100:30:10:2.5 Storage: -20°C	
1439	1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000 (Na⁺ salt) DSPE-MPEG-2000 CAS#: 147867-65-0	100 mg
	Source: synthetic Purity: 98+% by TLC Appearance: solid Solubility: chloroform Storage: -20°C	

Phosphatidylinositols

The metabolism of inositol lipids is involved in the signal transduction of many hormones, neurotransmitters and growth factors (1, 2). In the classical pathway, phosphatidylinositol-specific phospholipase C (PI-PLC) hydrolyzes phosphatidyl 4,5-biphosphate (PIP₂) to yield 1,2-diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃). The role of IP₃ and DAG as second messengers is well recognized.

In a second, more recently discovered pathway, the activation of phosphoinositide (PI) 3-kinase results in the formation of three novel phosphatidyl (PI) lipids phosphorylated at the D3 position of the inositol ring: PI-3-P, PI-3,4-P₂ and PI-3,4,5-P₃ (3). These D3 lipids are not known substrates for any of the phospholipase C enzymes and function as second messengers. PI 3-kinase activity is correlated with many cellular processes, including the regulation of cell growth, oncogenic transformation, chemotaxis and receptor down-regulation among others (4, 5, 7). The recent paper on the effect of PI3,4-P₂ on the *Akt* proto-oncogene product (38) also contains protocols for applying PIP's to cell cultures. Matreya's synthetic phosphatidylinositols and inositol phosphates are excellent tools for investigating these second messengers, understanding the enzyme mechanisms involved in phosphoinositide metabolism (39, 40) and for designing therapeutic pharmacological agents. The compounds are evaluated by ¹H and ³¹P NMR to guarantee enantiomeric purity of >98%. **See Literature References on page 99.**



Catalog number 1779

1779 1779-1	Phosphatidylinositol, dipalmitoyl, (NH₄⁺ salt) PI; DPPI (NH ₄ ⁺ salt) C ₄₁ H ₇₈ O ₁₃ P•NH ₄ ⁺	0.5 mg 1 mg
Source: synthetic Mol. Wt.: 828 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: solid Solubility: chloroform/methanol/DI water, 5:1:0.1 Storage: -20°C		
1773 1773-1 1773-5	Phosphatidylinositol 3-phosphate, dipalmitoyl, (NH₄⁺ salt) DPPI-3-P; PI-3-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	100 µg 1 mg 5 mg
Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR, Appearance: solid Solubility: chloroform/methanol/DI water, 1:1:0.3 Storage: -20°C		
1919 1919-1 1919-5	Phosphatidylinositol 4-phosphate, dipalmitoyl, (NH₄⁺ salt) DPPI-4-P; PI-4-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	100 µg 1 mg 5 mg
Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: solid Solubility: methanol, chloroform/methanol/DI water, 1:1:0.3; slightly soluble in DI water Storage: -20°C		

1784 **Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (NH₄⁺ salt)** **100 µg**
1784-1 DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (NH₄⁺ salt) C₂₅H₄₉O₁₉P₃•5NH₄ **1 mg**
1784-5 **5 mg**

Source: synthetic **Mol. Wt.:** 831 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: solid **Solubility:** chloroform/methanol/DI water, 1:1:0.3
Storage: -20°C

1778 **Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (Na⁺ salt)** **100 µg**
1778-1 DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (Na⁺ salt) C₂₅H₄₄O₁₉P₃•5Na **1 mg**
1778-5 **5 mg**

Source: synthetic **Mol. Wt.:** 856 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: solid **Solubility:** DI water **Storage:** -20°C

1783 **Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (NH₄⁺ salt)** **100 µg**
1783-1 DPPI-3,4,5-P3; PI-3,4,5-P3 dipalmitoyl (NH₄⁺ salt) C₄₁H₇₅O₂₂P₄•7NH₄ **1 mg**
1783-5 **5 mg**

Source: synthetic **Mol. Wt.:** 1170 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: solid **Solubility:** chloroform/methanol/DI water, 1:1:0.3
Storage: -20°C

1775 **Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (Na⁺ salt)** **100 µg**
1775-1 DPPI-3,4,5-P3; PI-3,4,5-P3 dipalmitoyl (Na⁺ salt) C₄₁H₇₅O₂₂P₄•7Na **1 mg**
1775-5 **5 mg**

Source: synthetic **Mol. Wt.:** 1205 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: solid **Solubility:** DI water **Storage:** -20°C

Bacterial Tetraethers

1303 **Main phospholipid (MPL) of *Thermoplasma acidophilum*, (>95% pure)** **5 mg**

Purified MPL of *Thermoplasma acidophilum* (>95% pure) C₉₅H₁₈₈O₁₆P


Source: natural, Archaeobacteria **Mol. Wt.:** 1618 **Purity:** >95% by TLC, HPLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1; hexane/2-propanol/DI water, 30:40:5 **Storage:** 4-8°C

1303-2 **Main phospholipid (MPL) of *Thermoplasma acidophilum*, (>50% pure)** **50 mg**

MPL of *Thermoplasma acidophilum* (>50% pure) C₉₅H₁₈₈O₁₆P

Source: natural, Archaeobacteria **Mol. Wt.:** 1618 **Purity:** >50% by TLC
Appearance: liquid **Solubility:** chloroform/methanol, 2:1; hexane/2-propanol/DI water, 30:40:5 **Storage:** 4-8°C highly hygroscopic

Glycosyl Glycerides



Catalog number 1058

1058	Monogalactosyldiglyceride	10 mg
-------------	----------------------------------	--------------

MGDG (hydrogenated) $C_{45}H_{86}O_{10}$ CAS#: 41670-62-6

Source: natural, plant **Mol. Wt.:** 787 (stearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 4:1:0.1
Storage: $-20^{\circ}C$

1059	Digalactosyldiglyceride	5 mg
-------------	--------------------------------	-------------

DGDG (hydrogenated) $C_{51}H_{96}O_{15}$ CAS#: 92457-02-8

Source: natural, plant **Mol. Wt.:** 949 (distearoyl) **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 4:1:0.1
Storage: $-20^{\circ}C$

Fatty Acids

Simple Fatty Acids

Saturated Fatty Acids and Methyl Esters

These products are 99% pure by GC. They are stable at room temperature and are supplied neat.

1200	Methyl hexanoate	1 g
-------------	-------------------------	------------

Methyl caproate; C6:0 Methyl ester $C_7H_{14}O_2$ CAS#: 106-70-7

Source: natural, plant **Mol. Wt.:** 130 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, ethyl ether **Storage:** room temperature

1196	Heptanoic acid	1 g
-------------	-----------------------	------------

C7:0 Fatty acid $C_7H_{14}O_2$ CAS#: 111-14-8

Source: natural, plant **Mol. Wt.:** 130 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, ethyl ether **Storage:** room temperature

1197	Methyl heptanoate	1 g
-------------	--------------------------	------------

C7:0 Methyl ester $C_8H_{16}O_2$ CAS#: 106-73-0

Source: natural, plant **Mol. Wt.:** 144 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, ethyl ether **Storage:** room temperature

1198	Octanoic acid Caprylic acid; C8:0 Fatty acid $C_8H_{16}O_2$ CAS#: 124-07-2	1 g
	Source: natural, plant Mol. Wt.: 144 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1199	Methyl octanoate Methyl caprylate; C8:0 Methyl ester $C_9H_{18}O_2$ CAS#: 111-11-5	1 g
	Source: natural, plant Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1163	Nonanoic acid C9:0 Fatty acid; Pelargonic acid $C_9H_{18}O_2$ CAS#: 112-05-0	100 mg
	Source: synthetic Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1164	Methyl nonanoate C9:0 Methyl ester $C_{10}H_{20}O_2$ CAS#: 1731-84-6	100 mg
	Source: synthetic Mol. Wt.: 172 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1261	Methyl decanoate Methyl caprate; C10:0 Methyl ester $C_{11}H_{22}O_2$ CAS#: 110-42-9	500 mg
	Source: natural, plant Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane Storage: room temperature	
1165	Undecanoic acid C11:0 Fatty acid $C_{11}H_{22}O_2$ CAS#: 112-37-8	100 mg
	Source: synthetic Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1166	Methyl undecanoate C11:0 Methyl ester $C_{12}H_{24}O_2$ CAS#: 1731-86-8	100 mg
	Source: synthetic Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1008	Dodecanoic acid Lauric acid; C12:0 Fatty acid $C_{12}H_{24}O_2$ CAS#: 143-07-7	1 g
	Source: natural, plant Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1009	Methyl dodecanoate Methyl laurate; C12:0 Methyl ester $C_{13}H_{26}O_2$ CAS#: 111-82-0	1 g
	Source: natural, plant Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	

1161	Tridecanoic acid C13:0 Fatty acid C ₁₃ H ₂₆ O ₂ CAS#: 638-53-9	100 mg
	Source: synthetic Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1162	Methyl tridecanoate C13:0 Methyl ester C ₁₄ H ₂₈ O ₂ CAS#: 1731-88-0	100 mg
	Source: synthetic Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1010	Tetradecanoic acid Myristic acid; C14:0 Fatty acid C ₁₄ H ₂₈ O ₂ CAS#: 544-63-8	1 g
	Source: natural, plant Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1011	Methyl tetradecanoate Methyl myristate; C14:0 Methyl ester C ₁₅ H ₃₀ O ₂ CAS#: 124-10-7	1 g
	Source: natural, plant Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1012	Pentadecanoic acid C15:0 Fatty acid C ₁₅ H ₃₀ O ₂ CAS#: 1002-84-2	1 g
	Source: synthetic Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1013	Methyl pentadecanoate C15:0 Methyl ester C ₁₆ H ₃₂ O ₂ CAS#: 7132-64-1	1 g
	Source: synthetic Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1014	Hexadecanoic acid Palmitic acid; C16:0 Fatty acid C ₁₆ H ₃₂ O ₂ CAS#: 57-10-3	1 g
	Source: natural, plant Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1015	Methyl hexadecanoate Methyl palmitate; C16:0 Methyl ester C ₁₇ H ₃₄ O ₂ CAS#: 112-39-0	1 g
	Source: natural, plant Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1018	Heptadecanoic acid Margaric acid; C17:0 Fatty acid C ₁₇ H ₃₄ O ₂ CAS#: 506-12-7	1 g
	Source: synthetic Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	

1019	Methyl heptadecanoate Methyl margarate; C17:0 Methyl ester $C_{18}H_{36}O_2$ CAS#: 1731-92-6	1 g
	Source: synthetic Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1020	Octadecanoic acid Stearic acid; C18:0 Fatty acid $C_{18}H_{36}O_2$ CAS#: 57-11-4	1 g
	Source: natural, plant Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1021	Methyl octadecanoate Methyl stearate; C18:0 Methyl ester $C_{19}H_{38}O_2$ CAS#: 112-61-8	1 g
	Source: natural, plant Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1028	Nonadecanoic acid C19:0 Fatty acid $C_{19}H_{38}O_2$ CAS#: 646-30-0	100 mg
	Source: synthetic Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1029	Methyl nonadecanoate C19:0 Methyl ester $C_{20}H_{40}O_2$ CAS#: 1731-94-8	100 mg
	Source: synthetic Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1030	Eicosanoic acid Arachidic acid; C20:0 Fatty acid $C_{20}H_{40}O_2$ CAS#: 506-30-9	500 mg
	Source: natural, plant Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1031	Methyl eicosanoate Methyl arachidate; C20:0 Methyl ester $C_{21}H_{42}O_2$ CAS#: 1120-28-1	500 mg
	Source: natural, plant Mol. Wt.: 327 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1241	Heneicosanoic acid C21:0 Fatty acid $C_{21}H_{42}O_2$ CAS#: 2363-71-5	100 mg
	Source: synthetic Mol. Wt.: 326 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1242	Methyl heneicosanoate C21:0 Methyl ester $C_{22}H_{44}O_2$ CAS#: 6064-90-0	100 mg
	Source: synthetic Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	

1035	Docosanoic acid Behenic acid; C22:0 Fatty acid $C_{22}H_{44}O_2$ CAS#: 112-85-6	500 mg
	Source: natural, plant Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1036	Methyl docosanoate Methyl behenate; C22:0 Methyl ester $C_{23}H_{46}O_2$ CAS#: 929-77-1	500 mg
	Source: natural, plant Mol. Wt.: 354 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1186	Tricosanoic acid C23:0 Fatty acid $C_{23}H_{46}O_2$ CAS#: 2433-96-7	100 mg
	Source: synthetic Mol. Wt.: 355 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1187	Methyl tricosanoate C23:0 Methyl ester $C_{24}H_{48}O_2$ CAS#: 2433-97-8	100 mg
	Source: synthetic Mol. Wt.: 368 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1037	Tetracosanoic acid Lignoceric acid; C24:0 Fatty acid $C_{24}H_{48}O_2$ CAS#: 557-59-5	100 mg
	Source: synthetic Mol. Wt.: 369 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1038	Methyl tetracosanoate Methyl lignocerate; C24:0 Methyl ester $C_{25}H_{50}O_2$ CAS#: 2442-49-1	100 mg
	Source: synthetic Mol. Wt.: 382 Purity: 99% by GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1251	Hexacosanoic acid Cerotic acid; C26:0 Fatty acid $C_{26}H_{52}O_2$ CAS#: 506-46-7	25 mg
	Source: synthetic Mol. Wt.: 397 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1252	Methyl hexacosanoate Methyl cerotate; C26:0 Methyl ester $C_{27}H_{54}O_2$ CAS#: 5802-82-4	25 mg
	Source: synthetic Mol. Wt.: 411 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: room temperature	
1271	Methyl octacosanoate Methyl montanate; C28:0 Methyl ester $C_{29}H_{58}O_2$ CAS#: 55682-92-3	50 mg
	Source: synthetic Mol. Wt.: 439 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methylene chloride Storage: room temperature	

1273 **Methyl triacontanoate** **50 mg**
Methyl melissate; C30:0 Methyl ester $C_{31}H_{62}O_2$ **CAS#:** 629-83-4

Source: synthetic **Mol. Wt.:** 467 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, methylene chloride **Storage:** room temperature

1275 **Methyl dotriacontanoate** **50 mg**
Methyl lacceroate; C32:0 Methyl ester $C_{33}H_{66}O_2$ **CAS#:** 41755-79-7

Source: synthetic **Mol. Wt.:** 495 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: chloroform, methylene chloride **Storage:** room temperature

Unsaturated Fatty Acids and Methyl Esters

Unsaturated fatty acids are easily oxidized. Flush open containers with argon or nitrogen and store at $-20^{\circ}C$, in dark.

1157 **Tetradecenoic acid (*cis*-9)** **100 mg**
Myristoleic acid; C14:1 (*cis*-9) fatty acid $C_{14}H_{26}O_2$ **CAS#:** 544-64-9

Source: natural, plant **Mol. Wt.:** 226 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** $-20^{\circ}C$

1040 **Methyl tetradecenoate (*cis*-9)** **100 mg**
Methyl myristoleate; C14:1 (*cis*-9) Methyl ester $C_{15}H_{28}O_2$
CAS#: 56219-06-8

Source: natural, plant **Mol. Wt.:** 240 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** $-20^{\circ}C$

1243 **Hexadecenoic acid (*cis*-6)** **25 mg**
Sapienic acid $C_{16}H_{30}O_2$

Source: synthetic **Mol. Wt.:** 254 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** ethanol, methanol, chloroform, ethyl ether
Storage: $-20^{\circ}C$

1016 **Hexadecenoic acid (*cis*-9)** **100 mg**
Palmitoleic acid; C16:1 (*cis*-9) Fatty acid $C_{16}H_{30}O_2$ **CAS#:** 373-49-9

Source: natural, plant **Mol. Wt.:** 254 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** $-20^{\circ}C$

1017 **Methyl hexadecenoate (*cis*-9)** **100 mg**
Methyl palmitoleate; C16:1 (*cis*-9) Methyl ester $C_{17}H_{32}O_2$
CAS#: 1120-25-8

Source: natural, plant **Mol. Wt.:** 268 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** $-20^{\circ}C$

1147 **Hexadecenoic acid (*trans*-9)** **100 mg**
Palmitelaidic acid; C16:1 (*trans*-9) Fatty acid $C_{16}H_{30}O_2$
CAS#: 10030-73-6

Source: synthetic **Mol. Wt.:** 254 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** $-20^{\circ}C$

1148	Methyl hexadecenoate (<i>trans</i>-9) Methyl palmitelaidate; C16:1 (<i>trans</i> -9) Methyl ester C ₁₇ H ₃₂ O ₂ CAS#: 10030-74-7 Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1208	11-Hexadecenoic acid (92% <i>cis</i>, 8% <i>trans</i>) C16:1 (<i>cis</i> -11) Fatty acid C ₁₆ H ₃₀ O ₂ Source: synthetic Mol. Wt.: 254 Purity: >98%, by TLC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C 92% <i>cis</i> , 8% <i>trans</i> by GC	50 mg
1204	Heptadecenoic acid (<i>cis</i>-10) C17:1 (<i>cis</i> -10) Fatty acid C ₁₇ H ₃₂ O ₂ CAS#: 29743-97-3 Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1203	Methyl heptadecenoate (<i>cis</i>-10) Methyl heptadecenoate; C17:1 (<i>cis</i> -10) Methyl ester C ₁₈ H ₃₄ O ₂ CAS#: 75190-82-8 Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1022	Octadecenoic acid (<i>cis</i>-9) Oleic acid; C18:1 (<i>cis</i> -9) Fatty acid C ₁₈ H ₃₄ O ₂ CAS#: 112-80-1 Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	1 g
1023	Methyl octadecenoate (<i>cis</i>-9) Methyl oleate; C18:1 (<i>cis</i> -9) Methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 112-62-9 Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	1 g
1149	Octadecenoic acid (<i>trans</i>-9) Elaidic acid; C18:1 (<i>trans</i> -9) Fatty acid C ₁₈ H ₃₄ O ₂ CAS#: 112-79-8 Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	1 g
1150	Methyl octadecenoate (<i>trans</i>-9) Methyl elaidate; C18:1 (<i>trans</i> -9) Methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 1937-62-8 Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	1 g

1266	Octadecenoic acid (<i>cis</i>-11) <i>cis</i> -Vaccenic acid; C18:1(<i>cis</i> -11) Fatty acid C ₁₈ H ₃₄ O ₂ CAS#: 506-17-2	100 mg
	Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1267	Methyl octadecenoate (<i>cis</i>-11) Methyl <i>cis</i> -vaccenate; C18:1(<i>cis</i> -11) Methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 1937-63-9	100 mg
	Source: semisynthetic, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1262	Octadecenoic acid (<i>trans</i>-11) <i>trans</i> Vaccenic acid; C18:1 (<i>trans</i> -11) Fatty acid C ₁₈ H ₃₄ O ₂ CAS#: 693-72-1	100 mg
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1263	Methyl octadecenoate (<i>trans</i>-11) Methyl <i>trans</i> vaccenate; C18:1 (<i>trans</i> -11) Methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 6198-58-9	100 mg
	Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1024	Octadecadienoic acid (all <i>cis</i>-9,12) Linoleic acid; C18:2 (all <i>cis</i> -9,12) Fatty acid C ₁₈ H ₃₂ O ₂ CAS#: 60-33-3	1 g
	Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, ethanol, hexane Storage: -20°C	
1025	Methyl octadecadienoate (all <i>cis</i>-9,12) Methyl linoleate; C18:2 (all <i>cis</i> -9,12) Methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 112-63-0	1 g
	Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1151	Linoelaidic acid (all <i>trans</i>-9,12) C18:2 (all <i>trans</i> -9, 12) Fatty acid C ₁₈ H ₃₂ O ₂ CAS#: 506-21-8	100 mg
	Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1152	Methyl octadecadienoate (all <i>trans</i>-9,12) Methyl linoelaidate; C18:2 (all <i>trans</i> -9,12) Methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 2566-97-4	100 mg
	Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1026	Octadecatrienoic acid (all <i>cis</i>-9,12,15) Linolenic acid; C18:3 (all <i>cis</i> -9,12,15) Fatty acid C ₁₈ H ₃₀ O ₂ CAS#: 463-40-1 Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	500 mg
1027	Methyl octadecatrienoate (all <i>cis</i>-9,12,15) Methyl linolenate; C18:3 (all <i>cis</i> -9,12,15) Methyl ester C ₁₉ H ₃₂ O ₂ CAS#: 301-00-8 Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	500 mg
1153	Octadecatrienoic acid (all <i>cis</i>-6,9,12) <i>gamma</i> -Linolenic acid; C18:3 (all <i>cis</i> -6,9,12) Fatty acid C ₁₈ H ₃₀ O ₂ CAS#: 506-26-3 Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1154	Methyl octadecatrienoate (all <i>cis</i>-6,9,12) Methyl <i>gamma</i> -linolenate; C18:3 (all <i>cis</i> -6,9,12) Methyl ester C ₁₉ H ₃₂ O ₂ CAS#: 16326-32-2 Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1276	Stearidonic acid (all <i>cis</i>-6,9,12,15) Morotic acid; C18:4 (all <i>cis</i> -6,9,12,15) Fatty acid C ₁₈ H ₂₈ O ₂ CAS#: 20290-75-9 Source: natural, plant Mol. Wt.: 276 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	25 mg
1277	Methyl stearidonate (all <i>cis</i>-6,9,12,15) Morotic acid methyl ester; C18:4 (all <i>cis</i> -6,9,12,15) Methyl ester C ₁₉ H ₃₀ O ₂ CAS#: 73097-00-4 Source: natural, plant Mol. Wt.: 290 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	25 mg
1205	Nonadecenoic acid (<i>cis</i>-10) C19:1 (<i>cis</i> -10) Fatty acid C ₁₉ H ₃₆ O ₂ CAS#: 73033-09-7 Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1206	Methyl nonadecenoate (<i>cis</i>-10) C19:1 (<i>cis</i> -10) Methyl ester C ₂₀ H ₃₈ O ₂ CAS#: 19788-74-0 Source: synthetic Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg

1032	Eicosenoic acid (<i>cis</i>-11) C20:1 (<i>cis</i> -11) Fatty acid C ₂₀ H ₃₈ O ₂ CAS#: 5561-99-9 Source: natural, plant Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1033	Methyl eicosenoate (<i>cis</i>-11) Methyl eicosenoate; C20:1 (<i>cis</i> -11) Methyl ester C ₂₁ H ₄₀ O ₂ CAS#: 2390-09-2 Source: natural, plant Mol. Wt.: 324 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1192	Eicosadienoic acid (all <i>cis</i>-11,14) C20:2 (all <i>cis</i> -11,14) Fatty acid C ₂₀ H ₃₆ O ₂ CAS#: 2091-39-6 Source: synthetic Mol. Wt.: 309 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1193	Methyl eicosadienoate (all <i>cis</i>-11,14) Methyl eicosadienoate; C20:2 (all <i>cis</i> -11,14) Methyl ester C ₂₁ H ₃₈ O ₂ CAS#: 2463-02-7 Source: synthetic Mol. Wt.: 322 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1179	Methyl eicosatrienoate (all <i>cis</i>-5,8,11) Mead acid methyl ester; C20:3 (all <i>cis</i> -5,8,11) Methyl ester C ₂₁ H ₃₆ O ₂ CAS#: 14602-39-2 Source: natural, plant Mol. Wt.: 320 Purity: 90% by TLC, GC Appearance: liquid Solvent: hexane Solubility: chloroform, hexane, ethyl ether Storage: -20°C	1 mg/ml, 1 ml
1269	Methyl eicosatrienoate (all <i>cis</i>-8,11,14) Methyl homogamma linolenate; C20:3 (all <i>cis</i> -8,11,14) Methyl ester C ₂₁ H ₃₆ O ₂ Source: semi-synthetic, plant Mol. Wt.: 321 Purity: 99% by TLC, GC Appearance: liquid Solubility: hexane, ethyl ether, chloroform Storage: -20°C	50 mg
1042	Arachidonic acid (all <i>cis</i>-5,8,11,14) C20:4 (all <i>cis</i> -5,8,11,14) Fatty acid C ₂₀ H ₃₂ O ₂ CAS#: 506-32-1 Source: natural, fungal Mol. Wt.: 304 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride Storage: -20°C Dry Ice Charge Applies	100 mg
1034	Methyl eicosatetraenoate (all <i>cis</i>-5,8,11,14) Methyl arachidonate; C20:4 (all <i>cis</i> -5,8,11,14) Methyl ester C ₂₁ H ₃₄ O ₂ CAS#: 2566-89-4 Source: natural, fungal Mol. Wt.: 318 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	100 mg

1167	Eicosapentaenoic acid (all <i>cis</i>-5,8,11,14,17) EPA ; <i>omega</i> -3 Fatty acid; C20:5 (all <i>cis</i> -5,8,11,14,17) Fatty acid C ₂₀ H ₃₀ O ₂ CAS#: 10417-94-4 Source: natural, fish oil Mol. Wt.: 302 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies Anti-hyperlipoproteinemic agent; 5-LOX inhibitor	25 mg
1194	Methyl eicosapentaenoate (all <i>cis</i>-5,8,11,14,17) Methyl ester of <i>omega</i> -3 Fatty acid; C20:5 (all <i>cis</i> -5,8,11,14,17) Methyl ester C ₂₁ H ₃₂ O ₂ CAS#: 2734-47-6 Source: natural, fish oil Mol. Wt.: 316 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies	25 mg
1264	Docosenoic acid (<i>cis</i>-13) Erucic acid; C22:1 (<i>cis</i> -13) Fatty acid C ₂₂ H ₄₂ O ₂ CAS#: 112-86-7 Source: natural, plant Mol. Wt.: 339 Purity: >99% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether, hexane Storage: -20°C	100 mg
1265	Methyl docosenoate (<i>cis</i>-13) Methyl erucate; C22:1 (<i>cis</i> -13) Methyl ester C ₂₃ H ₄₄ O ₂ CAS#: 1120-34-9 Source: natural, plant Mol. Wt.: 352 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C	100 mg
1175	Docosapentaenoic acid (all <i>cis</i>-7,10,13,16,19) C22:5 (all <i>cis</i> -7,10,13,16,19) Fatty acid C ₂₂ H ₃₄ O ₂ CAS#: 24880-45-3 Source: semi-synthetic Mol. Wt.: 330 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies	25 mg
1244	Methyl docosapentaenoate (all <i>cis</i>-7,10,13,16,19) C22:5 (all <i>cis</i> -7,10,13,16,19) Methyl ester C ₂₃ H ₃₆ O ₂ CAS#: 108698-02-8 Source: semi-synthetic Mol. Wt.: 344 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, ethanol, hexane, Storage: -20°C Dry Ice Charge Applies	25 mg
1136	Docosahexaenoic acid (all <i>cis</i>-4,7,10,13,16,19) DHA; C22:6 (all <i>cis</i> -4,7,10,13,16,19) <i>omega</i> -3 Fatty acid C ₂₂ H ₃₂ O ₂ CAS#: 6217-54-5 Source: natural, algae Mol. Wt.: 328 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies	100 mg

1041 **Methyl docosaehaenoate (all *cis*-4,7,10,13,16,19)** **100 mg**
Methyl ester of *omega*-3 fatty acid; C22:6 (all *cis*-4,7,10,13,16,19) Methyl ester $C_{23}H_{34}O_2$ **CAS#:** 2566-90-7

Source: natural, algae **Mol. Wt.:** 342 **Purity:** 99% by TLC, GC
Appearance: liquid **Solubility:** chloroform, hexane, ethyl ether **Storage:** -20°C
Dry Ice Charge Applies

1155 **Nervonic acid (*cis*-15)** **100 mg**
Tetracosenoic acid (*cis*-15); C24:1 (*cis*-15) Fatty acid $C_{24}H_{46}O_2$ **CAS#:** 506-37-6

Source: synthetic **Mol. Wt.:** 367 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1156 **Methyl tetracosenoate (*cis*-15)** **100 mg**
Methyl nervonate; C24:1 (*cis*-15) Methyl ester $C_{25}H_{48}O_2$ **CAS#:** 2733-88-2

Source: synthetic **Mol. Wt.:** 381 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

Trans Fatty Acids and Methyl Esters

1147 **Hexadecenoic acid (*trans*-9)** **100 mg**
Palmitelaidic acid; C16:1 (*trans*-9) Fatty acid $C_{16}H_{30}O_2$ **CAS#:** 10030-73-6

Source: synthetic **Mol. Wt.:** 254 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1148 **Methyl hexadecenoate (*trans*-9)** **100 mg**
Methyl palmitelaidate; C16:1 (*trans*-9) Methyl ester $C_{17}H_{32}O_2$ **CAS#:** 10030-74-7

Source: synthetic **Mol. Wt.:** 268 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1149 **Octadecenoic acid (*trans*-9)** **1 g**
Elaidic acid; C18:1 (*trans*-9) Fatty acid $C_{18}H_{34}O_2$ **CAS#:** 112-79-8

Source: synthetic **Mol. Wt.:** 282 **Purity:** 99% by TLC, GC **Appearance:** solid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1150 **Methyl octadecenoate (*trans*-9)** **1 g**
Methyl elaidate; C18:1 (*trans*-9) Methyl ester $C_{19}H_{36}O_2$ **CAS#:** 1937-62-8

Source: natural, plant **Mol. Wt.:** 296 **Purity:** 99% by TLC, GC **Appearance:** liquid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1262 **Octadecenoic acid (*trans*-11)** **100 mg**
trans Vaccenic acid; C18:1 (*trans*-11) Fatty acid $C_{18}H_{34}O_2$ **CAS#:** 693-72-1

Source: synthetic **Mol. Wt.:** 282 **Purity:** 99% by TLC, GC **Appearance:** solid
Solubility: chloroform, hexane, ethyl ether **Storage:** -20°C

1263	Methyl octadecenoate (<i>trans</i>-11) Methyl <i>trans</i> vaccenate; C18:1 (<i>trans</i> -11) Methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 6198-58-9 Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1151	Linoelaidic acid (all <i>trans</i>-9,12) C18:2 (all <i>trans</i> -9,12) Fatty acid C ₁₈ H ₃₂ O ₂ CAS#: 506-21-8 Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1152	Methyl octadecadienoate (all <i>trans</i>-9,12) Methyl linoelaidate; C18:2 (all <i>trans</i> -9,12) Methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 2566-97-4 Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	100 mg
1131	Cis-Trans Isomer Standard Mixture Source: margarine Appearance: liquid Solvent: 5ml methylene chloride Solubility: methylene chloride, chloroform Storage: -20°C Analysis of positional cis-trans fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of cis-trans fatty acid isomers as methyl esters in methylene chloride. This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 trans isomers (4 peaks), C18:1 cis & trans isomers (2 peaks), C18:1 cis isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0	5 mg/ml, 5 ml
1181	9(E),11(E)-Octadecadienoic acid 9- <i>trans</i> , 11- <i>trans</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-71-8 Source: synthetic Mol. Wt.: 280 Melting Point (°C): 55-57 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	25 mg

Conjugated Linoleic Acid Isomers (CLA)


Linoleic acid is an essential fatty acid (18:2 ω6) of which several naturally occurring conjugated derivatives have been identified. These derivatives, called “conjugated linoleic acid” or CLA can have the two double bonds mainly in the 9 and 11 or in the 10 and 12 positions, resulting in eight possible geometric isomers. CLA occurs in meat (41) and dairy products (42, 43). In both cases, the 9(Z),11(E)-isomer is predominant and is thought to be the biologically active form. CLA assimilated through the diet of animals is found in the intestinal musosa, liver and adipose tissue (44). See also review article by Parodi (43). CLA has several biological properties. It’s anti-carcinogenic activity has been demonstrated by its ability to inhibit chemically induced tumor formation in animal models of carcinogenesis (41,45-47). The addition of CLA to culture medium suppresses the *in vitro* growth of human melanoma, colorectal and breast cancer cells (48). CLA also exhibits anti-atherogenic activity. Addition of CLA to a controlled atherogenic diet significantly reduced the development of atherosclerosis in hamsters and rabbits (49,50). Animals fed a diet containing CLA also had lower levels of low-density-lipoprotein (LDL) cholesterol. CLA may be involved in regulating fat and protein metabolism (51,52). Several species of animals fed CLA-supplemented diets showed improved feed efficiency. Lean body mass increased while body fat was reduced. This seems to be due, mainly or exclusively, to the 10(E),12(Z)-isomer (catalog # 1249, see below). CLA competes with linoleate for Δ6 desaturase (53). Dietary CLA normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat (54) *via* activation of PPAR γ, a result which bears on the possible ameliorization or prevention of NIDDM. The 11(Z),13(E)-isomer (catalog # 1259) has been shown to be concentrated in the heart and in mitochondria.

See Literature References on page 99.

CLA Research is Being Redone With Our Highly Pure Isomers

Most studies to date have utilized a mixture of CLA isomers containing less than 30% of the presumed active 9(Z),11(E)-isomer (55,56). In addition to the 9,11- and 10,12-isomers, 8,10- and 11,13-isomers have recently been identified in the widely used mixture (56,57). Matreya offers a highly pure CLA which is 98+% the active 9,11-“cis, trans” isomer. The corresponding “trans,trans” and “cis,cis” isomers are also available. In addition, we now offer the pure 10(E),12(Z)-isomer, which has been widely sought for comparison studies.

See Literature References on page 99.



1245	9(Z),11(E)-Octadecadienoic acid	25 mg
1245-1	<i>9-cis, 11-trans</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	1 g
1245-10		10 g

Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, hexane, methanol, DMSO Storage: -20°C

1255	Methyl 9(Z), 11(E)-octadecadienoate	25 mg
	Methyl ester of CLA (<i>9-cis, 11-trans</i>) C ₁₉ H ₃₄ O ₂	

Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C

1181	9(E),11(E)-Octadecadienoic acid	25 mg
	<i>9-trans, 11-trans</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-71-8	

Source: synthetic Mol. Wt.: 280 Melting Point (°C): 55-57 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, hexane, methanol
Storage: -20°C

1257	Methyl 9(E),11(E)-octadecadienoate Methyl ester of CLA (9- <i>trans</i> , 11- <i>trans</i>) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1248 1248-1	9(Z),11(Z)-Octadecadienoic acid 9- <i>cis</i> , 11- <i>cis</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-40-7	25 mg 1 g
	Source: synthetic Mol. Wt.: 280 Melting Point (°C): 40-42 Purity: 96+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol, ethyl ether Storage: -20°C	
1256	Methyl 9(Z), 11(Z)-octadecadienoate Methyl ester of CLA (9- <i>cis</i> , 11- <i>cis</i>) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 96+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1249 1249-1 1249-10	10(E),12(Z)-Octadecadienoic acid 10- <i>trans</i> , 12- <i>cis</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 2420-44-2	25 mg 1 g 10 g
	Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1254	Methyl 10(E), 12(Z)-octadecadienoate Methyl ester of CLA (10- <i>trans</i> , 12- <i>cis</i>) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1259	11(Z), 13(E)-Octadecadienoic acid 11- <i>cis</i> , 13- <i>trans</i> CLA C ₁₈ H ₃₂ O ₂	25 mg
	Source: synthetic Mol. Wt.: 280 Purity: 77% <i>cis</i> , <i>trans</i> ; 2 % <i>cis</i> , <i>cis</i> ; 6% <i>trans</i> , <i>trans</i> by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1247-1 1247-10	9(Z),11(E)-Octadecadienoic acid 9- <i>cis</i> , 11- <i>trans</i> CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	1 g 10 g
	Source: synthetic Mol. Wt.: 280 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1%(E),(E) by TLC, GC Appearance: liquid Solubility: ethanol, ethyl ether, hexane Storage: -20°C	
1258	Methyl 9(Z),11(E)-octadecadienoate Methyl ester of CLA (9- <i>cis</i> , 11- <i>trans</i>) C ₁₉ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1% (E),(E) by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C	


Other CLA Products and Derivatives

1409	1-Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C		
1410	1-Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C		
1411	1-Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C		

Hydroxy Fatty Acids

2-Hydroxy Fatty Acids and Methyl Esters

These products are racemic and 98+% pure by GC and TLC. The 2-hydroxy fatty acids are components of glycosphingolipids and are involved in fatty acid degradation. They are stable and are supplied neat in vials.



Catalog number 1758


1758 1758-1	2-Hydroxydecanoic acid 2-Hydroxy C10:0 fatty acid C ₁₀ H ₂₀ O ₃ CAS#: 5393-81-7	50 mg 1 g
Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methanol Storage: -20°C		
1759 1759-1	Methyl 2-hydroxydecanoate 2-Hydroxy C10:0 methyl ester C ₁₁ H ₂₂ O ₃ CAS#: 71271-24-4	50 mg 1 g
Source: synthetic Mol. Wt.: 202 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methanol Storage: -20°C		

1701 1701-1	2-Hydroxydodecanoic acid 2-Hydroxy C12:0 fatty acid C ₁₂ H ₂₄ O ₃ CAS#: 2984-55-6	50 mg 1 g
Source: synthetic Mol. Wt.: 216 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methanol Storage: -20°C		
1702 1702-1	Methyl 2-hydroxydodecanoate 2-Hydroxy C12:0 methyl ester C ₁₃ H ₂₆ O ₃ CAS#: 51067-85-7	50 mg 1 g
Source: synthetic Mol. Wt.: 230 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C		
1703 1703-1	2-Hydroxytetradecanoic acid 2-Hydroxy C14:0 fatty acid C ₁₄ H ₂₈ O ₃ CAS#: 2507-55-3	50 mg 1 g
Source: synthetic Mol. Wt.: 244 Melting Point (°C): 81-82 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, methanol Storage: -20°C		
1704 1704-1	Methyl 2-hydroxytetradecanoate 2-Hydroxy C14:0 methyl ester C ₁₅ H ₃₀ O ₃ CAS#: 56009-40-6	50 mg 1 g
Source: synthetic Mol. Wt.: 258 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C		
1705 1705-1	2-Hydroxyhexadecanoic acid 2-Hydroxy C16:0 fatty acid C ₁₆ H ₃₂ O ₃ CAS#: 764-67-0	50 mg 1 g
Source: synthetic Mol. Wt.: 272 Melting Point (°C): 86-87 Purity: 98+% by TLC, GC Appearance: solid Solubility: methanol, chloroform/methanol, 2:1 Storage: -20°C		
1706 1706-1	Methyl 2-hydroxyhexadecanoate 2-Hydroxy C16:0 methyl ester C ₁₇ H ₃₄ O ₃ CAS#: 16742-51-1	50 mg 1 g
Source: synthetic Mol. Wt.: 286 Melting Point (°C): 59-60 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C		
1707 1707-1	2-Hydroxyoctadecanoic acid 2-Hydroxy C18:0 fatty acid C ₁₈ H ₃₆ O ₃ CAS#: 629-22-1	50 mg 1 g
Source: synthetic Mol. Wt.: 300 Melting Point (°C): 92-93 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		
1708 1708-1	Methyl 2-hydroxyoctadecanoate 2-Hydroxy C18:0 methyl ester C ₁₉ H ₃₈ O ₃ CAS#: 2420-35-1	50 mg 1 g
Source: synthetic Mol. Wt.: 315 Melting Point (°C): 64-66 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C		
1709 1709-0.5	2-Hydroxyeicosanoic acid 2-Hydroxy C20:0 fatty acid C ₂₀ H ₄₀ O ₃ CAS#: 16742-48-6	25 mg 0.5 g
Source: synthetic Mol. Wt.: 329 Melting Point (°C): 91-92 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		

1710 1710-0.5	Methyl 2-hydroxyeicosanoate 2-Hydroxy C20:0 methyl ester $C_{21}H_{42}O_3$ CAS#: 16742-49-7	25 mg 0.5 g
Source: synthetic Mol. Wt.: 343 Melting Point (°C): 62-64 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: -20°C		
1711 1711-0.5	2-Hydroxydocosanoic acid 2-Hydroxy C22:0 fatty acid $C_{22}H_{44}O_3$ CAS#: 13980-14-8	25 mg 0.5 g
Source: synthetic Mol. Wt.: 366 Melting Point (°C): 96-97 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		
1712 1712-0.5	Methyl 2-hydroxydocosanoate 2-Hydroxy C22:0 methyl ester $C_{23}H_{46}O_3$ CAS#: 13980-17-1	25 mg 0.5 g
Source: synthetic Mol. Wt.: 371 Melting Point (°C): 72-73 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: -20°C		
1713	2-Hydroxytricosanoic acid 2-Hydroxy C23:0 fatty acid $C_{23}H_{46}O_3$ CAS#: 2718-37-8	10 mg
Source: synthetic Mol. Wt.: 371 Melting Point (°C): 98-99 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		
1714	Methyl 2-hydroxytricosanoate 2-Hydroxy C23:0 methyl ester $C_{24}H_{48}O_3$ CAS#: 118745-41-8	10 mg
Source: synthetic Mol. Wt.: 385 Melting Point (°C): 68-70 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: -20°C		
1715	2-Hydroxytetracosanoic acid 2-Hydroxy C24:0 fatty acid; Cerebronic acid $C_{24}H_{48}O_3$ CAS#: 544-57-0	5 mg
Source: synthetic Mol. Wt.: 385 Melting Point (°C): 101-104 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		
1716	Methyl 2-hydroxytetracosanoate 2-Hydroxy C24:0 methyl ester $C_{25}H_{50}O_3$ CAS#: 2433-95-6	5 mg
Source: synthetic Mol. Wt.: 399 Melting Point (°C): 77-80 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethyl ether Storage: -20°C		
1722	2-Hydroxy Methyl Ester Mixture Quantitative mixture	10 mg/ml, 1 ml
Source: synthetic Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C		
Contains: 2-OH C14:0, 20.0%; 2-OH C16:0, 20.0%; 2-OH C18:0, 15.0%; 2-OH C20:0, 15.0%; 2-OH C22:0, 10.0%; 2-OH C23:0, 10.0%; 2-OH C24:0, 10.0%		

3-Hydroxy Fatty Acids and Methyl Esters

These products are racemic and 98+% pure by GC and TLC. 3-Hydroxy fatty acids occur in the lipid fraction of many microorganisms and are useful in the typing of microbial isolates. They are stable and are supplied neat in vials.



Catalog number 1747

1747 **3-Hydroxyhexanoic acid** **25 mg**
1747-0.5 3-Hydroxy C6:0 fatty acid C₆H₁₂O₃ CAS#: 10191-24-9 **0.5 g**

Source: synthetic Mol. Wt.: 132 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, methanol Storage: -20°C

1748 **Methyl 3-hydroxyhexanoate** **25 mg**
1748-0.5 3-Hydroxy C6:0 methyl ester C₇H₁₄O₃ CAS#: 21188-58-9 **0.5 g**

Source: synthetic Mol. Wt.: 146 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, methanol Storage: -20°C

1745 **3-Hydroxyoctanoic acid** **25 mg**
1745-0.5 3-Hydroxy C8:0 fatty acid C₈H₁₆O₃ CAS#: 88930-08-9 **0.5 g**

Source: synthetic Mol. Wt.: 160 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, methanol Storage: -20°C

1746 **Methyl 3-hydroxyoctanoate** **25 mg**
1746-0.5 3-Hydroxy C8:0 methyl ester C₉H₁₈O₃ CAS#: 85549-54-8 **0.5 g**

Source: synthetic Mol. Wt.: 174 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, ethyl ether Storage: -20°C

1725 **3-Hydroxynonanoic acid** **25 mg**
1725-0.5 3-Hydroxy C9:0 fatty acid C₉H₁₈O₃ CAS#: 88930-09-0 **0.5 g**

Source: synthetic Mol. Wt.: 174 Melting Point (°C): 60-62 Purity: 98+% by TLC, GC
Appearance: solid Solubility: chloroform, ethanol, methanol
Storage: -20°C

1726 **Methyl 3-hydroxynonanoate** **25 mg**
1726-0.5 3-Hydroxy C9:0 methyl ester C₁₀H₂₀O₃ CAS#: 83968-06-3 **0.5 g**

Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: liquid
Solubility: chloroform, ethanol, ethyl ether Storage: -20°C

1727 **3-Hydroxydecanoic acid** **25 mg**
1727-0.5 3-Hydroxy C10:0 fatty acid C₁₀H₂₀O₃ CAS#: 5561-87-5 **0.5 g**

Source: synthetic Mol. Wt.: 188 Melting Point (°C): 57-60 Purity: 98+% by TLC, GC
Appearance: solid Solubility: chloroform, ethanol, methanol Storage: -20°C

1728 **Methyl 3-hydroxydecanoate** **25 mg**
1728-0.5 3-Hydroxy C10:0 methyl ester $C_{11}H_{22}O_3$ **CAS#:** 62675-82-5 **0.5 g**

Source: synthetic **Mol. Wt.:** 202 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

1729 **3-Hydroxyundecanoic acid** **25 mg**
1729-0.5 3-Hydroxy C11:0 fatty acid $C_{11}H_{22}O_3$ **CAS#:** 40165-88-6 **0.5 g**

Source: synthetic **Mol. Wt.:** 202 **Melting Point (°C):** 74-76 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1730 **Methyl 3-hydroxyundecanoate** **25 mg**
1730-0.5 3-Hydroxy C11:0 methyl ester $C_{12}H_{24}O_3$ **CAS#:** 127593-21-9 **0.5 g**

Source: synthetic **Mol. Wt.:** 216 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

1731 **3-Hydroxydodecanoic acid** **25 mg**
1731-0.5 3-Hydroxy C12:0 fatty acid $C_{12}H_{24}O_3$ **CAS#:** 1883-13-2 **0.5 g**

Source: synthetic **Mol. Wt.:** 216 **Melting Point (°C):** 71-72 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** ethanol, methanol **Storage:** -20°C

1732 **Methyl 3-hydroxydodecanoate** **25 mg**
1732-0.5 3-Hydroxy C12:0 methyl ester $C_{13}H_{26}O_3$ **CAS#:** 85464-97-7 **0.5 g**

Source: synthetic **Mol. Wt.:** 230 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, ethyl ether **Storage:** -20°C

1733 **3-Hydroxytridecanoic acid** **25 mg**
1733-0.5 3-Hydroxy C13:0 fatty acid $C_{13}H_{26}O_3$ **CAS#:** 32602-69-0 **0.5 g**

Source: synthetic **Mol. Wt.:** 230 **Melting Point (°C):** 80-83 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1734 **Methyl 3-hydroxytridecanoate** **25 mg**
1734-0.5 3-Hydroxy C13:0 methyl ester $C_{14}H_{28}O_3$ **0.5 g**

Source: synthetic **Mol. Wt.:** 244 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethyl ether **Storage:** -20°C

1735 **3-Hydroxytetradecanoic acid** **25 mg**
1735-0.5 3-Hydroxy C14:0 fatty acid $C_{14}H_{28}O_3$ **CAS#:** 3422-31-9 **0.5 g**

Source: synthetic **Mol. Wt.:** 244 **Melting Point (°C):** 80-81 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1736 **Methyl 3-hydroxytetradecanoate** **25 mg**
1736-0.5 3-Hydroxy C14:0 methyl ester $C_{15}H_{30}O_3$ **CAS#:** 55682-83-2 **0.5 g**

Source: synthetic **Mol. Wt.:** 258 **Melting Point (°C):** 36-37 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform, ethyl ether, methanol
Storage: -20°C

1739 **3-Hydroxyhexadecanoic acid** **25 mg**
1739-0.5 3-Hydroxy C16:0 fatty acid C₁₆H₃₂O₃. CAS#: 928-17-6 **0.5 g**

Source: synthetic Mol. Wt.: 272 Melting Point (°C): 85-86 Purity: 98+% by TLC,
GC Appearance: solid Solubility: chloroform, ethanol, methanol Storage: -20°C

1740 **Methyl 3-hydroxyhexadecanoate** **25 mg**
1740-0.5 3-Hydroxy C16:0 methyl ester C₁₇H₃₄O₃ CAS#: 51883-36-4 **0.5 g**

Source: synthetic Mol. Wt.: 286 Melting Point (°C): 43-45 Purity: 98+% by TLC,
GC Appearance: solid Solubility: ethanol, methanol Storage: -20°C

1741 **3-Hydroxyheptadecanoic acid** **25 mg**
1741-0.5 3-Hydroxy C17:0 fatty acid C₁₇H₃₄O₃ CAS#: 40165-89-7 **0.5 g**

Source: synthetic Mol. Wt.: 286 Melting Point (°C): 93-95 Purity: 98+% by TLC,
GC Appearance: solid Solubility: ethanol, methanol Storage: -20°C

1742 **Methyl 3-hydroxyheptadecanoate** **25 mg**
1742-0.5 3-Hydroxy C17:0 methyl ester C₁₈H₃₆O₃ **0.5 g**

Source: synthetic Mol. Wt.: 300 Melting Point (°C): 53-55 Purity: 98+% by TLC,
GC Appearance: solid Solubility: ethanol, methanol Storage: -20°C


1743 **3-Hydroxyoctadecanoic acid** **25 mg**
1743-0.5 3-Hydroxy C18:0 fatty acid C₁₈H₃₆O₃ CAS#: 45261-96-9 **0.5 g**

Source: synthetic Mol. Wt.: 300 Melting Point (°C): 52-54 Purity: 98+% by TLC,
GC Appearance: solid Solubility: ethanol, methanol Storage: -20°C

1744 **Methyl 3-hydroxyoctadecanoate** **25 mg**
1744-0.5 3-Hydroxy C18:0 methyl ester C₁₉H₃₈O₃ CAS#: 14531-40-9 **0.5 g**

Source: synthetic Mol. Wt.: 314 Melting Point (°C): 52-54 Purity: 98+% by TLC,
GC Appearance: solid Solubility: ethanol, methanol Storage: -20°C

Omega Hydroxy Fatty Acids



Catalog number 1754

1754 **Royal Jelly acid** **50 mg**
10-Hydroxy-2-(E)-decenoic acid; *omega*-Hydroxy C10:1 (2-*trans*) fatty acid
C₁₀H₁₈O₃ CAS#: 14113-05-4

Source: synthetic Mol. Wt.: 186 Melting Point (°C): 63-65 Purity: 98+% by TLC,
GC Appearance: solid Solubility: chloroform, ethanol, methanol Storage: -20°C

1881	15-Hydroxypentadecanoic acid <i>omega</i> -Hydroxy C15:0 fatty acid C ₁₅ H ₃₀ O ₃ CAS#: 4617-33-8	25 mg
	Source: synthetic Mol. Wt.: 258 Melting Point (°C): 84-86 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, ethanol, methanol Storage: room temperature	
1882	Methyl 15-hydroxypentadecanoate <i>omega</i> -Hydroxy C15:0 methyl ester C ₁₆ H ₃₂ O ₃ CAS#: 76529-42-5	25 mg
	Source: synthetic Mol. Wt.: 272 Melting Point (°C): 50-52 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1760	17-Hydroxyheptadecanoic acid <i>omega</i> -Hydroxy C17:0 fatty acid C ₁₇ H ₃₄ O ₃ CAS#: 13099-34-8	25 mg
	Source: synthetic Mol. Wt.: 286 Melting Point (°C): 93-95 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1761	Methyl 17-hydroxyheptadecanoate <i>omega</i> -Hydroxy C17:0 methyl ester C ₁₈ H ₃₆ O ₃ CAS#: 94036-00-7	25 mg
	Source: synthetic Mol. Wt.: 300 Melting Point (°C): 59-63 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1877	20-Hydroxyeicosanoic acid <i>omega</i> -Hydroxy C20:0 fatty acid C ₂₀ H ₄₀ O ₃	25 mg
	Source: synthetic Mol. Wt.: 328 Melting Point (°C): 96-98 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol Storage: room temperature	
1878	Methyl 20-hydroxyeicosanoate <i>omega</i> -Hydroxy C20:0 methyl ester C ₂₁ H ₄₂ O ₃	25 mg
	Source: synthetic Mol. Wt.: 342 Melting Point (°C): 69-71 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1880	Methyl 21-hydroxyheneicosanoate <i>omega</i> -Hydroxy C21:0 methyl ester C ₂₂ H ₄₄ O ₃	25 mg
	Source: synthetic Mol. Wt.: 356 Melting Point (°C): 73-76 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1818	22-Hydroxydocosanoic acid <i>omega</i> -Hydroxy C22:0 fatty acid C ₂₂ H ₄₄ O ₃	25 mg
	Source: synthetic Mol. Wt.: 356 Melting Point (°C): 100-102 Purity: 98+% by TLC, GC Appearance: solid Solubility: chloroform, warm ethanol Storage: room temperature	

1819 **Methyl 22-hydroxydocosanoate** **25 mg**
omega-Hydroxy C22:0 methyl ester C₂₃H₄₆O₃

Source: synthetic **Mol. Wt.:** 370 **Melting Point (°C):** 73-75 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform, warm ethanol, ethyl ether
Storage: room temperature

1883 **Methyl 27-hydroxyheptacosanoate** **25 mg**
omega-Hydroxy C27:0 methyl ester C₂₈H₅₆O₃

Source: synthetic **Mol. Wt.:** 440 **Melting Point (°C):** 85-89 **Purity:** 97+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform **Storage:** room temperature

1884 **Methyl 30-hydroxytriacontanoate** **25 mg**
omega-Hydroxy C30:0 methyl ester C₃₁H₆₂O₃

Source: synthetic **Mol. Wt.:** 482 **Melting Point (°C):** 88-91 **Purity:** 97+% by TLC,
GC **Appearance:** solid **Solubility:** chloroform **Storage:** room temperature

Other Hydroxy Fatty Acids

1182 **Ricinelaidic acid** **100 mg**
12-Hydroxy C18:1 (9-*trans*) fatty acid C₁₈H₃₄O₃ **CAS#:** 82188-83-8

Source: synthetic **Mol. Wt.:** 298 **Melting Point (°C):** 50-53 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** ethanol, methanol **Storage:** -20°C

1183 **Methyl ricinelaidate** **100 mg**
12-Hydroxy C18:1 (9-*trans*) methyl ester C₁₉H₃₆O₃ **CAS#:** 7706-01-6


Source: synthetic **Mol. Wt.:** 312 **Purity:** 98+% by TLC, GC **Appearance:** solid
Solubility: ethanol, methanol **Storage:** -20°C

1766 **6-Hydroxyoctadecanoic acid** **10 mg**
6-Hydroxy C18:0 fatty acid C₁₈H₃₆O₃

Source: synthetic **Mol. Wt.:** 300 **Melting Point (°C):** 80-82 **Purity:** 98+% by TLC,
GC **Appearance:** solid **Solubility:** ethanol, methanol **Storage:** room temperature

Branched and Cyclic Fatty Acids

Iso-Fatty Acids and Esters




Catalog number 1656

1656 **Methyl 11-methyldodecanoate** **20 mg**
iso-Tridecanoic methyl ester; iso C13 Methyl ester C₁₄H₂₈O₂
CAS#: 5129-57-7

Source: synthetic **Mol. Wt.:** 228 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: hexane, ethyl ether, methylene chloride **Storage:** -20°C

1657	Methyl 12-methyltridecanoate iso-Tetradecanoic methyl ester; iso C14 Methyl ester $C_{15}H_{30}O_2$ CAS#: 5129-58-8 Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1605	13-Methyltetradecanoic acid iso-Pentadecanoic acid; iso C15 Fatty acid $C_{15}H_{30}O_2$ CAS#: 27836-87-9 Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: solid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1600	Methyl 13-methyltetradecanoate iso-Pentadecanoic methyl ester; iso C15 Methyl ester $C_{16}H_{32}O_2$ CAS#: 5129-59-9 Source: synthetic Mol. Wt.: 256 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1601	Methyl 14-methylpentadecanoate iso-Palmitic methyl ester; iso C16 Methyl ester $C_{17}H_{34}O_2$ CAS#: 5129-60-2 Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1606	15-Methylhexadecanoic acid iso-Heptadecanoic acid; iso C17 Fatty acid $C_{17}H_{34}O_2$ CAS#: 1603-03-8 Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: solid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1602	Methyl 15-methylhexadecanoate iso-Heptadecanoic methyl ester; iso C17 Methyl ester $C_{18}H_{36}O_2$ CAS#: 6929-04-0 Source: synthetic Mol. Wt.: 284 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg
1603	Methyl 17-methyloctadecanoate iso-Nonadecanoic methyl ester; iso C19 Methyl ester $C_{20}H_{40}O_2$ CAS#: 55124-97-5 Source: synthetic Mol. Wt.: 313 Purity: 98+% by GC Appearance: solid Solubility: chloroform, ethyl ether, ethanol Storage: $-20^{\circ}C$	20 mg

Anteiso-Fatty Acids and Esters



1615 **12-Methyltetradecanoic acid** **20 mg**

anteiso-Pentadecanoic acid; anteiso C15 Fatty acid $C_{15}H_{30}O_2$ **CAS#:** 5502-94-3

Source: synthetic **Mol. Wt.:** 242 **Purity:** 98+% by GC **Appearance:** solid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1612 **Methyl 12-methyltetradecanoate** **20 mg**

anteiso-Pentadecanoic methyl ester; anteiso C15 Methyl ester $C_{16}H_{32}O_2$
CAS#: 5129-66-8

Source: synthetic **Mol. Wt.:** 256 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1613 **Methyl 13-methylpentadecanoate** **20 mg**

anteiso-Palmitic methyl ester; anteiso C16 Methyl ester $C_{17}H_{34}O_2$
CAS#: 5487-50-3

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1616 **14-Methylhexadecanoic acid** **20 mg**

anteiso-Heptadecanoic acid; anteiso C17 Fatty acid $C_{17}H_{34}O_2$ **CAS#:** 5918-29-6

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by GC **Appearance:** solid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

1614 **Methyl 14-methylhexadecanoate** **20 mg**

anteiso-Heptadecanoic methyl ester; anteiso C17 Methyl ester $C_{18}H_{36}O_2$
CAS#: 2490-49-5

Source: synthetic **Mol. Wt.:** 284 **Purity:** 98+% by GC **Appearance:** liquid
Solubility: chloroform, ethyl ether, ethanol **Storage:** $-20^{\circ}C$

Methylated Fatty Acids

1207 **D,L-2,6-Dimethylheptanoic acid** **50 mg**

2,6-Dimethyl C7:0 fatty acid $C_9H_{18}O_2$

Source: synthetic **Mol. Wt.:** 158 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform **Storage:** room temperature

1791 **10-Methylhexadecanoic acid** **25 mg**
10-Methyl C16:0 fatty acid C₁₇H₃₄O₂

Source: synthetic **Mol. Wt.:** 270 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform **Storage:** room temperature

1792 **Methyl 10-methylhexadecanoate** **25 mg**
10-Methyl C16:0 methyl ester C₁₈H₃₆O₂

Source: synthetic **Mol. Wt.:** 284 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform **Storage:** room temperature

1195 **Phytanic acid** **25 mg**
3,7,11,15-Tetramethylhexadecanoic acid C₂₀H₄₀O₂ **CAS#:** 14721-66-5

Source: semi-synthetic **Mol. Wt.:** 312 **Purity:** 97+% by GC **Appearance:** solid
Solubility: chloroform, methanol **Storage:** -20°C

Cyclopropyl Fatty Acids and Esters

1822 **Methyleneoctadecanoic acid (all *cis*-9,10)** **25 mg**
Dihydrosterculic acid C₁₉H₃₆O₂ **CAS#:** 4675-61-0

Source: synthetic **Mol. Wt.:** 296 **Melting Point (°C):** 38-42 **Purity:** 98+% by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, hexane
Storage: -20°C

1823 **Methyl *cis*-9,10-methyleneoctadecanoate, C19:0 *delta* (all *cis*-9,10)** **25 mg**
Methyl dihydrosterculate C₂₀H₃₈O₂ **CAS#:** 3971-54-8

Source: synthetic **Mol. Wt.:** 310 **Purity:** 98+% by TLC, GC **Appearance:** liquid
Solubility: chloroform, ethanol, methanol, hexane **Storage:** -20°C

Unusual Fatty Acids and Derivatives

1751 **N-Oleoylethanolamine** **100 mg**
NOE C₂₀H₃₉NO₂ **CAS#:** 111-58-0

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, ethyl ether, DMSO
Storage: -20°C

Activity: acid ceramidase inhibitor

1786 **N-Hexadecanoylethanolamine** **100 mg**
C₁₈H₃₇NO₂ **CAS#** 544-31-0


Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform, ethanol, methanol, **Storage:** -20°C

Activity: inactive as acid ceramidase inhibitor

Other Lipids

Tocopherols

Catalog number 1072



1072


rac-alpha-Tocopherol

50 mg/ml, 1 ml

5,7,8-Trimethyltocol C₂₉H₅₀O₂ CAS#: 59-02-9

Source: synthetic **Mol. Wt.:** 431 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C

Catalog number 1071



1071


rac-beta-Tocopherol

50 mg/ml, 1 ml

5,8-Dimethyltocol C₂₈H₄₈O₂ CAS#: 148-03-8

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C

Catalog number 1073




1073

rac-gamma-Tocopherol

50 mg/ml, 1 ml


7,8-Dimethyltocol C₂₈H₄₈O₂ CAS#: 73980-80-0

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 97% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C




1074 **rac-5,7-Dimethyltocol** **50 mg/ml, 1 ml**
*C*₂₈H₄₈O₂ CAS#: 493-35-6

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, ethyl ether, chloroform, alcohols **Storage:** -20°C



1790 **(+)-delta-Tocopherol** **50 mg/ml, 1 ml**
8-Methyltocol *C*₂₇H₄₆O₂ CAS#: 119-13-1

Source: natural, plant **Mol. Wt.:** 403 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C




1797 **Tocol** **50 mg/ml, 1 ml**
rac-Tocol *C*₂₆H₄₄O₂

Source: synthetic **Mol. Wt.:** 389 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, methanol, ethanol **Storage:** -20°C

Cholestane Derivatives

Catalog number 1006



1006 **Cholesterol** **500 mg**
*C*₂₇*H*₄₆*O* CAS#: 57-88-5

Source: natural, ovine **Mol. Wt.:** 387 **Melting Point (°C):** 147-148 **Purity:** 98+%
by TLC, GC **Appearance:** solid **Solubility:** chloroform, ethanol **Storage:** -20°C

1115 **5-*alpha*-Cholestane** **100 mg**
*C*₂₇*H*₄₈ CAS#: 481-21-0

Source: synthetic **Mol. Wt.:** 373 **Purity:** 98+% by GC **Appearance:** solid
Solubility: chloroform, ethyl ether, hexane **Storage:** -20°C

1116 **Coprostanol** **25 mg**
5-beta-Cholestan-3-*beta*-ol *C*₂₇*H*₄₈*O* CAS#: 360-68-9

Source: semisynthetic **Mol. Wt.:** 389 **Melting Point (°C):** 101-103 **Purity:** 98+%
by GC **Appearance:** solid **Solubility:** chloroform, ethyl ether, warm methanol
Storage: -20°C

Plant Sterols and Steryl Glucosides

1119 **Plant Sterol Mixture** **25 mg/ml, 1 ml**
Sterol mixture, qualitative

Source: natural, plant **Appearance:** liquid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains: Brassicasterol, Campesterol, Stigmasterol, *beta*-Sitosterol, in order of elution

1123 **Plant Sterols Kit** **1 kit**

Source: synthetic or plant **Appearance:** liquid/solid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains in individual packages: steryl glucosides 25 mg, esterified steryl glucosides 10 mg, plant sterol mixture 25 mg, *beta*-sitosterol (55%) 100 mg, desmosterol (85%) 2 mg, lanosterol (55%) 100 mg, stigmasterol 25 mg, ergosterol 25 mg, coprostanol 5 mg, cholestanol 100 mg

1113 **beta-Sitostanol** **50 mg**
Stigmastanol C₂₉H₅₂O CAS#: 19466-47-8

Source: synthetic **Mol. Wt.:** 417 **Melting Point (°C):** 127-132 **Purity:** 98+% by TLC, 97+% by GC **Appearance:** solid **Solubility:** chloroform **Storage:** -20°C

1120 **Lanosterol** **500 mg**
C₃₀H₅₀O CAS#: 79-63-0

Source: synthetic or plant **Mol. Wt.:** 427 **Purity:** 55% by TLC, GC **Appearance:** solid **Solubility:** chloroform **Storage:** -20°C

1121 **Stigmasterol** **100 mg**
5,22-Cholestadien-24-beta-ethyl-3-beta-ol C₂₉H₄₈O CAS#: 83-48-7

Source: synthetic **Mol. Wt.:** 413 **Melting Point (°C):** 165-167 **Purity:** 95% by TLC, GC **Appearance:** solid **Solubility:** chloroform **Storage:** -20°C

1122 **Ergosterol** **100 mg**
C₂₈H₄₄O CAS#: 57-87-4

Source: synthetic or plant **Mol. Wt.:** 397 **Melting Point (°C):** 156-158 **Purity:** 95% by TLC, GC **Appearance:** solid **Solubility:** chloroform **Storage:** -20°C

1117 **Steryl glucosides** **25 mg**
C₃₅H₆₀O₆

Source: natural, plant **Mol. Wt.:** 577 (based on β-sitosteryl glucoside)
Melting Point (°C): 283-287 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform/methanol/DI water, 2:1:0.1 (warm) **Storage:** -20°C

1118 **Esterified Steryl Glucosides** **10 mg**
1:1:1, sterol:glucose:fatty acid C₅₁H₉₀O₇

Source: natural, plant **Mol. Wt.:** 815 (based on β-sitosteryl glucoside palmitate)
Purity: 98+% by TLC **Appearance:** solid **Solubility:** chloroform, ethyl ether, pyridine **Storage:** -20°C

Sterol, glucose and fatty acid in a molar ratio 1:1:1.

Standards and Reference Compounds

Food Industry Mixtures

Each methyl ester mixture is carefully prepared by weight.

4210 **KEL-FIM-FAME-5 Mixture** **15.5 mg/ml, 1 ml**
Methyl ester mixture

Source: synthetic or plant **Appearance:** liquid **Solvent:** heptane **Solubility:** heptane
Storage: -20°C

Contains the methyl esters of the following fatty acids (mg/ml in brackets): C8:0 [0.3], C10:0 [0.5], C12:0 [1.0], C13:0 [0.5], C14:0 [0.5], C14:1 [0.3], C15:0 [0.3], C16:0 [2.0], C16:1 [1.0], C17:0 [0.5], C18:0 [1.0], C18:1tr [0.4], C18:1c [3.0], C18:2 [2.0], C20:0 [0.3], C18:3 [1.0], C20:1 [0.3], C22:0 [0.3], C22:1 [0.3], listed in order of their elution.

2009 **FIM-FAME-6 Mixture** **33 mg/ml, 1 ml**
Methyl ester mixture

Source: synthetic or plant **Appearance:** liquid **Solvent:** heptane **Storage:** -20°C

Contains the methyl esters of these fatty acids. Each methyl ester is 3.03% of the mixture except C16:0 which is 6.06%. C4:0, C6:0, C8:0, C10:0, C11:0, C12:0, C13:0, C14:0, C14:1(*cis*-9), C15:0, C15:1(*cis*-10), C16:0, C16:1(*cis*-9), C17:0, C17:1(*cis*-10), C18:0, C18:1(*trans*-9), C18:1(*cis*-9), C18:2(all *cis*-9,12), C20:0, C18:3(all *cis*-6,9,12), C20:1(*cis*-11), C18:3(all *cis*-9,12,15), C20:2(all *cis*-11,14), C22:0, C20:3(all *cis*-8,11,14), C22:1(*cis*-13), C20:3(all *cis*-11,14,17), C20:4(all *cis*-5,8,11,14), C22:2(all *cis*-13,16), C24:1(*cis*-15), C22:6(all *cis*-4,7,10,13,16,19), listed in order of their elution.

2010 **FIM-FAME-7 Mixture** **30 mg/ml, 1 ml**
Methyl ester mixture

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Contains the methyl esters of these fatty acids (weight percent in [brackets]): C4:0 [4.0], C6:0 [4.0], C8:0 [4.0], C10:0 [4.0], C11:0 [2.0], C12:0 [4.0], C13:0 [2.0], C14:0 [4.0], C14:1(*cis*-9) [2.0], C15:0 [2.0], C15:1(*cis*-10) [2.0], C16:0 [6.0], C16:1(*cis*-9) [2.0], C17:0 [2.0], C17:1(*cis*-10) [2.0], C18:0 [4.0], C18:1(*trans*-9) [2.0], C18:1(*cis*-9) [4.0], C18:2(all *trans*-9,12) [2.0], C18:2(all *cis*-9,12) [2.0], C20:0 [4.0], C18:3(all *cis*-6,9,12) [2.0], C20:1(*cis*-11) [2.0], C18:3(all *cis*-9,12,15) [2.0], C21:0 [2.0], C20:2(all *cis*-11,14) [2.0], C22:0 [4.0], C20:3 (all *cis*-8,11,14) [2.0], C22:1(*cis*-13) [2.0], C20:3(all *cis*-11,14,17) [2.0], C20:4(all *cis*-5,8,11,14) [2.0], C23:0 [2.0], C22:2(all *cis*-13,16) [2.0], C24:0 [4.0], C20:5(all *cis*-5,8,11,14,17) [2.0], C24:1(*cis*-15) [2.0], C22:6(all *cis*-4,7,10,13,16,19) [2.0], listed in order of their elution.

Polyunsaturated Fatty Acid Methyl Esters Mixtures

These are complex qualitative standard mixtures of polyunsaturated fatty acid methyl esters. Because they are extracted from natural materials, relative peak sizes may vary from lot to lot.

1093 **PUFA-1** **100 mg**
Qualitative mixture

Source: natural, fish oil **Appearance:** liquid **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C

Contains: C14:0, C16:0, C16:1 ω 7, C18:1 ω 9, C18:1 ω 7, C18:2 ω 6, C20:1 ω 9, C18:4 ω 3, C22:1 ω 11, C22:1 ω 9, C20:5 ω 3, C22:5 ω 3, C22:6 ω 3

1081 **PUFA-2** **100 mg**
Qualitative mixture

Source: natural, porcine **Appearance:** liquid **Solubility:** alcohols, hexane, chloroform **Storage:** -20°C

Contains: C14:0, C16:0, C16:1 ω 7, C18:0, C18:1 ω 9, C18:1 ω 7, C18:2 ω 6, C18:3 ω 6, C18:3 ω 3, C20:1 ω 9, C20:2 ω 6, C20:3 ω 6, C20:4 ω 6, C20:5 ω 3, C22:4 ω 6, C22:5 ω 3, C22:6 ω 3

1177 **PUFA-3** **100 mg**
Qualitative mixture

Source: natural, menhaden oil **Appearance:** liquid **Solubility:** alcohols, hexane, chloroform **Storage:** -20°C

Contains: C14:0, C16:0, C16:1 ω 7, C16:2 ω 4, C16:3 ω 4, C16:4 ω 1, C18:0, C18:1 ω 9, C18:1 ω 7, C18:2 ω 6, C18:2 ω 4, C18:3 ω 4, C18:3 ω 3, C18:4 ω 3, C20:1 ω 9, C20:4 ω 6, C20:4 ω 3, C20:5 ω 3, C21:5 ω 3, C22:5 ω 3, C22:6 ω 3

Carbohydrate Mixtures

1124 **Alditol Acetate Mixture-1** **50 mg/ml, 1 ml**
Quantitative carbohydrate mixture

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

Contains: rhamnitol, fucitol, ribitol and arabinitol pentaacetates, 12.5 mg/ml each

1125 **Alditol Acetate Mixture-2** **50 mg/ml, 1 ml**
Quantitative carbohydrate mixture

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

Contains: mannitol, galactitol, glucitol and inositol hexaacetates, 12.5 mg/ml each

Other Fatty Acid Methyl Ester Mixtures

1722 **2-Hydroxy Methyl Ester Mixture** **10 mg/ml, 1 ml**

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform
Storage: -20°C

Quantitative mix contains: C14:0, 20.0%; C16:0, 20.0%; C18:0, 15.0%; C20:0, 15.0%;
C22:0, 10.0%; C23:0, 10.0%; C24:0, 10.0%

1131 **Cis-Trans Isomer Standard Mixture** **5 mg/ml, 5 ml**

Source: margarine **Appearance:** liquid **Solvent:** 5ml methylene chloride
Solubility: methylene chloride, chloroform **Storage:** -20°C

Analysis of positional *cis-trans* fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mixture to ensure proper operation of your column for this tricky separation. Mixture consists of *cis-trans* fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 *trans* isomers (4 peaks), C18:1 *cis* & *trans* isomers (2 peaks), C18:1 *cis* isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

2011 **Long Chain Fatty Acid Methyl Ester Mixture** **25 mg/ml, 1 ml**

C24:0, C26:0, C28:0, C30:0, C32:0 Fatty acid methyl ester mixture

Source: synthetic **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Quantitative mixture contains:
C24:0, 20.0%; C26:0, 20.0%; C28:0, 20.0%; C30:0, 20.0%; C32:0, 20.0%

AOCS Animal and Vegetable Oil Reference Mixtures (RM Mixtures)

By studying problems with the quantitative analysis of animal and vegetable oils and fats, the American Oil Chemists' Society has found certain mixtures to be useful as reference standards. The composition of each mixture (see Table I below) is similar to the fatty acid distribution of certain oils. All mixtures are in methyl ester form and ready for GC analysis

Table I. AOCS Oil Reference Mixtures

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated in the Table.

Mix No. Catalog No.	RM-1 1084	RM-2 1085	RM-3 1086	Rapeseed 1083	RM-4 1087	RM-5 1088	RM-6 1089
C8:0 Caprylate						7.0	
C10:0 Caprate						5.0	
C12:0 Laurate						48.0	
C14:0 Myristate			1.0	1.0		15.0	2.0
C16:0 Palmitate	6.0	7.0	4.0	4.0	11.0	7.0	30.0
C16:1 Palmitoleate (<i>cis</i> -9)							3.0
C18:0 Stearate	3.0	5.0	3.0	3.0	3.0	3.0	14.0
C18:1 Oleate (<i>cis</i> -9)	35.0	18.0	45.0	60.0	80.0	12.0	41.0
C18:2 Linoleate (all <i>cis</i> -9,12)	50.0	36.0	15.0	12.0	6.0	3.0	7.0
C18:3 Linolenate (all <i>cis</i> -9,12,15)	3.0	34.0	3.0	5.0			3.0
C20:0 Arachidate	3.0		3.0	3.0			
C20:1 Eicosenoate (<i>cis</i> -11)				1.0			
C22:0 Behenate			3.0	3.0			
C22:1 Erucate (<i>cis</i> -13)			20.0	5.0			
C24:0 Lignocerate			3.0	3.0			

1083 Rapeseed Oil Reference Mixture

25 mg/ml, 1 ml

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: ethyl ether, methylene chloride **Storage:** -20°C

Suitable standard for low erucic acid oil

1084 RM-1 Mixture

50 mg


Source: synthetic or plant **Appearance:** liquid **Solubility:** chloroform, ethyl ether
Storage: -20°C

Suitable standard for corn, cottonseed, soybean, safflower, sesame, poppy seed, walnut kapok, and rice oils

1085	RM-2 Mixture	50 mg
<p>Source: synthetic or plant Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C</p> <p>Suitable standard for linseed, perilla, hempseed, and rubberseed oils</p>		
1086	RM-3 Mixture	50 mg/ml, 1 ml
<p>Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: ethyl ether, methylene chloride Storage: -20°C</p> <p>Suitable standards for peanut, rapeseed, and mustard seed oils</p>		
1087	RM-4 Mixture	50 mg
<p>Source: synthetic or plant Appearance: liquid Solubility: chloroform, ethyl ether Storage: -20°C</p> <p>Suitable standard for olive, teaseed, and neatsfoot oils</p>		
1088	RM-5 Mixture	50 mg
<p>Source: synthetic or plant Appearance: liquid Solubility: chloroform Storage: -20°C</p> <p>Suitable standard for coconut, palm kernel, babassu and ouri-ouri oils</p>		
1089	RM-6 Mixture	50 mg
<p>Source: synthetic or plant Appearance: liquid Solubility: ethyl ether, methylene chloride Storage: -20°C</p> <p>Suitable standard for lard, beef tallow, mutton tallow, and palm oil</p>		

Custom Mixtures

Custom fatty acid methyl ester mixtures can be prepared to your specification. Minimum quantity requirements apply to these orders.



Cat# 4210 spiked with 0.4 mg/ml C18:2t ester (methyl linoleidate) and chromatographed on a Supelco SP 2330 fused silica column.

Peak number	FAME
1	C8:0
2	C10:0
3	C12:0
4	C13:0
5	C14:0
6	C14:1
7	C15:0
8	C16:0
9	C16:1
10	C17:0
11	C18:0
12	C18:1t-9
13	C18:1c-9
14	C18:2t,t-9,12
15	C18:2c,c-9,12
16	C20:0
17	C18:3
18	C20:1
19	C22:0
20	C22:1

Table II. Standards for GC Analysis

GLC Standard Mixtures

GLC-10 through GLC-100 standards are **equal weight measures** of fatty acid methyl esters. They are **quantitative standards**, useful for determining relative **retention times and response factors**.

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated below. All double bonds are *cis*.

Mixture Number Catalog Number	GLC-10 1095	GLC-30 1097	GLC-40 1098	GLC-50 1099	GLC-60 1100	GLC-70 1101	GLC-80 1102	GLC-90 1103	GLC-100 1104
C8:0 Caprylate		20.0				20.0			
C9:0 Nonanoate						20.0			
C10:0 Caprate		20.0				20.0			
C11:0 Undecanoate						20.0			
C12:0 Laurate		20.0				20.0			
C13:0 Tridecanoate							20.0	20.0	
C14:0 Myristate		20.0					20.0		
C15:0 Pentadecanoate							20.0	20.0	
C16:0 Palmitate	20.0	20.0	25.0				20.0		
C16:1 Palmitoleate (<i>cis</i> -9)				20.0					
C17:0 Heptadecanoate							20.0	20.0	
C18:0 Stearate	20.0		25.0						20.0
C18:1 Oleate (<i>cis</i> -9)	20.0			20.0					
C18:2 Linoleate (all <i>cis</i> -9,12)	20.0								
C18:3 Linolenate (all <i>cis</i> -9,12,15)	20.0								
C19:0 Nonadecanoate								20.0	20.0
C20:0 Arachidate			25.0		25.0				20.0
C20:1 Eicosenoate (<i>cis</i> -11)				20.0	25.0				
C20:2 Eicosadienoate (all <i>cis</i> -11,14)					25.0				
C20:3 Eicosatrienoate (all <i>cis</i> -11,14,17)					25.0				
C21:0 Heneicosanoate								20.0	20.0
C22:0 Behenate			25.0						20.0
C22:1 Erucate (<i>cis</i> -13)				20.0					
C24:1 Nervonate (<i>cis</i> -15)				20.0					

1095 GLC-10 Mixture 50 mg

Source: synthetic or plant Appearance: liquid Solubility: methylene chloride
Storage: -20°C

1097 GLC-30 Mixture 50 mg

Source: synthetic or plant Appearance: liquid Solubility: methylene chloride
Storage: -20°C

1098 GLC-40 Mixture 50 mg/ml, 1 ml

Source: synthetic or plant Appearance: liquid Solvent: methylene chloride
Solubility: methylene chloride Storage: -20°C

1099 **GLC-50 Mixture** **50 mg/ml, 1 ml**

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

1100 **GLC-60 Mixture** **50 mg/ml, 1 ml**

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

1101 **GLC-70 Mixture** **50 mg**

Source: synthetic or plant **Appearance:** liquid **Solubility:** methylene chloride
Storage: -20°C

1102 **GLC-80 Mixture** **50 mg/ml, 1 ml**

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

1103 **GLC-90 Mixture** **50 mg/ml, 1 ml**

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

1104 **GLC-100 Mixture** **50 mg/ml, 1 ml**

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Water Soluble Fatty Acid Mixtures

1106 **WSFA-2 Mixture** **5 ml**

Water soluble fatty acid qualitative mixture

Appearance: liquid **Solvent:** DI water **Solubility:** DI water **Storage:** Room Temp

Contains: acetic, propionic, isobutyric, n-butyric, isovaleric and n-valeric acids

1108 **WSFA-4 Mixture** **5 ml**

Water soluble fatty acid qualitative mixture

Appearance: liquid **Solvent:** DI water **Solubility:** DI water **Storage:** Room Temp

Contains: acetic, propionic, isobutyric, n-butyric, 2-methylbutyric, isovaleric and n-valeric acids

Microbiology Standard Mixtures

1105 **GLC-110 Mixture** **10 mg/ml, 1 ml**

Bacterial lipid standard, qualitative mixture

Source: various **Appearance:** liquid **Solvent:** chloroform **Solubility:** methylene chloride, chloroform **Storage:** –20°C

Contains:

Methyl 12-methyltridecanoate	(iso-C14:0)	Methyl 14-methylpentadecanoate	(iso-C16:0)
Methyl tetradecanoate (myristate)	(C14:0)	Methyl hexadecanoate (palmitate)	(C16:0)
Methyl 12-methyltetradecanoate	(anteiso-C15:0)	Methyl 14-methylhexadecanoate	(anteiso-C17:0)
Methyl pentadecanoate	(C15:0)		

1114 **Bacterial Acid Methyl Esters CP Mixture** **10 mg/ml, 1 ml**

Qualitative mixture

Source: various **Appearance:** liquid **Solvent:** methyl caproate **Solubility:** hexane, ethanol, methanol **Storage:** –20°C

A qualitative standard. Mixture consists of equal amounts of the compounds listed.

Methyl undecanoate	C11:0	Methyl cis-9-hexadecenoate (palmitoleate)	C16:1(cis-9)
Methyl 2-hydroxydecanoate	2-OH C10:0	Methyl hexadecanoate (palmitate)	C16:0
Methyl dodecanoate (laurate)	C12:0	Methyl 15-methylhexadecanoate	iso-C17:0
Methyl tridecanoate	C13:0	Methyl cis-9,10-methylenehexadecanoate	C17:0Δ (all cis-9,10)
Methyl 2-hydroxydodecanoate	2-OH C12:0	Methyl heptadecanoate (margarate)	C17:0
Methyl 3-hydroxydodecanoate	3-OH C12:0	Methyl 2-hydroxyhexadecanoate	2-OH C16:0
Methyl tetradecanoate (myristate)	C14:0	Methyl cis-9,12-octadecadienoate (linoleate)	C18:2 (all cis-9,12)
Methyl 13-methyltetradecanoate	iso-C15:0	Methyl cis-9-octadecenoate (oleate)	C18:1(cis-9)
Methyl 12-methyltetradecanoate	anteiso-C15:0	Methyl trans-9-octadecenoate (elaidate)	C18:1 (trans-9)
Methyl pentadecanoate	C15:0	Methyl octadecanoate (stearate)	C18:0
Methyl 2-hydroxytetradecanoate	2-OH C14:0	Methyl cis-9,10-methyleneoctadecanoate	C19:0Δ (all cis-9,10)
Methyl 3-hydroxytetradecanoate	3-OH C14:0	Methyl nonadecanoate	C19:0
Methyl 14-methylpentadecanoate	iso-C16:0	Methyl eicosanoate (arachidate)	C20:0

1075 **Volatile Acid Mixture** **100 ml**

Qualitative mixture

Appearance: liquid **Solvent:** DI water **Solubility:** DI water
Storage: 4-8°C

Contains: formic, acetic, propionic, isobutyric, n-butyric, isovaleric, n-valeric, isocaproic, n-caproic, and heptanoic acids

1077 **Non-Volatile Acid Mixture** **100 ml**

Qualitative mixture

Appearance: liquid **Solvent:** DI water **Solubility:** DI water
Storage: 4-8°C

Contains: pyruvic, lactic, oxalacetic, oxalic, methyl malonic, malonic, fumaric and succinic acids.

Biochemical Research Standard Mixtures

These mixtures are prepared by precise gravimetric technique. All mixtures contain equal amounts of listed components. A data sheet is supplied with each mixture.

1127 **Polar Lipid Mixture** **25 mg/ml, 1 ml**
TLC standards mixture

Source: natural, egg, ovine **Appearance:** liquid **Solvent:** chloroform/methanol, 2:1
Solubility: chloroform/methanol, 2:1 **Storage:** -20°C

Contains: cholesterol, phosphatidylethanolamine, lecithin, and *lyso*-lecithin

1128 **Sphingolipid Mixture** **25 mg/ml, 1 ml**
TLC standards mixture

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol, 2:1
Solubility: chloroform/methanol, 2:1 **Storage:** -20°C

Contains: cerebrosides, sulfatides, and sphingomyelin

1129 **Non-Polar Lipid Mixture A** **25 mg/ml, 1 ml**
TLC standards mixture

Source: natural, plant, ovine **Appearance:** liquid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains: cholesteryl palmitate, tripalmitin, palmitic acid, and cholesterol

1130 **Non-Polar Lipid Mixture B** **25 mg/ml, 1 ml**
TLC standards mixture

Source: natural, plant, ovine **Appearance:** liquid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains: cholesteryl oleate, methyl oleate, triolein, oleic acid, and cholesterol

Glycosphingolipid Reference Mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505 **Neutral Glycosphingolipid Mixture** **1 mg/ml, 1 ml**
Glycosylceramides, qualitative mixture

Source: natural, bovine and porcine **Appearance:** liquid
Solvent: chloroform/methanol, 2:1 **Solubility:** chloroform/methanol, 2:1
Storage: -20°C

Contains: cerebrosides, lactosylceramides, ceramide trihexosides, globosides

1508	Monosialoganglioside Mixture	0.5 mg/ml, 1 ml
<p>Source: natural, bovine, human Appearance: liquid Solvent: chloroform/methanol/DI water, 2:1:0.1 Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C</p> <p>Contains: GM₃, GM₂, GM₁</p>		
1509	Disialoganglioside Mixture	0.5 mg/ml, 1 ml
<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/DI water, 2:1:0.1 Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C</p> <p>Contains: GD₃, GD_{1a}, GD_{1b}</p>		
1510	Lactosylceramides and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml
<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/DI water, 2:1:0.1 Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C</p> <p>Contains: LC, GM₃, GD₃</p>		
1511	Gangliotetraosylceramide and Sialosyl Derivatives Mixture	0.5 mg/ml, 1 ml
<p>Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/DI water, 2:1:0.1 Solubility: chloroform/methanol/DI water, 2:1:0.1 Storage: -20°C</p> <p>Contains: asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b}</p>		

Biochemicals and Reagents

Stable Isotope Labeled Compounds

1914	N-Octadecanoyl-D₃₅-psychosine, (perdeuterated, C18:0 fatty acid) N-C18:0-D ₃₅ -Cerebrosides, perdeuterated; N-Stearoyl-D ₃₅ -psychosine, perdeuterated C ₄₂ H ₄₆ D ₃₅ NO ₈	5 mg
<p>Source: semisynthetic, bovine Mol. Wt.: 762 Purity: 98+% by TLC Appearance: solid Solubility: chloroform, hot ethanol, chloroform/methanol, 2:1 Storage: -20°C</p>		
1533	N-Hexadecanoyl-D₃-glucopsychosine, deuterated N-C16:0-D ₃ -Glucopsychosine, deuterated; N-C16:0-D ₃ -Glucocerebroside, deuterated; N-Palmitoyl-D ₃ -glucopsychosine, deuterated C ₄₀ H ₇₄ D ₃ NO ₈	1 mg
<p>Source: semisynthetic, bovine buttermilk Mol. Wt.: 703 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol, 2:1 Storage: -20°C</p>		
1534	N-Hexadecanoyl-D₃-lactosylceramide, deuterated N-C16:0-D ₃ -Lactosylceramide, deuterated; N-Palmitoyl-D ₃ -lactosylceramide, deuterated C ₄₆ H ₈₄ D ₃ NO ₁₃	1 mg
<p>Source: semisynthetic, bovine buttermilk Mol. Wt.: 865 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol/DI water, 5:1:0.1 Storage: -20°C</p>		

2200 **N-1-¹³C-Hexadecanoyl-sphingosylphosphorylcholine** **1 mg**
D-erythro-Sphingomyelin with 1-¹³C-palmitic acid; N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine ¹²C₃₈¹³CH₇₉N₂O₆P

Source: semisynthetic, bovine **Mol. Wt.:** 703 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1536 **N-Octadecanoyl-D₃-sulfatide, deuterated** **1 mg**
N-C18:0-D₃-Sulfatide, deuterated; N-Stearoyl-D₃-sulfatide, deuterated
C₄₂H₇₈D₃NO₁₁S

Source: semisynthetic, bovine **Mol. Wt.:** 833 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1
Storage: -20°C

1537 **N-Octadecanoyl-D₃-ceramide trihexoside, deuterated** **0.5 mg**
C18:0-D₃-CTH, deuterated; N-C18:0-D₃-Gb3, deuterated; N-Octadecanoyl-D₃-globotriaosylceramide, deuterated; N-Stearoyl-D₃-ceramide trihexoside, deuterated C₅₄H₉₈D₃NO₁₈

Source: semisynthetic, porcine **Mol. Wt.:** 1055 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO, chloroform/methanol, 2:1 **Storage:** -20°C

2050 **N-omega-CD₃-Octadecanoyl monosialoganglioside GM₁ (NH₄⁺ salt)** **0.5 mg**
N-CD₃-Stearoyl GM₁, C₇₃H₁₂₈N₃O₃₁D₃•NH₃

Source: semisynthetic, bovine **Mol. Wt.:** 1550 + NH₃ **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

2051 **N-omega-CD₃-Octadecanoyl monosialoganglioside GM₂ (NH₄⁺ salt)** **250 µg**
N-CD₃-Stearoyl GM₂ C₆₇H₁₁₈D₃N₃O₂₆•NH₃

Source: semisynthetic, human Tay-Sachs **Mol. Wt.:** 1388 + NH₃ **Purity:** 98+% by TLC, MS **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.1; forms micellar solution in water **Storage:** -20°C

2052 **N-omega-CD₃-Octadecanoyl monosialoganglioside GM₃ (NH₄⁺ salt)** **250 µg**
N-CD₃-Stearoyl GM₃ C₅₉H₁₀₅D₃N₂O₂₁•NH₃


Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 1185 + NH₃ **Purity:** 98+% by TLC, MS **Appearance:** solid **Solubility:** chloroform/methanol/DI water, 2:1:0.2; forms micellar solution in water **Storage:** -20°C

Fluorescent Compounds

1841 **N-Hexanoyl-NBD-D-erythro-sphingosine** **100 µg**
1841-001 N-C6:0-NBD-Ceramide; N-C6:0-NBD-D-erythro-Sphingosine C₃₀H₄₉N₅O₆
CAS#: 86701-10-2

Source: synthetic **Mol. Wt.:** 576 **Melting Point (°C):** 85-88 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Catalog number 1618



Excitation: 460 nm
Emission: 535 nm

1618 **N-Dodecanoyl-NBD-D-erythro-sphingosine** **100 µg**
1618-001 N-C12:0-NBD-Ceramide; N-C12:0-NBD-D-erythro-Sphingosine **1 mg**
C₃₆H₆₁N₅O₆
Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1857 **N-Hexanoyl-NBD-L-threo-sphingosine** **100 µg**
1857-001 N-C6:0-NBD-Ceramide; N-C6:0-NBD-L-threo-Sphingosine **1 mg**
C₃₀H₄₉N₅O₆
Source: synthetic **Mol. Wt.:** 575 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

1620 **N-Dodecanoyl-NBD-L-threo-sphingosine** **100 µg**
1620-001 N-C12:0-NBD-Ceramide; N-C12:0-NBD-L-threo-Sphingosine **1 mg**
C₃₆H₆₁N₅O₆
Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1624 **N-Hexanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
1624-001 N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-L-threo-Dihydrosphingosine **1 mg**
C₃₀H₅₁N₅O₆
Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1623 **N-Dodecanoyl-NBD-L-threo-dihydrosphingosine** **100 µg**
1623-001 N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-L-threo-Dihydrosphingosine **1 mg**
C₃₆H₆₃N₅O₆
Source: synthetic **Mol. Wt.:** 662 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1626 **N-Hexanoyl-NBD-D-erythro-dihydrosphingosine** **100 µg**
1626-001 N-C6:0-NBD-Dihydroceramide; N-C6:0-NBD-D-erythro-Dihydrosphingosine **1 mg**
C₃₀H₅₁N₅O₆
Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1625 **N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine** **100 µg**
1625-001 N-C12:0-NBD-Dihydroceramide; N-C12:0-NBD-D-erythro-
Dihydrosphingosine **1 mg**
C₃₆H₆₃N₅O₆
Source: synthetic **Mol. Wt.:** 662 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1628 **N-Hexanoyl-NBD-phytosphingosine** **100 µg**
1628-001 N-C6:0-NBD-Phytoceramide; N-C6:0-NBD-Phytosphingosine C₃₀H₅₁N₅O₇ **1 mg**

Source: semisynthetic, bacteria **Mol. Wt.:** 594 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1627 **N-Dodecanoyl-NBD-phytosphingosine** **100 µg**
1627-001 N-C12:0-NBD-Phytoceramide; N-C12:0-NBD-Phytosphingosine C₃₆H₆₃N₅O₇ **1 mg**

Source: semisynthetic, bacteria **Mol. Wt.:** 678 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

1912 **N-Hexanoyl-NBD-sphingosylphosphorylcholine** **100 µg**
1912-001 N-C6:0-NBD-Sphingomyelin; N-C6:0-NBD-Sphingosylphosphorylcholine
C₃₅H₆₁N₆O₉P **CAS#:** 94885-04-8 **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 740 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Mixture of D-erythro and L-threo isomers

1619 **N-Dodecanoyl-NBD-sphingosylphosphorylcholine** **100 µg**
1619-001 N-C12:0-NBD-Sphingomyelin; N-C12:0-NBD-Sphingosylphosphorylcholine
C₄₁H₇₃N₆O₉P **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 825 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 2:1 **Storage:** -20°C

Mixture of D-erythro and L-threo isomers

1621 **N-Hexanoyl-NBD-galactosylceramide** **100 µg**
1621-001 N-C6:0-NBD-*beta*-D-Galactosylsphingosine; N-C6:0-NBD-Cerebrosides
C₃₆H₅₉N₅O₁₁ **1 mg**

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1 **Storage:** -20°C

1633 **N-Dodecanoyl-NBD-galactosylceramide** **100 µg**
1633-001 N-C12:0-NBD-*beta*-D-Galactosylsphingosine; N-C12:0-NBD-Cerebroside
C₄₂H₇₁N₅O₁₁ **1 mg**

Source: semisynthetic, bovine spinal cord **Mol. Wt.:** 822 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform, DMSO, chloroform/methanol, 2:1
Storage: -20°C

1622 **N-Hexanoyl-NBD-glucosylceramide** **100 µg**
1622-001 N-C6:0-NBD-*beta*-D-Glucosylsphingosine; N-C6:0-NBD-Glucosylceramide,
fluorescent C₃₆H₅₉N₅O₁₁ **1 mg**

Source: semisynthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** methanol, chloroform/methanol, 5:1 **Storage:** -20°C

1629 **N-Hexanoyl-NBD-lactosylceramide** **50 µg**
1629-001 N-Hexanoyl-NBD-*beta*-D-lactosylsphingosine; N-C6:0-NBD-*beta*-D-
Lactosylsphingosine; N-C6:0-NBD-Lactosylceramide C₄₂H₆₉N₅O₁₆ **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 900 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1630 **N-Dodecanoyl-NBD-lactosylceramide** **50 µg**
1630-001 N-Dodecanoyl-NBD-*beta*-D-lactosylsphingosine; N-C12:0-NBD-*beta*-D-
Lactosylsphingosine; N-C12:0-NBD-Lactosylceramide C₄₈H₈₁N₅O₁₆ **1 mg**

Source: semisynthetic, bovine buttermilk **Mol. Wt.:** 984 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1631 **N-Dodecanoyl-NBD-ceramide trihexoside** **100 µg**
1631-001 N-C12:0-NBD-CTH; N-C12:0-NBD-Globotriaosylceramide C₅₄H₉₁N₅O₂₁ **1 mg**

Source: semisynthetic, porcine **Mol. Wt.:** 1145 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** DMSO; hot methanol, chloroform/methanol, 2:1
Storage: -20°C

1632 **N-Dodecanoyl-NBD-sulfatide** **100 µg**
1632-001 N-C12:0-NBD-Sulfatide; N-Dodecanoyl-NBD-*lyso*-sulfatide; N-Dodecanoyl-
NBD-sphingosyl-*beta*-D-galactoside-3-sulfate C₄₂H₇₁N₅O₁₄S **1 mg**

Source: semisynthetic, bovine **Mol. Wt.:** 901 **Purity:** 98+% by TLC
Appearance: solid **Solubility:** chloroform/methanol, 2:1 **Storage:** -20°C

1634 **omega-N-NBD-D-erythro-C14-Sphingosine** **1 mg**
omega-N-(7-nitrobenzo-2-oxa-1,3-diazol-4-yl)-(2S)-amino-tetradec-(4E)-ene-
(1,3R)-diol C₂₀H₃₁N₅O₅

Source: synthetic **Mol. Wt.:** 422 **Purity:** 98+% by TLC **Appearance:** solid
Solubility: methanol, ethanol, chloroform/methanol, 9:1 **Storage:** -20°C

Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC
 (actual composition may vary according to dietary history and growth condition of the source)

	Lecithin (egg)	Phosphatidyl-ethanolamine (egg)	lyso-Lecithin (egg)	Phosphatidylserine (bovine)	Phosphatidylinositol (plant)	Sulfatides (bovine)	Cerebrosides (bovine)	Sphingomyelin (bovine)	Phosphatidic acid (semi-synthetic)	Ceramides (bovine)
Catalog Number	#1044	#1045	#1046	#1047	#1048	#1049	#1050	#1051	#1053	#1056
Fatty Acids										
C14:0										
C16:0	31	17	72	1	42	trace	trace	4	39	trace
C16:1										
C18:0	16	29	24	42		5	4	40	12	4
C18:1	31	17	3	27	6	trace			34	
C18:2	16	11			47				15	
C18:3					5					
C20:0				1		1	1	3		1
C20:1				4						
C20:4		12		4						
C21:0										
C22:0				1		7	4	13		4
C22:1				1		trace				
C22:6				7						
C23:0							2	2		2
C24:0						18	10	9		10
C24:1						29	15	22		15
C25:0						2	3			9
C25:1						2	1			1
C26:0						1	2			2
C26:1						3	1			1
C27:0						1	2			2
C27:1							2			2
C14:0 2-OH										
C16:0 2-OH										
C18:0 2-OH						5	15			15
C20:0 2-OH						trace	1			1
C22:0 2-OH						3	6			6
C23:0 2-OH							5			5
C24:0 2-OH						10	17			17
C24:1 2-OH						6	6			
C25:0 2-OH						2	3			3
C25:1 2-OH										
C26:0 2-OH										
C26:1 2-OH										
C16 cis 9,10 methylene										
C18 cis 9,10 methylene										
Others	6	14	1	12	0	5	0	7	0	0
Total	100	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Glucocerebrosides (Gaucher's spleen)	Monogalactosyl- diglycerides (plant)	Digalactosyl- diglyceride (plant)	Monosialo- ganglioside GM ₁	Disialoganglioside GD _{1a}	Trisialogang- lioside GT _{1b}	Gangliotetraosyl- ceramide	Purified mixed gangliosides (bovine)	Cerebrosides Kerasin (bovine)	Ceramide trihexoside (porcine)
Catalog Number	#1057	#1058	#1059	#1061	#1062	#1063	#1064	#1065	#1066	#1067
Fatty Acids										
C14:0					1		trace	trace		
C16:0	26	23	9	2	1	1	1	1	trace	3
C16:1										
C18:0	9	77	91	90	89	87	86	86	5	2
C18:1						1	3	3		2
C18:2										
C18:3										
C20:0	5			3	2	4	4	4	1	2
C20:1										
C20:4										
C21:0										
C22:0	26			1	1	1	2	2	9	17
C22:1									trace	
C22:6										
C23:0	5					1	1	1	5	1
C24:0	22					1	1	1	25	29
C24:1	6			1		1	2	2	43	5
C25:0									3	
C25:1									3	
C26:0									2	
C26:1									4	
C27:0										
C27:1										
C14:0 2-OH										
C16:0 2-OH										
C18:0 2-OH										
C20:0 2-OH										
C22:0 2-OH										3
C23:0 2-OH										1
C24:0 2-OH										19
C24:1 2-OH										10
C25:0 2-OH										
C25:1 2-OH										
C26:0 2-OH										
C26:1 2-OH										
C16 cis 9,10 methylene										
C18 cis 9,10 methylene										
Others	1	0	0	3	6	3	0	0	0	6
Total	100	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Globosides (porcine)	Lecithin (bovine)	Esterified steryl glucoside	Cerebrosides Phrenosin (bovine)	Phosphatidyl- ethanolamine (plant)	Lecithin (plant)	Ceramides (non- hydroxy)	Ceramides (hydroxy)	Sphingomyelin (porcine RBC)	Sphingomyelin (buttermilk)
Catalog Number	#1068	#1070	#1118	#1138	#1301	#1302	#1322	#1323	#1328	#1329
Fatty Acids										
C14:0		trace								1
C16:0	2	35	34		22	14			25	14
C16:1		1								
C18:0	2	14	8		3	4	11		7	3
C18:1		33	8		7	11				
C18:2			36		60	65				
C18:3			4		8	6				
C20:0	2		1				2		3	1
C20:1										
C20:4										
C21:0										
C22:0	20		4				10		9	26
C22:1										
C22:6										
C23:0	2		2				6		1	30
C24:0	33		2				24		22	21
C24:1	5						31		22	3
C25:0							3			
C25:1							3			
C26:0	2						2			
C26:1							3			
C27:0										
C27:1										
C14:0 2-OH										
C16:0 2-OH										
C18:0 2-OH				36				24		
C20:0 2-OH				1				1		
C22:0 2-OH	4			8				8		
C23:0 2-OH				6				6		
C24:0 2-OH	19			25				35		
C24:1 2-OH	9			9				17		
C25:0 2-OH				4				4		
C25:1 2-OH				2						
C26:0 2-OH				2						
C26:1 2-OH				2				2		
C16 cis 9,10 methylene										
C18 cis 9,10 methylene										
Others	0	17	1	5	0	0	5	3	11	1
Total	100	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Spingomyelin, (egg, chicken)	Phosphatidylinositol	Lactosyl ceramide (porcine)	Disialoganglioside GD _{1b}	Monosialoganglioside GM2	Monosialoganglioside GM3 (buttermilk)	Disialoganglioside GD3 (buttermilk)	Lactosyl ceramide (buttermilk)	Ceramide trihexosides (top spot)
Catalog Number	#1332	#1336	#1500	#1501	#1502	#1503	#1504	#1507	#1513
Fatty Acids									
C14:0	trace			trace					
C16:0	72	32	14	1	2	6	8	12	1
C16:1									
C18:0	8	7	6	86	82	1	1	1	1
C18:1	3	7	4	3					
C18:2		47							
C18:3		6							
C20:0	2		1	4	7	1	1	1	2
C20:1									
C20:4									
C21:0						1	2		
C22:0	5		9	2	4	23	24	25	22
C22:1									
C22:6									
C23:0	1		1	1	trace	36	35	36	2
C24:0	2		15	1	1	22	21	21	58
C24:1	4		5	2	2	3	3		7
C25:0								1	1
C25:1									
C26:0									5
C26:1									
C27:0									
C27:1									
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH			trace						
C20:0 2-OH									
C22:0 2-OH			8						
C23:0 2-OH									
C24:0 2-OH			24						
C24:1 2-OH			13						
C25:0 2-OH									
C25:1 2-OH									
C26:0 2-OH									
C26:1 2-OH									
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	3	1	0	0	2	7	5	3	1
Total	100	100	100	100	100	100	100	100	100


Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
(actual composition may vary according to dietary history and growth condition of the source)

	Ceramide trihexosides (bottom spot)	Tetrasialoganglioside GQ1 _b	Glucocerebrosides (buttermilk)	Glucocerebrosides (plant)	Mixed Gangliosides, purified (porcine)	Fucosylated monosialoganglioside GM ₁	Disialogangliosides GD ₂	Monosialoganglioside GM ₄
Catalog Number	#1514	#1516	#1521	#1522	#1525	#1526	#1527	#1535
Fatty Acids								
C14:0								
C16:0	3	5	7		1	8	1	4
C16:1		1						
C18:0		80	2		87	2	89	2
C18:1		2						
C18:2		3						
C18:3								
C20:0		4	1		4	13	7	trace
C20:1								trace
C20:4								
C21:0			1					
C22:0	2	2	27		1	43	1	3
C22:1								4
C22:6								
C23:0			36		1	3	1	4
C24:0	3		23		1	26		6
C24:1					2	5	1	4
C25:0			1					
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH				trace				
C16:0 2-OH				79				
C18:0 2-OH	1			trace				1
C20:0 2-OH	1							3
C22:0 2-OH	11			8				25
C23:0 2-OH	1			1				17
C24:0 2-OH	52			9				18
C24:1 2-OH	25							7
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	1	3	2	3	3			2
Total	100	100	100	100	100	100	100	100

Sphingolipid Metabolism Pathways

Matreya
Lipids for Research

168 Tressler Street
Pleasant Gap, PA 16823 USA
Tel: 814 359 5060 Fax: 814 359 5062
e-mail: customer-service@matreya.com Web: www.matreya.com



Sphingolipid Structures and Pathways Wall Chart

In a clear and straightforward manner, this wall chart indicates the structures and relationships between most commonly discussed sphingolipids. Full size copies (approximately 35 x 26 inches) are available on request to customer service.

Literature References

1. B.A. Fenderson, E.M. Eddy, S.Hakomori, *BioEssays* 12, 173, 1990
2. A. Gorio et al., *Exp. Brain Res.*, Suppl. 13, 283, 1984
3. F. Di Gregorio et al., *Neuropediatrics*, Suppl. 15, 93, 1984
4. J.S. Schneider et al., *Science*, 256, 843, 1992
5. R.W. Leeden, R.K. Yu, *Methods Enzymol.*, 83, 139, 1982
6. Borman, S., *Chemical and Engineering News* 82:32, 31-35, 2004
7. M. Faucher et al., *J. Biol. Chem.* 263, 5319, 1988
8. B. M. Buehrer, R.M. Bell, *Adv. in Lipid Res.*, 26, 59, 1993
9. C. W. Sachs et al., *J. Biol. Chem.*, 270, 26639, 1995
10. Radin, N. S., *Biochem. J.* 371:243, 2003
11. Radin, N. S., *Biorg. Med. Chem.* 11:2123, 2003
12. Radin, N. S., *Cancer Investigation* 20:1, 2002
13. Sietsma H., Veldman R.J., Vander Kolk D., Ausema B., Nijhof W., Kamps W., Vellenga E., Kok J.W. *Clin. Cancer Res.* 6, 942-948, 2000
14. Shayman, J. A., *Kidney Int.* 58, 11-26, 2000
15. Kolesnick, R. N. et al. *J. Cell Physiol.* 184, 285-300, 2000
16. Smith, W. L. and Merrill, A. H., Jr. *J. Biol. Chem.* 277, 25841-25842, 2002
17. Pyne, S. And Pyne, N. J. *Biochem. J.* 349, 385-402, 2000
18. Merrill, A. H. Jr. *J. Biol. Chem.* 277, 25843-25846, 2002
19. Hannun, Y. A. and Obeid, L., *J. Biol. Chem.* 277, 25847-25850, 2002
20. van Meer, G. And Lisman, Q. *J. Biol. Chem.* 277, 25855-25858, 2002
21. Kolter, T., Proia, R. L. and Sandhoff, K. *J. Biol. Chem.* 277, 25859-25862, 2002
22. Radin, N. S., *Cancer Invest.* 20, 779-786, 2002
23. Tettamanti, G., Bassi R., Viani, P., Riboni, L., *Biochimie* 85(3-4), 423-437, 2003
24. Fredman, P., Hedberg, K., Brezicka, T., *Biodrugs* 17(3) 155-167, 2003
25. Hoeckstra, D., et al. *J. Lipid Res.* 44, 869-877, 2003
26. R.R, Vunnam, N.S. Radin, *Chem. Phys. Lipids*, 265, 1980
27. J. Inokuchi and N.S. Radin, *J. Lipid Res.*, 28, 565, 1987
28. N.S. Radin et al. *J. Biochem.*, 111, 191, 1992
29. J. Inokuchi et al., *Cancer letters*, 38, 23, 1987
30. Y. Lavie et al., *J. Biol. Chem.*, 271, 19530, 1996
31. Y. Lavie et al., *J. Biol. Chem.*, 271, in press, 1996
32. Y. Hannun et al., *Science*, 235, 670, 1987
33. S. Spiegel et al., *Proc. Intern. Conf. Biol. Function Glycosphingolipids*, Santa Barbara, CA, 1990
34. Jacqueline M. Kravka, Li Li, Zdzislaw M. Szulc, Jacek Bielawski, Besium Ogretmen, Yusuf A. Hannun, Lina M. Obeid, and Alicja Bielawska. *J. Biol. Chem.*, 10, 1074/jbc. M700647200, February 5, 2007
35. M. Sugita et al. *Biochim. Biophys. Acta*, 398, 125, 1975
36. A. Bielawska et al. *J. Biol. Chem.*, 271, 12646, 1996
37. A. Bielawska et al. *ibid.* 267, 18493, 1992
38. J. M. L. Hauser et al., *J. Biol. Chem.* 269, 6803, 1994
39. Y. Hannun, *ibid.*, 269, 3125, 1994
40. R. Kolesnick, D.W. Golde, *Cell*, 77, 325, 1984
41. M.W. Pariza et al., *Cancer Res.* 43, 2444s, 1983
42. Y. L. Ha et al., *J. Agr. Food Chem.*, 37, 75, 1989
43. P.W. Parodi, *Austral. J. Dairy Tech.*, 49, 93, 1994
44. S. Banni et al. *Abstr. 87th AOCS Mtg.*, 28, 1996
45. C. Ip et al., *Cancer Res.* 51, 6118, 1991
46. M. A. Belury, *Nutr. Rev.* 53 (4 Pt. 1), 83, 1995
47. C. Liew et al. *Carcinogenesis*, 16, 3037, 1995
48. T. D. Shultz et al. *Cancer Let.*, 63, 125, 1992
49. R.J. Nicolosi, *Abstr. 87th AOCS Mtg.*, 1996
50. K.N. Lee et al., *Atheroscl.*, 108(1), 19, 1994
51. B.F. Haumann, *Inform.*, 7(2), 152, 1996
52. M.W. Pariza et al., *Abstr.*, 87th AOCS Mtg., 1996
53. M.A. Belury et al., *Lipids*, 32, 199, 1997
54. Houseknecht, K.L. et al., *Biochem. Biophys. Res. Comm.*, 224, 678, 1998
55. W.W. Christie et al. *JAOCS*, 74, 1231, 1997
56. Sehat, N. et al., *Lipids*, 33, 217, 1998
57. Sébédio, J. L. et al., *Biochim. Biophys. Acta*, 1345,5, 1997

Note: We are pleased and honored to have the above list of scientists in the field of Lipid Research. In case your name and contribution is not listed above, we apologize. If you would like your publication listed in our next catalog, please send your name and publication to the attention of Marketing Department.

Cross Reference for Product Numbers and Catalog Pages

1006	76	1077	86	1192	57
1008	49	1081	79	1193	57
1009	49	1083	81	1194	58
1010	50	1084	81	1195	73
1011	50	1085	81, 82	1196	48
1012	50	1086	81, 82	1197	48
1013	50	1087	81, 82	1198	49
1014	50	1088	81, 82	1199	49
1015	50	1089	81, 82	1200	48
1016	53	1093	79	1203	54
1017	53	1095	84	1204	54
1018	50	1097	84	1205	56
1019	51	1098	84	1206	56
1020	51	1099	84, 85	1207	72
1021	51	1100	84, 85	1208	54
1022	54	1101	84, 85	1241	51
1023	54	1102	84, 85	1242	51
1024	55	1103	84, 85	1243	53
1025	55	1104	84, 85	1244	58
1026	56	1105	86	1245	61
1027	56	1106	85	1245-1	61
1028	51	1108	85	1245-10	61
1029	51	1113	77	1247-1	62
1030	51	1114	86	1247-10	62
1031	51	1115	76	1248	62
1032	57	1116	76	1248-1	62
1033	57	1117	77	1249	62
1034	57	1118	77, 97	1249-1	62
1035	52	1119	76	1249-10	62
1036	52	1120	77	1251	52
1037	52	1121	77	1252	52
1038	52	1122	77	1254	62
1040	53	1123	76	1255	61
1041	59	1124	79	1256	62
1042	57	1125	79	1257	62
1044	40, 94	1127	87	1258	62
1045	41, 94	1128	87	1259	62
1046	40, 94	1129	87	1261	49
1047	40, 94	1130	87	1262	55, 59
1048	41, 94	1131	60, 80	1263	55, 59
1049	22, 96	1136	58	1264	58
1050	19, 96	1138	20, 96	1265	58
1051	15, 95	1147	53, 59	1266	55
1051-1	15	1148	54, 59	1267	55
1052	41	1149	54, 59	1269	57
1053	41, 95	1150	54, 59	1271	52
1056	11, 97	1151	55, 60	1273	53
1057	21, 98	1152	55, 60	1275	53
1057-25	21	1153	56	1276	56
1058	48, 95	1154	56	1277	56
1059	48, 95	1155	59	1301	41, 94
1061	29, 95	1156	59	1302	40, 94
1061-50	29	1157	53	1303	47
1062	31, 95	1161	50	1303-2	47
1063	31, 96	1162	50	1305	20
1064	29, 96	1163	49	1306	22
1065	32, 96	1164	49	1310	22
1066	19, 96	1165	49	1318	17
1067	25, 97	1166	49	1319	17
1067-10	25	1167	58	1320	5
1068	26, 97	1175	58	1321	17
1069	41	1177	79	1321-05	17
1070	40, 94	1179	57	1322	11, 97
1071	74	1181	60, 61	1322-05	11
1072	74	1182	70	1323	12, 97
1073	74	1183	70	1323-05	12
1074	75	1186	52	1324	4
1075	86	1187	52	1325	20

1326	4	1537	26, 27, 89	1713	65
1327	16	1538	25	1714	65
1328	15, 95	1600	71	1715	65
1329	15, 98	1601	71	1716	65
1329-1	15	1602	71	1717	39
1330	5	1603	71	1718	39
1330-1	5	1605	71	1719	37
1332	15, 97	1606	71	1720	37
1332-1	15	1612	72	1722	65, 80
1333	8	1613	72	1725	66
1333-100	8	1614	72	1725-0.5	66
1334	20	1615	72	1726	66
1334-50	20	1616	72	1726-0.5	66
1335	20	1618	13, 90	1727	66
1336	41	1618-001	13, 90	1727-0.5	66
1400	43	1619	18, 91	1728	67
1409	43, 63	1619-001	18, 91	1728-0.5	67
1410	43, 63	1620	13, 90	1729	67
1411	44, 63	1620-001	13, 90	1729-0.5	67
1425	43	1621	21, 27, 91	1730	67
1426	43	1621-001	21, 27, 91	1730-0.5	67
1427	43	1622	21, 28, 91	1731	67
1428	42	1622-001	21, 28, 91	1731-0.5	67
1429	42	1623	14, 90	1732	67
1430	42	1623-001	14, 90	1732-0.5	67
1431	44	1624	13, 90	1733	67
1432	44	1624-001	13, 90	1733-0.5	67
1433	44	1625	14, 90	1734	67
1434	45	1625-001	14, 90	1734-0.5	67
1435	45	1626	14, 90	1735	67
1436	45	1626-001	14, 90	1735-0.5	67
1437	43	1627	14, 91	1736	67
1438	44	1627-001	14, 91	1736-0.5	67
1439	45	1628	14, 91	1739	68
1442	42	1628-001	14, 91	1739-0.5	68
1443	44	1629	25, 28, 91	1740	68
1444	45	1629-001	25, 28, 91	1740-0.5	68
1445	43	1630	25, 28, 92	1741	68
1500	24, 98	1630-001	25, 28, 92	1741-0.5	68
1501	31, 96	1631	26, 28, 92	1742	68
1502	30, 98	1631-001	26, 28, 92	1742-0.5	68
1503	30, 98	1632	24, 28, 92	1743	68
1504	31, 98	1632-01	24, 28, 92	1743-0.5	68
1505	32, 87	1633	21, 28, 91	1744	68
1507	24, 98	1633-001	21, 28, 91	1744-0.5	68
1507-50	24	1634	2, 92	1745	66
1508	32, 88	1656	70	1745-0.5	66
1509	32, 88	1657	71	1746	66
1510	33, 88	1701	64	1746-0.5	66
1511	33, 88	1701-1	64	1747	66
1512	29	1702	64	1747-0.5	66
1513	25	1702-1	64	1748	66
1514	25	1703	64	1748-0.5	66
1516	32, 97	1703-1	64	1749	37
1517	24	1704	64	1750	39
1518	30	1704-1	64	1751	35, 73
1520	26	1705	64	1752	39
1521	21, 98	1705-1	64	1753	38
1521-50	21	1706	64	1754	68
1522	21, 98	1706-1	64	1755	38
1522-100	21	1707	64	1756	38
1523	26	1707-1	64	1758	63
1524	26	1708	64	1758-1	63
1525	32	1708-1	64	1759	64
1526	30	1709	65	1759-1	64
1527	31	1709-0.5	65	1760	69
1531	22	1710	65	1761	69
1532	24	1710-0.5	65	1766	70
1533	22, 27, 88	1711	65	1773	46
1534	24, 27, 88	1711-0.5	65	1773-1	46
1535	31, 97	1712	65	1773-5	46
1536	23, 27, 89	1712-0.5	65	1775	47

1775-1	47	1882	69	2046	18
1775-5	47	1883	70	2047	11
1778	47	1884	70	2050	30,89
1778-1	47	1886	36	2051	30,89
1778-5	47	1886-005	36	2052	30,89
1779	46	1887	36	2076	23
1779-1	46	1887-005	36	2077	10
1783	47	1888	23	2200	16, 89
1783-1	47	1889	37	4210	78
1783-5	47	1890	16		
1784	46	1891	5		
1784-1	46	1892	5		
1784-5	46	1893	5		
1786	35, 73	1894	12		
1790	75	1895	12		
1791	73	1896	6		
1792	73	1897	12		
1797	75	1900	7		
1800	38	1900-100	7		
1802	2	1901	7		
1803	17	1901-100	7		
1805	6	1903	8		
1806	2	1903-100	8		
1807	3, 35	1904	23		
1807-025	3, 35	1907	15		
1809	8	1909	15		
1810	8	1910	10		
1818	69	1911	16		
1819	70	1912	18, 91		
1822	73	1912-001	18, 91		
1823	73	1913	17		
1826	2	1914	20, 27, 88		
1827	2	1915	8		
1828	7	1915-100	8		
1829	7	1916	9		
1830	8	1916-25	9		
1831	3	1917	16		
1831-1	3	1918	16		
1832	9	1919	46		
1832-100	9	1919-1	46		
1833	2	1919-5	46		
1834	10	1931	23		
1835	3	1930	9		
1838	3	1930-25	9		
1839	4	1932	23		
1840	3	1933	23		
1841	13, 89	1950	33		
1841-001	13, 89	1951	33		
1842	10	1954	33		
1843	9	1957	34		
1845	4	1960	34		
1846	4	1961	34		
1847	7	1963	34		
1848	7	1964	34		
1850	9	1977	33		
1851	4	2009	78		
1852	18	2010	78		
1854	10	2011	80		
1855	9	2034	12		
1857	13, 90	2035	12		
1857-001	13, 90	2036	13		
1858	39	2037	8		
1859	36	2037-100	8		
1860	36	2038	8		
1865	38	2038-100	8		
1868	38	2039	9		
1875	23	2039-100	9		
1876	4	2041	11		
1877	69	2042	9		
1878	69	2043	11		
1880	69	2044	10		
1881	69	2045	11		

Product Name Index

10-Methylhexadecanoic acid	73
12-Methyltetradecanoic acid	72
13-Methyltetradecanoic acid	71
14-Methylhexadecanoic acid	72
15-Hydroxypentadecanoic acid	69
15-Hydroxypentadecanoic acid methyl ester	69
15-Methylhexadecanoic acid	71
17-Hydroxyheptadecanoic acid	69
17-Hydroxyheptadecanoic acid methyl ester	69
20-Hydroxyeicosanoic acid	69
20-Hydroxyeicosanoic acid methyl ester	69
21-Hydroxyheneicosanoic acid methyl ester	69
22-Hydroxydocosanoic acid	69
22-Hydroxydocosanoic acid methyl ester	69
27-Hydroxyheptacosanoic acid methyl ester	70
30-Hydroxytriacontanoic acid methyl ester	70
6-Hydroxyoctadecanoic acid	70

A

Acetyl-4-(1R,2S,3R,4-tetrahydroxybutyl)-imidazole	39
Alditol Acetate Mixture-1	79
Alditol Acetate Mixture-2	79
anteiso-Heptadecanoic acid methyl ester	72
anteiso-Palmitic acid methyl ester	72
anteiso-Pentadecanoic acid methyl ester	72
anteiso-Pentadecanoic acid methyl ester	72
Anti-ganglioside asialo GM ₁	33
Anti-ganglioside asialo GM ₂	33
Anti-ganglioside GD _{1b}	34
Anti-ganglioside GD ₂	34
Anti-ganglioside GD ₃	33
Anti-ganglioside GM ₁	33
Anti-ganglioside GM ₂ (NANA)	34
Anti-ganglioside GM ₄	34
Anti-globoside GL-4	34
Antibody asialo GM ₁	33
Antibody asialo GM ₂	33
Antibody disialoganglioside GD _{1b}	34
Antibody disialoganglioside GD ₂	34
Antibody ganglioside GD ₃	33
Antibody globoside	34
Antibody monosialoganglioside GM ₁	33
Antibody monosialoganglioside GM ₂ (NANA)	34
Antibody monosialoganglioside GM ₄	34
Arachidic acid	51
Arachidic acid methyl ester	51
Arachidonic acid	57
Arachidonic acid methyl ester	57
Asialo GM ₁	29
Asialo GM ₂	29

B

Bacterial Acid Methyl Esters CP Mixture	86
---	----

Behenic acid	52
Behenic acid methyl ester	52

C

Capric acid methyl ester	49
Caproic acid methyl ester	48
Caprylic acid	49
Caprylic acid methyl ester	49
Castanospermine	38
Ceramide-1-phosphate, N-C16:0-D-erythro	18
Ceramide, N-C10:0-D-erythro	8
Ceramide, N-C12:0-NBD-D-erythro	13,90
Ceramide, N-C12:0-NBD-L-threo	13,90
Ceramide, N-C15:0-D-erythro	8
Ceramide, N-C16:0-D-erythro	8
Ceramide, N-C17:0-D-erythro	8
Ceramide, N-C18:0-D-erythro	9
Ceramide, N-C18:0-D-threo	9
Ceramide, N-C18:0-L-erythro	9
Ceramide, N-C18:0-L-threo	9
Ceramide, N-C19:0-D-erythro	9
Ceramide, N-C2:0-D-erythro	7
Ceramide, N-C2:0-D-erythro (C14 sphingoid base)	10
Ceramide, N-C2:0-L-erythro	7
Ceramide, N-C2:0-L-threo	7
Ceramide, N-C24:0-D-erythro	9
Ceramide, N-C24:1 (cis-15)-D-erythro	9
Ceramide, N-C6:0-D-erythro	7
Ceramide, N-C6:0-L-erythro	7
Ceramide, N-C6:0-L-threo	7
Ceramide, N-C6:0-NBD-D-erythro	13,89
Ceramide, N-C6:0-NBD-L-threo	13,90
Ceramide, N-C8:0-D-erythro	8
Ceramide, N-C8:0-D-threo	8
Ceramide, N-C8:0-L-threo	8
Ceramide, N-(R,S)-alpha-hydroxy-C12:0-D-erythro	9
Ceramide, N-(R,S)-alpha-hydroxy-C18:0-D-erythro	10
Ceramide, N-hexadecanoyl-D-erythro (C16 sphingoid base)	10
Ceramide, N-hexanoyl-D-threo	8
Ceramide trihexoside, lyso	26
Ceramide trihexoside, N-C17:0	26
Ceramide trihexoside, N-C12:0-NBD	26,28,92
Ceramide trihexoside, N-C18:0-D3	26,27,89
Ceramide trihexoside, N-C23:0	26
Ceramide trihexoside, N-dodecanoyl-NBD	26,28,92
Ceramide trihexoside, N-heptadecanoyl	26
Ceramide trihexoside, N-octadecanoyl-D ₃	26,27,89
Ceramide trihexoside, N-stearoyl-D ₃	26,27
Ceramide trihexoside, N-tricosanoyl	26
Ceramide trihexosides	25,95

Ceramide trihexosides (bottom spot)	25,98
Ceramide trihexosides (top spot)	25,97
Ceramides (hydroxy and non-hydroxy acyl groups)	11,94
Ceramides (mostly hydroxy acyl groups)	12,96
Ceramides (mostly non-hydroxy acyl groups)	11,96
Cerebroside; Kerasin (top spot)	19,95
Cerebroside, N-C12:0-NBD	21,28,91
Cerebroside, N-C15:0	20
Cerebroside, N-C18:0-D ₃₅	20,27,88
Cerebroside, N-C2:0	20
Cerebroside, N-C6:0-NBD	21,27,91
Cerebroside, N-C8:0	20
Cerebroside, Phrenosin (bottom spot)	20,96
Cerebroside sulfate	22,94
Cerebrosides	19,94
Cerotic acid	52
Cerotic acid methyl ester	52
Cholestane, 5-alpha	76
Cholesterol	76
Cis-Trans Isomer Standard Mixture	60,80
Conduritol B Epoxide	37
Coprostanol	76
Cyclopropenylceramide, C8:0-N	36
Cyclopropenylceramide, C16:0-N	36

D

D-MAPP	36
Decanoic acid methyl ester	49
Diffuoropalmitic acid	39
Digalactosyldiglyceride	48,95
Diheptadecanoyl-sn-glycero-3-phosphorylcholine	43
Dihydroceramide, N-C12:0-NBD-D-erythro	14,90
Dihydroceramide, N-C12:0-NBD-L-threo	14,90
Dihydroceramide, N-C18:0-D-erythro	11
Dihydroceramide, N-C2:0-D-erythro	10
Dihydroceramide, N-C6:0-D-erythro	10
Dihydroceramide, N-C6:0-NBD-D-erythro	14,90
Dihydroceramide, N-C6:0-NBD-L-threo	13,90
Dihydroceramide, N-C8:0-D-erythro	10
Dihydroceramide, N-(R,S)-alpha-hydroxy-C16:0-D-erythro	11
Dihydroceramide, N-(R,S)-alpha-hydroxy-C18:0-D-erythro	11
Dihydroceramide, N-(R,S)-alpha-hydroxyoctadecanoyl-D-erythro	11
Dihydrosphingosine, D-erythro-C ₂₀	3
Dihydrosphingosine, D-erythro-C ₂₀	4
Dihydrosphingosine, D-threo	4
Dihydrosphingosine, D,L-C ₁₆	4
Dihydrosphingosine, D,L-erythro	4
Dihydrosphingosine, D,L-erythro-C ₂₀	4
Dihydrosphingosine, L-erythro	4

Hydroxydocosanoic acid (α)	65
Hydroxydocosanoic acid (α) methyl ester	65
Hydroxydodecanoic acid (α)	64
Hydroxydodecanoic acid (α) methyl ester	64
Hydroxydodecanoic acid (β)	67
Hydroxydodecanoic acid (β) methyl ester	67
Hydroxyeicosanoic acid (α)	64
Hydroxyeicosanoic acid (α) methyl ester	65
Hydroxyheptadecanoic acid (β)	68
Hydroxyheptadecanoic acid (β) methyl ester	68
Hydroxyhexadecanoic acid (α)	64
Hydroxyhexadecanoic acid (α) methyl ester	64
Hydroxyhexadecanoic acid (β)	68
Hydroxyhexadecanoic acid (β) methyl ester	68
Hydroxyhexanoic acid (β)	66
Hydroxyhexanoic acid (β) methyl ester	66
Hydroxynonanoic acid (β)	66
Hydroxynonanoic acid (β) methyl ester	66,67
Hydroxyoctadecanoic acid (α)	64
Hydroxyoctadecanoic acid (α) methyl ester	64
Hydroxyoctadecanoic acid (β)	68
Hydroxyoctadecanoic acid (β) methyl ester	68
Hydroxyoctanoic acid (β)	66
Hydroxyoctanoic acid (β) methyl ester	66
Hydroxytetracosanoic acid (α)	65
Hydroxytetracosanoic acid (α) methyl ester	65
Hydroxytetradecanoic acid (α)	64
Hydroxytetradecanoic acid (α) methyl ester	64
Hydroxytetradecanoic acid (β)	67
Hydroxytetradecanoic acid (β) methyl ester	67
Hydroxytricosanoic acid (α)	65
Hydroxytricosanoic acid (α) methyl ester	65
Hydroxytridecanoic acid (β)	67
Hydroxytridecanoic acid (β) methyl ester	67
Hydroxyundecanoic acid (β)	67
Hydroxyundecanoic acid (β) methyl ester	67
I	
iso-Heptadecanoic acid	71
iso-Heptadecanoic acid methyl ester	71
iso-Nonadecanoic acid	71
iso-Nonadecanoic acid methyl ester	71
iso-Palmitic acid methyl ester	71
iso-Pentadecanoic acid	71
iso-Pentadecanoic acid methyl ester	71
iso-Tetradecanoic acid	71
iso-Tetradecanoic acid methyl ester	71
iso-Tridecanoic acid methyl ester	70

K	
KEL-FIM-FAME-5 Mixture	78
L	
L-MAPP	36
Lacceroic acid methyl ester	53
Lactosylceramide and Sialosyl Derivatives Mixture	33,88
Lactosylceramide, <i>lyso</i>	24
Lactosylceramide, N-C12:0-NBD	25,28,92
Lactosylceramide, N-C16:0	24
Lactosylceramide, N-C16:0-D ₃	24,27,88
Lactosylceramide, N-C17:0	25
Lactosylceramide, N-C6:0-NBD	25,28,91
Lactosylceramide, N-dodecanoyl-NBD	25,28,92
Lactosylceramide, N-heptadecanoyl	25
Lactosylceramide, N-hexadecanoyl	24
Lactosylceramide, N-hexadecanoyl-D ₃	24,27,88
Lactosylceramide, N-hexanol-NBD	25,28,91
Lactosylceramide, N-palmitoyl	24
Lactosylceramide, N-palmitoyl-D ₃	24,27,88
Lactosylceramides, bovine buttermilk	24,97
Lactosylceramides, porcine	24,97
Lactocerebrosides, bovine buttermilk	24,97
Lactocerebrosides, porcine	24,97
Lanosterol	77
Lauric acid	49
Lauric acid methyl ester	49
Lecithin, bovine	40,96
Lecithin, egg	40,94
Lecithin, <i>lyso</i> , egg	40,94
Lecithin, plant	40,96
Lignoceric acid	52
Lignoceric acid methyl ester	52
Linoelaidic acid	55,60
Linoelaidic acid methyl ester	55,60
Linoleic acid	55
Linoleic acid methyl ester	55
Linolenic acid	56
Linolenic acid- <i>gamma</i>	56
Linolenic acid- <i>gamma</i> methyl ester	56
Linolenic acid methyl ester	56
Long Chain Fatty Acid Methyl Ester Mixture	80
Loxastatin	39
<i>lyso</i> -Ceramide trihexoside	26
<i>lyso</i> -Cerebroside	20
<i>lyso</i> -Dihydrosphingomyelin	17
<i>lyso</i> -Glucocerebroside, bovine buttermilk	22
<i>lyso</i> -Glucocerebroside, plant	22
<i>lyso</i> -Lactosylceramide	24
<i>lyso</i> -Lecithin, egg	40,94
<i>lyso</i> -Monosialoganglioside GM ₁	30
<i>lyso</i> -Phosphatidylcholine, egg	40,94
<i>lyso</i> -Sphingomyelin	17
<i>lyso</i> -Sulfatide	23

M	
Margaric acid	50
Margaric acid methyl ester	51
Mead acid methyl ester	57
Melissoic acid methyl ester	53
Methyl 10(E),12(Z)-octadecadienoate	62
Methyl 10-methylhexadecanoate	73
Methyl 11-methyldodecanoate	70
Methyl 12-methyltetradecanoate	72
Methyl 12-methyltridecanoate	71
Methyl 13-methylpentadecanoate	72
Methyl 13-methyltetradecanoate	71
Methyl 14-methylhexadecanoate	72
Methyl 14-methylpentadecanoate	71
Methyl 15-hydroxypentadecanoate	69
Methyl 15-methylhexadecanoate	71
Methyl 17-hydroxyheptadecanoate	69
Methyl 17-methyloctadecanoate	71
Methyl 20-hydroxyeicosanoate	69
Methyl 21-hydroxyheneicosanoate	69
Methyl 22-hydroxydocosanoate	70
Methyl 27-hydroxyheptacosanoate	70
Methyl 2-fluoropalmitate	39
Methyl 2-hydroxydecanoate	63
Methyl 2-hydroxydocosanoate	65
Methyl 2-hydroxydodecanoate	64
Methyl 2-hydroxyeicosanoate	65
Methyl 2-hydroxyhexadecanoate	64
Methyl 2-hydroxyoctadecanoate	64
Methyl 2-hydroxytetraacosanoate	65
Methyl 2-hydroxytetradecanoate	64
Methyl 2-hydroxytricosanoate	65
Methyl 30-hydroxytriacontanoate	70
Methyl 3-hydroxydecanoate	67
Methyl 3-hydroxydodecanoate	67
Methyl 3-hydroxyheptadecanoate	68
Methyl 3-hydroxyhexadecanoate	68
Methyl 3-hydroxyhexanoate	66
Methyl 3-hydroxynonanoate	66
Methyl 3-hydroxyoctadecanoate	68
Methyl 3-hydroxyoctanoate	66
Methyl 3-hydroxytetradecanoate	67
Methyl 3-hydroxytridecanoate	67
Methyl 3-hydroxyundecanoate	67
Methyl 9(E),11(E)-octadecadienoate	62
Methyl 9(Z) 11(E)-octadecadienoate	61,62
Methyl 9(Z) 11(Z)-octadecadienoate	62
Methyl arachidate	51
Methyl arachidonate	57
Methyl behenate	52
Methyl caprate	49
Methyl caproate	48
Methyl caprylate	49
Methyl cerotate	52
Methyl <i>cis</i> -9,10-methylenecetadecanoate, C19:0 <i>delta</i>	73
Methyl decanoate	49
Methyl dihydrosterculate	73
Methyl docosaehaenoate	59
Methyl docosanoate	52
Methyl docosapentaenoate	58
Methyl docosenoate	58
Methyl dodecanoate	49
Methyl dotriacontanoate	53
Methyl eicosadienoate	57
Methyl eicosanoate	51
Methyl eicosapentaenoate	58

Methyl eicosatetraenoate	57	Methyltocol-8	75	Octadecatrienoic acid	
Methyl eicosatrienoate		Monogalatosyldiglyceride	48,95	(all <i>cis</i> -6,9,12)	56
(all <i>cis</i> -5,8,11)	57	Monosialoganglioside GM ₁	29,95	Octadecatrienoic acid	
Methyl eicosatrienoate		Monosialoganglioside GM1, fucosyl		(all <i>cis</i> -9,12,15)	56
(all <i>cis</i> -8,11,14)	57		30	Octadecatrienoic acid methyl ester	
Methyl eicosenoate	57	Monosialoganglioside GM ₁ , <i>lyso</i>	30	(all <i>cis</i> -6,9,12)	56
Methyl elaidate	54	Monosialoganglioside GM ₁ , N-		Octadecatrienoic acid methyl ester	
Methyl erucate	58	<i>omega</i> -CD ₃ -octadecanoyl	30,89	(all <i>cis</i> -9,12,15)	56
Methyl <i>gamma</i> -linolenate	56	Monosialoganglioside GM ₁ , N-		Octadecenoic acid (<i>cis</i> -11)	55
Methyl heneicosanoate	51	<i>omega</i> -CD ₃ -stearoyl	30,89	Octadecenoic acid (<i>cis</i> -9)	54
Methyl heptadecanoate	51	Monosialoganglioside GM ₂	30,97	Octadecenoic acid methyl ester	
Methyl heptadecenoate	54	Monosialoganglioside GM ₂ , N-		(<i>cis</i> -9)	54
Methyl heptanoate	48	<i>omega</i> -CD ₃ -octadecanoyl	30,89	Octadecenoic acid methyl ester	
Methyl hexacosanoate	52	Monosialoganglioside GM ₂ , N-		(<i>cis</i> -11)	55
Methyl hexadecanoate	50	<i>omega</i> -CD ₃ -stearoyl	30,89	Octadecenoic acid methyl ester	
Methyl hexadecenoate (<i>cis</i> -9)	53	Monosialoganglioside GM ₃	30,97	(<i>trans</i> -11)	55,60
Methyl hexadecenoate (<i>trans</i> -9)		Monosialoganglioside GM ₃ , N-		Octadecenoic acid methyl ester	
	54,59	<i>omega</i> -CD ₃ -octadecanoyl	30,89	(<i>trans</i> -9)	54,59
Methyl hexanoate	48	Monosialoganglioside GM ₃ , N-		Octadecenoic acid (<i>trans</i> -11)	55,59
Methyl homogamma linolenate	57	<i>omega</i> -CD ₃ -stearoyl	30,89	Octadecenoic acid (<i>trans</i> -9)	54,59
Methyl lacceroate	53	Monosialoganglioside GM ₄	31,98	Octanoic acid	49
Methyl laurate	49	Monosialoganglioside Mixture	32,88	Octanoic acid methyl ester	49
Methyl lignocerate	52	Montanoic acid methyl ester	52	Oleic acid	54
Methyl linoelaidate	55,60	Morotic acid	56	Oleic acid methyl ester	54
Methyl linoleate	55	Morotic acid methyl ester	56	Oleoylethanolamine, N	35,73
Methyl linolenate	56	Myristic acid	50		
Methyl margarate	51	Myristic acid methyl ester	50	P	
Methyl melissate	53	Myristoleic acid	53	Palmitelaidic acid	53,59
Methyl montanate	52	Myristoleic acid methyl ester	53	Palmitelaidic acid methyl ester	
Methyl moroctate	56				53,54,59
Methyl myristate	50	N		Palmitic acid	50
Methyl myristoleate	53	Nervonic acid	59	Palmitic acid methyl ester	50
Methyl nervonate	59	NOE	35,73	Palmitoleic acid	53
Methyl nonadecanoate	51	Non-Polar Lipid Mixture A	87	Palmitoleic acid methyl ester	53
Methyl nonadecenoate	56	Non-Polar Lipid Mixture B	87	Palmitoyl-2-oleoyl-sn-glycero-3-	
Methyl nonanoate	49	Non-Volatile Acid Mixture	86	phosphorylcholine	43
Methyl octacosanoate	52	Nonadecanoic acid	51	Palmitoyl-2-oleoyl-sn-glycero-3-	
Methyl octadecadienoate		Nonadecanoic acid methyl ester	51	phosphorylglycerol	44
(all <i>cis</i> -9,12)	55	Nonadecenoic acid	56	Palmitoyl serinol, N	6
Methyl octadecadienoate		Nonadecenoic acid methyl ester	56	Palmitoyl-sn-glycero-3-	
(all <i>trans</i> -9,12)	55,60	Nonanoic acid	49	phosphorylcholine	43
Methyl octadecanoate	51	Nonanoic acid methyl ester	49	PDMP, D- <i>threo</i>	38
Methyl octadecatrienoate				PDMP, D,L- <i>threo</i>	37
(all <i>cis</i> -6,9,12)	56	O		PDMP, D,L- <i>erythro</i>	38
Methyl octadecatrienoate		Octacosanoic acid methyl ester	52	PDMP, L- <i>threo</i>	37
(all <i>cis</i> -9,12,15)	56	Octadecadienoic acid-10(E),12(Z)	62	Pelargonic acid	49
Methyl octadecenoate (<i>cis</i> -9)	54	Octadecadienoic acid-10(E),12(Z)		Pelargonic acid methyl ester	49
Methyl octadecenoate (<i>cis</i> -11)	54	methyl ester	62	Pentadecanoic acid	50
Methyl octadecenoate (<i>trans</i> -11)		Octadecadienoic acid-11(Z),13(E)	62	Pentadecanoic acid methyl ester	50
	55,60	Octadecadienoic acid-9(E),11(E)		Phosphatidic acid, egg	41,94
Methyl octadecenoate (<i>trans</i> -9)	54,59	Octadecadienoic acid-9(E),11(E)		Phosphatidylcholine, bovine	40,96
Methyl octanoate	49	methyl ester	62	Phosphatidylcholine, egg	40,96
Methyl oleate	54	Octadecadienoic acid-9(Z),11(E)		Phosphatidylcholine, <i>lyso</i> , egg	40,94
Methyl palmitate	50	Octadecadienoic acid-9(Z),11(E)		Phosphatidylcholine, plant	40,96
Methyl palmitelaidate	54,59	Octadecadienoic acid-9(Z),11(E)		Phosphatidylethanolamine,	
Methyl palmitoleate	53	methyl ester	61,62	bovine	41
Methyl pelargonate	49	Octadecadienoic acid-9(Z),11(E)		Phosphatidylethanolamine,	
Methyl pentadecanoate	50	methyl ester	61,62	egg	41,94
Methyl ricinelaidate	70	Octadecadienoic acid-9(Z),11(Z)	62	Phosphatidylethanolamine,	
Methyl stearate	51	Octadecadienoic acid-9(Z),11(Z)		plant	41,96
Methyl stearidonate	56	methyl ester	62	Phosphatidylinositol 3-phosphate,	
Methyl tetracosanoate	52	Octadecadienoic acid		dipalmitoyl	46
Methyl tetracosenoate	59	(all <i>cis</i> -9,12)	55	Phosphatidylinositol 4-phosphate,	
Methyl tetradecanoate	50	Octadecadienoic acid		dipalmitoyl	46
Methyl tetradecenoate	53	(all <i>trans</i> -9,12)	55,60	Phosphatidylinositol bis-4,5-	
Methyl triacontanoate	53	Octadecadienoic acid methyl ester		phosphate, dioctanoyl	
Methyl tricosanoate	52	(all <i>cis</i> -9,12)	55	(Na ⁺ salt)	47
Methyl tridecanoate	50	Octadecadienoic acid methyl ester		Phosphatidylinositol bis-4,5-	
Methyl undecanoate	49	(all <i>trans</i> -9,12)	55,60	phosphate, dioctanoyl	
Methyl vaccenate (<i>cis</i> -11)	55	Octadecanoic acid	51	(NH ₄ ⁺ salt)	47
Methyl vaccenate (<i>trans</i> -11)	55,59	Octadecanoic acid methyl ester	51	Phosphatidylinositol, plant	41,94
Methyleneoctadecanoic acid	73			Phosphatidylinositol, dipalmitoyl	46

Phosphatidylinositol, plant, soy	41,97	Sphinganine, D- <i>erythro</i> -C ₂₀	4	Sphingosine, N-heptadecanoyl-D- <i>erythro</i>	8
Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl (Na ⁺ salt)	47	Sphinganine, D- <i>threo</i>	4	Sphingosine, N-hexadecanoyl-D- <i>erythro</i>	8
Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl (NH ₄ ⁺ salt)	47	Sphinganine, D,L-C ₁₆	4	Sphingosine, N-hexadecanoyl-D- <i>erythro</i> (C16 sphingoid base)	10
Phosphatidylserine, bovine	40,94	Sphinganine, D,L- <i>erythro</i> -C ₂₀	4	Sphingosine, N-hexanoyl-D- <i>erythro</i>	7
Phosphoglycerides Kit	41	Sphinganine, L- <i>erythro</i>	4	Sphingosine, N-hexanoyl-D- <i>threo</i>	8
Phospholipid (MPL) of <i>Thermoplasma acidophilum</i> (>50%)	47	Sphinganine, L- <i>threo</i>	3,35	Sphingosine, N-hexanoyl-L- <i>erythro</i>	7
Phospholipid (MPL) of <i>Thermoplasma acidophilum</i> (>95%)	47	Sphinganine, N-acetyl-D- <i>erythro</i>	10	Sphingosine, N-hexanoyl-L- <i>threo</i>	7
Phytanic acid	73	Sphinganine, N-dodecanoyl-NBD-D- <i>erythro</i>	14,90	Sphingosine, N-hexanoyl-NBD-D- <i>erythro</i>	13,89
Phytoceramide, N-C12:0-NBD	14,91	Sphinganine, N-dodecanoyl-NBD-L- <i>threo</i>	14,90	Sphingosine, N-hexanoyl-NBD-L- <i>threo</i>	13,90
Phytoceramide, N-C16:0	12	Sphinganine, N-hexanoyl-D- <i>erythro</i>	10	Sphingosine, N-N, dihexyl-D- <i>erythro</i>	6
Phytoceramide, N-C18:0	12	Sphinganine, N-hexanoyl-NBD-D- <i>erythro</i>	14,90	Sphingosine, N,N-dimethyl-D- <i>erythro</i>	5
Phytoceramide, N-C2:0	12	Sphinganine, N-hexanoyl-NBD-L- <i>threo</i>	13,90	Sphingosine, N-nonadecanoyl-D- <i>erythro</i>	9
Phytoceramide, N-C24:0	13	Sphinganine, N-octadecanoyl-D- <i>erythro</i>	11	Sphingosine, N-octadecanoyl-D- <i>erythro</i>	9
Phytoceramide, N-C6:0	12	Sphinganine, N-octanoyl-D- <i>erythro</i>	10	Sphingosine, N-octadecanoyl-D- <i>threo</i>	9
Phytoceramide, N-C6:0-NBD	14,91	Sphinganine, N-(R,S)- <i>alpha</i> -hydroxydodecanoyl-D- <i>erythro</i>	11	Sphingosine, N-octadecanoyl-L- <i>erythro</i>	9
Phytoceramide, N-C8:0	12	Sphinganine, N-(R,S)- <i>alpha</i> -hydroxyhexadecanoyl-D- <i>erythro</i>	11	Sphingosine, N-octadecanoyl-L- <i>threo</i>	9
Phytosphingosine	5	Sphinganine, N-(R,S)- <i>alpha</i> -hydroxyoctadecanoyl-D- <i>erythro</i>	11	Sphingosine, N-octanoyl-D- <i>erythro</i>	8
Phytosphingosine, N-acetyl	12	Sphingolipid Mixture	87	Sphingosine, N-octanoyl-D- <i>threo</i>	8
Phytosphingosine, N-dodecanoyl-NBD	14,91	Sphingomyelin, bovine	15,94	Sphingosine, N-octanoyl-L- <i>threo</i>	8
Phytosphingosine, N-hexadecanoyl	12	Sphingomyelin, bovine buttermilk	15,96	Sphingosine, <i>omega</i> -N-NBD-D- <i>erythro</i> -C ₁₄	2,92
Phytosphingosine, N-hexanoyl	12	Sphingomyelin, D- <i>erythro</i> with 1- ¹³ C-palmitic acid	16,89	Sphingosine, N-pentadecanoyl-D- <i>erythro</i>	8
Phytosphingosine, N-hexanoyl-NBD	14,91	Sphingomyelin, egg	15,97	Sphingosine, N-(R,S)- <i>alpha</i> -hydroxydodecanoyl-D- <i>erythro</i>	9
Phytosphingosine, N-octadecanoyl	12	Sphingomyelin, N-C12:0-NBD	18,91	Sphingosine, N-(R,S)- <i>alpha</i> -hydroxyoctadecanoyl-D- <i>erythro</i>	10
Phytosphingosine, N-octanoyl	12	Sphingomyelin, N-C17:0	16	Sphingosine, N-tetracosanoyl-D- <i>erythro</i>	9
Phytosphingosine, N-tetracosanoyl	13	Sphingomyelin, N-C18:0	16	Sphingosine, N-tetracosenoyl-D- <i>erythro</i>	9
Plant Sterol Mixture	76	Sphingomyelin, N-C2:0	15	Sphingosylphosphorylcholine	17
Plant Sterols Kit	76	Sphingomyelin, N-C20:0-D- <i>erythro</i>	16	Sphingosylphosphorylcholine, D- <i>erythro</i>	17
Polar Lipid Mixture	87	Sphingomyelin, N-C22:0-D- <i>erythro</i>	16	Sphingosylphosphorylcholine, L- <i>threo</i>	17
PPMP, D- <i>threo</i>	38	Sphingosine-1-phosphate, N-hexadecanoyl-D- <i>erythro</i>	18	Sphingosylphosphorylcholine, N-1- ¹³ C-hexadecanoyl	16, 89
PPMP, D,L- <i>erythro</i>	38	Sphingomyelin, N-C6:0	15	Sphingosylphosphorylcholine, N-acetyl	15
PPMP, D,L- <i>threo</i>	37	Sphingomyelin, N-C6:0-NBD	18,91	Sphingosylphosphorylcholine, N-eicosanoyl-D- <i>erythro</i>	16
PPMP, L- <i>threo</i>	38	Sphingomyelin, porcine	15,96	Sphingosylphosphorylcholine, N-docosanoyl-D- <i>erythro</i>	16
Psychosine	20	Sphingosine-1-phosphate, D- <i>erythro</i>	17	Sphingosylphosphorylcholine, N-dodecanoyl-NBD	18,91
Psychosine, N-acetyl	20	Sphingosine, D- <i>erythro</i>	2	Sphingosylphosphorylcholine, N-heptadecanoyl	16
Psychosine, N-octadecanoyl-D ₃₅	20,27,88	Sphingosine, D- <i>erythro</i> -C ₁₂	3	Sphingosylphosphorylcholine, N-hexanoyl	15
Psychosine, N-pentadecanoyl	20	Sphingosine, D- <i>erythro</i> -C ₁₄	2	Sphingosylphosphorylcholine, N-octadecanoyl	16
PUFA-1 Mixture	79	Sphingosine, D- <i>erythro</i> -C ₁₆	3	Sphingosylphosphorylethanolamine, N-acyl	16
PUFA-2 Mixture	79	Sphingosine, D- <i>erythro</i> -C ₂₀	3	Stearic acid	51
PUFA-3 Mixture	79	Sphingosine, D- <i>erythro</i> -C ₂₀	3	Stearic acid methyl ester	51
R		Sphingosine, D- <i>threo</i>	2	Stearidonic acid	56
Rapeseed Oil Reference Mixture	81	Sphingosine, L- <i>erythro</i>	2	Stearidonic acid methyl ester	56
Ricinelaidic acid	70	Sphingosine, L- <i>threo</i>	2		
RM-1 Mixture	81	Sphingosine, N-acetyl-D- <i>erythro</i>	7		
RM-2 Mixture	82	Sphingosine, N-acetyl-D- <i>erythro</i> (C14 sphingoid base)	10		
RM-3 Mixture	82	Sphingosine, N-acetyl-L- <i>erythro</i>	7		
RM-4 Mixture	82	Sphingosine, N-acetyl-L- <i>threo</i>	7		
RM-5 Mixture	82	Sphingosine, N-decanoyl-D- <i>erythro</i>	8		
RM-6 Mixture	82	Sphingosine, N-dodecanoyl-NBD-D- <i>erythro</i>	13,90		
Royal Jelly acid	68	Sphingosine, N-dodecanoyl-NBD-L- <i>threo</i>	13,90		
S					
Safingol	3,35				
Sapienic acid	53				
Sitostanol- <i>beta</i>	77				
Sphinganine•HCl (3-keto)	4				
Sphinganine, D- <i>erythro</i>	3				

Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine	44,63
Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine	44,63
Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine	43,63
Steryl glucosides	77
Stigmastanol	77
Stigmasterol	77
Sulfatide-lyso	23
Sulfatide, N-acetyl	23
Sulfatide, N-C12:0-NBD	24,28,92
Sulfatide, N-C16:0	23
Sulfatide, N-C18:0	23
Sulfatide, N-C18:0-D ₃	23,27,89
Sulfatide, N-C18:1	23
Sulfatide, N-C2:0	23
Sulfatide, N-C24:0	23
Sulfatide, N-C24:1	23
Sulfatide, N-dodecanoyl-NBD	24,28,92
Sulfatide, N-hexadecanoyl	23
Sulfatide, N-lignoceroyl	23
Sulfatide, N-nervonyl	23
Sulfatide, N-octadecanoyl	23

Sulfatide, N-octadecanoyl-D ₃	23,27,89
Sulfatide, N-octadecenoyl	23
Sulfatide, N-oleoyl	23
Sulfatide, N-palmitoyl	23
Sulfatide, N-stearoyl	23
Sulfatide, N-stearoyl-D ₃	23,27,89
Sulfatide, N-tetracosanoyl	23
Sulfatide, tetracosenoyl	23
Sulfatides	22,94

T

Tetracontanoic acid methyl ester	53
Tetracosanoic acid	52
Tetracosanoic acid methyl ester	52
Tetracosenoic acid	59
Tetracosenoic acid methyl ester	59
Tetradecanoic acid	50
Tetradecanoic acid methyl ester	50
Tetradecenoic acid methyl ester	53
Tetrasialoganglioside GQ _{1b}	32,98
THI	39
Tocol- <i>rac</i>	75
Tocopherol, (+)- <i>delta</i>	75
Tocopherol, <i>rac-alpha</i>	74
Tocopherol, <i>rac-beta</i>	74
Tocopherol, <i>rac-gamma</i>	74

Tricosanoic acid	52
Tricosanoic acid methyl ester	52
Tridecanoic acid	50
Tridecanoic acid methyl ester	50
Trimethyltocol-5,7,8	74
Trisialoganglioside GT _{1b}	31,95

U

Undecanoic acid	49
Undecanoic acid methyl ester	49

V

Vaccenic acid (<i>cis</i> -11)	55
Vaccenic acid methyl ester (<i>cis</i> -11)	55
Vaccenic acid methyl ester (<i>trans</i> -11)	55,59
Vaccenic acid (<i>trans</i> -11)	55,59
Volatile Acid Mixture	86

W

WSFA-2 Mixture	85
WSFA-4 Mixture	85

Matreya LLC Ordering Information

Office Hours: Monday - Friday, 8:00am - 4:00pm, Eastern Time

• Ordering and Customer Service

Telephone (Toll Free)
Telephone (worldwide)
Fax (24/7 worldwide)
e-mail
e-commerce website

(800) 342 3595
(814) 359 5060
(814) 359 5062
customerservice@matreya.com
www.matreya.com

• Technical Service

Telephone (worldwide)
Fax (worldwide)
e-mail

(814) 359 5060
(814) 359 5062
techservice@matreya.com

• Terms

Prices and discounts are subject to change without notice. Freight charges are prepaid and added as a separate charge on the invoice. Orders placed are F.O.B. Pleasant Gap, PA. **Net 30 Days.**

• Shipping

Standard orders: FedEx 2nd-Day
Rush Orders: FedEx Priority or Standard overnight
International Orders: FedEx International Priority
Items requiring dry ice for shipping will be charged a \$40 fee.

• Return Policy

All returns must have prior authorization. Items incorrectly ordered by you and returned to Matreya are subject to a 25% (\$25 minimum) restocking charge. The following items may not be returned: frozen products, items which have passed their expiration dates, custom synthesized products and accommodation orders. If a product has been incorrectly sent to you due to an error on our part, a credit will be issued to your account immediately. Please inspect and verify your order upon receipt. No products will be returned after 30 days (assuming proper storage and handling by customer).

• Disclaimer

The information and data included in this catalog are correct and reliable to the best of our knowledge. However, we offer no guarantees and assume no responsibility for this information. No license or immunity under any patent is granted or implied through our sale of any material. Any information sent to us by the customer, relative to his or her interest in Matreya's products, must be submitted voluntarily and without obligation on the part of Matreya.

Matreya LLC
168 Tressler Street
Pleasant Gap, PA 16823 USA
Federal ID No. 20-1237500

International Dealers & Representatives

Australia (South Australia)

Adelab Scientific
36 Holland Street
Thebarton 5031
Tel: 61-88-234-7955
Fax: 61-88-234-7897
Email: info@adelab.com.au
Web: www.adelab.com.au

France

Biovalley
18 Route de Tournan
Marne la Vallée Cedex 3
Conches, 77601
Tel: 33-16-007-2020
Fax: 33-16-007-5051
Email: biovalley@biovalley.fr
Web: www.biovalley.fr

Japan

Techno Chemical Corp
27-9, Honkomagome 1 Chome
Bunkyo-ku, Tokyo 113
Tel: 81-33-947-7310
Fax: 81-33-947-7306
Email: info@technochemical.com
Web: www.technochemical.com

New Zealand

Phenomenex NZ Ltd.
PO Box 31-6-1
Milford, Auckland
Tel: 649-478-0951
Fax: 649-478-0952
Email: info@phenomenex.co.za
Web: www.phenomenex.com

Switzerland

VWR International AG-Life Sciences
Lerzenstrasse 16/18
Dietikon, 8953
Tel: 41-44-745-1436
Fax: 41-44-745-1410
Email: info@ch.vwr.com
Web: www.vwr.com

Canada

MJS Biolynx
PO Box 1150
300 Laurier Blvd
Brockville, Ontario K6V 5W1
Tel: 613-498-2126
Toll free: 888-593-5969
Fax: 613 342 1341
Email: sales@biolynx.ca
Web: www.biolynx.ca

Germany

BIOTREND Chemikalien GmbH
Eupenerstr. 157
Köln, 50933
Tel: 49-221-949-8320
Fax: 49-221-949-8325
Email: info@biotrend.com
Web: www.biotrend.com

Korea

Kim & Friends, Inc.
SK Twintech Tower B-304
345-9 Gasan-dong, Geumcheon-gu
Seoul 153-773
Tel: 82-2-2647-6611
Fax: 82-2-2647-6687
Email: kslee@kimnfriends.co.kr
Web: www.kimnfriends.co.kr

Spain

Teknokroma S.C.C.L.
Camí de Can Calders, 14
Apartado de Correos, 147
Sant Cugat del Vallès
Barcelona ES-08173
Tel: 34-93-674-8800
Fax: 34-93-675-2405
Email: commercial@teknokroma.es
Web: www.teknokroma.es

Taiwan

Super Chroma Enterprise Ltd
5F.226 Roosevelt Rd, Section 5
Taipei ROC, 116
Tel: 88-622-930-7110
Fax: 88-622-930-7112
Email: sc@superchroma.com.tw
Web: www.superchroma.com.tw

China

Shenzhen Bolin Chemical Co. LTD
Rm 1309, No. 1 News Bldg
2 Shennan Zhong Road
Shenzhen, 518027
Tel: 867-556-165-1418 (1428)
Fax: 867-558-209-6552
Email: lh@bolin-lkmlabs.com
Web: www.bolin-lkmlabs.com

Italy

Superchrom S.r.l.
Via C. Menotti, 11
Milano, 20129
Tel: 39-02-738-6315
Fax: 39-027-010-0100
Email: Superchrom@iol.it

Netherlands

Distrilab BV
Olmenlaan 6-C
Leusden, 3833 AV
Tel: 31-33-494-7834
Fax: 31-33-432-1441
Email: info@distrilab.nl
Web: www.distrilab.nl

Sweden

Larodan Fine Chemicals AB
Limhamnsgårdens Alle'9
Malmö, S-21616
Tel: 464-016-4155
Fax: 464-015-5498
Email: info@larodan.se
Web: www.larodan.se

United Kingdom (England)

Universal Biologicals (Cambridge) Ltd.
Passhouse Farmhouse
Papworth St. Agnes
Cambridge CB23 3QU
Tel: 44-148-083-9015
Fax: 44-148-803-1912
Email: info@universalbiologicals.com
Web: www.universalbiologicals.com