

Food and Beverage Standards

Vitamins

Vitamins

For use in HPLC, GC, and other traditional wet chemistry analyses.
All compounds have been thoroughly evaluated to ensure the utmost quality.

Vitamins – neat, unless otherwise noted.

DESCRIPTION	CAS NO.	QTY.	CAT. NO.	PRICE
WATER SOLUBLE VITAMINS				
Thiamine hydrochloride (B1)	67-03-8	1g	47858	
Riboflavin (B2)	83-88-5	1g	47861	
Pyridoxine hydrochloride (B6)	58-56-0	1g	47862	
L-Ascorbic acid	50-81-7	1g	47863	
Nicotinic acid	59-67-6	1g	47864	
Nicotinamide	98-92-0	1g	47865-U	
Folic acid	75708-92-8	500mg	47866	
Pantothenic acid (hemicalcium salt)	137-08-6	1g	47867	
D-Biotin	58-85-5	100mg	47868	
Cyanocobalamin (B12)	68-19-9	100mg	47869	
FAT SOLUBLE VITAMINS				
Retinol acetate	127-47-9	100mg	46958	
Retinol palmitate	79-81-2	100mg	46959-U	
D- α -Tocopherol succinate	4345-03-3	100mg	47782	
DL- α -Tocopherol	10191-41-0	100mg	47783	
rac- β -Tocopherol (50mg/mL in hexane)	148-03-8	1mL	46401-U	
δ -Tocopherol	119-13-1	100mg	47784	
γ -Tocopherol	54-28-4	10mg	47785	
DL- α -Tocopherol acetate	7695-91-2	100mg	47786	
Cholecalciferol (D3)	67-97-0	100mg	47763	
Ergocalciferol (D2)	50-14-6	100mg	47768	
Phylloquinone (K1)	84-80-0	100mg	47773	
Menaquinone (K2)	110232-49-8	100mg	47774	
Menadione (K3)	58-27-5	1000mg	47775	

Food and Beverage Standards

Mycotoxins

Mycotoxins

Our mycotoxin standards are specifically designed for use according to the Official Methods of the Association of Official Analytical Chemists (AOAC). We test all products by spectroscopy and/or HPLC against known standards and previous lots to ensure precision and uniformity, assuring that you are obtaining the finest mycotoxin standards available.

Ordering Note: Due to recent changes in U.S. and international laws, you could need a license to order aflatoxin standards. A company representative will notify you if a license is necessary.

Caution: Mycotoxins may be carcinogenic and, therefore, should be handled only by qualified personnel.

DESCRIPTION	CONCENTRATION/SOLVENT	QTY.	CAT. NO.	PRICE
AFLATOXIN STANDARDS – QUANTITATIVE STANDARDS DESIGNED FOR USE IN ACCORDANCE WITH AOAC METHOD 970.44.				
Aflatoxin B and Aflatoxin G Mixes				
Each ampul contains 1µg B ₁ , 1µg G ₁ , 0.3µg B ₂ , and 0.3µg G ₂ .	Benzene:acetonitrile (98:2) Methanol Methanol	5 x 1mL 5 x 1mL 5mL	46300-U■ 46304-U 46303	
Single Component Aflatoxin Standards				
Aflatoxin B ₁ [1162-65-8]	3µg/mL in benzene:acetonitrile (98:2)	1mL	46323-U	
Aflatoxin B ₂ [7220-81-7]	3µg/mL in benzene:acetonitrile (98:2)	1mL	46324-U	
Aflatoxin G ₁ [1165-39-5]	3µg/mL in benzene:acetonitrile (98:2)	1mL	46325-U	
Aflatoxin G ₂ [7241-98-7]	3µg/mL in benzene:acetonitrile (98:2)	1mL	46326-U	
Aflatoxin M ₁ [6795-23-9]	10µg/mL in acetonitrile	1mL	46319-U	
Aflatoxin M ₂ [6885-57-0]	1µg/mL in acetonitrile	1mL	46910-U	
OTHER MYCOTOXINS				
Deoxynivalenol (DON) [51481-10-8]	200µg/mL in ethyl acetate:methanol (95:5) For use in AOAC Methods 986.17 and 986.16	1mL	46911	
Ochratoxin A [303-47-9]	50µg/mL in benzene:acetic acid (99:1)	1mL	46912	
Ochratoxin B [4825-86-9]	50µg/mL in benzene:acetic acid (99:1) Ochratoxin A and B solutions have been produced according to the guidelines in AOAC Method 973.37	1mL	46913-U	
Patulin [149-29-1]	100µg/mL in chloroform Prepared for use with AOAC Method 974.18	1mL	46914-U	
Zearalenone [17924-92-4]	50µg/mL in acetonitrile AOAC Method 976.22 contains methodology for detecting zearalenone in corn. Method 985.18 lists a method for detecting and quantifying zearalenone and α-zearalenol in corn.	1mL	46916-U	

■ Intended for use in accordance with AOAC Method 971.22.

Food and Beverage Standards

Carbohydrates/Organic Acids/Sugar Alcohols

Carbohydrates/Organic Acids/Sugar Alcohols

Prepared, tested, and packaged using rigorous manufacturing procedures.

Food and Beverage Standards

Alditol Acetates, Sweeteners, Antimicrobials

Alditol Acetates

These are quantitative standards. Concentrations of each component are selected to give a relatively uniform chromatogram for use in TLC, GC, or HPLC analysis.

MIX	COMPONENTS	QTY.	CAT. NO.	PRICE
Alditol Acetate Mix 1	4 components, 50mg/mL total in chloroform Rhamnitol acetate (25%) Ribitol acetate (25%) Fucitol acetate (25%) Arabitol acetate (25%)	1mL	47880-U	
Alditol Acetate Mix 2	4 components, 50mg/mL total in chloroform Mannitol acetate (25%) Glucitol acetate (25%) Galactitol acetate (25%) Inositol acetate (25%)	1mL	47881	

Sweeteners

Sweeteners are classified into two main groups — caloric (nutritive) and non-caloric (non-nutritive). Nutritive sweeteners are metabolized by the body to provide energy. Non-nutritive sweeteners are not metabolized by the body and are unchanged before excretion. The nutritive sweeteners are carbohydrates, or derivatives of carbohydrates, such as sugar alcohols.

The study of sweeteners is increasing, due to the dietary requirements of diabetics and concerns about obesity. We offer a variety of sweeteners that the food industry either is using or is evaluating for potential use. We have evaluated these standards to provide the highest quality for your analytical needs.

COMPOUND	CAS NO.	QTY.	CAT. NO.	PRICE
Acesulfame K	55589-62-3	1g	47134	
Aspartame	22839-47-0	500mg	47135	
Sodium cyclamate	139-05-9	1g	47827	
D-(+)-Glucose	50-99-7	1g	47829	
Sodium saccharin	82385-42-0	1g	47839	
Saccharin (hemicalcium)	6381-91-5	1g	47840	
D-Sorbitol	50-70-4	1g	47841	
Xylitol	87-99-0	1g	47844	

Antimicrobials/Preservatives

These neat preservative standards have been evaluated to ensure that they are of the highest quality. We offer these standards in small quantities; there is no need to purchase a large quantity of chemicals, only to pay expensive disposal charges later.

Neat, each 1g.

COMPOUND	CAS NO.	QTY.	CAT. NO.	PRICE
Benzoic acid	65-85-0	1g	47849	
Sodium benzoate	532-32-1	1g	47850	
Methyl paraben	99-76-3	1g	47889	
Ascorbic acid	110-44-1	1g	47845	
Potassium sorbate	590-00-1	1g	47848	

Food and Beverage Standards

Lipids

Lipid Standards


Fatty Acid Methyl Esters (FAMEs)

A Word on Nomenclature - Common names are used in this catalog where brevity does not sacrifice clarity. Geneva names are used where possible.


Saturated and unsaturated fatty acids are named according to the number of carbon atoms of the parent hydrocarbon chain. Saturated fatty acids are named according to the modified Geneva system, by replacing the terminal "e" of the parent hydrocarbon with the "oic" (e.g., decane to decanoic). The common names of most of these fatty acids are also listed.

Unsaturated fatty acids are named by replacing the "e" of the corresponding unsaturated hydrocarbon with the suffix "oic" (e.g., to decenoic). The number of multiple double bonds is indicated by adding dienoic, and so on to the hydrocarbon name. For example, the 18 carbon chain acid with three double bonds is named octadecatrienoic.

Double bonds are also designated by position and geometric configuration. Naturally occurring fatty acids are usually ~~confined~~
configuration, unless stated ~~trans~~. Octadecenoic acid with the double bond in the nine position has ~~both~~ common name, oleic acid) and trans(elaidic acid) forms. The simplest way to name double-bond positions is to count carbons, starting with the carboxyl carbon, until you reach the double bond. Thus, oleic acid named by the Geneva system is octadecenoic acid:



and linoleic acid is cis9,12-octadecadienoic acid:



In the product descriptions for lipid mixes, chain length, followed by the number of double bonds, is indicated in parentheses component names. For example, linolenic acid, which has a chain length of 18 and 3 double bonds, is listed as:

Linolenic acid (18:3)

C18 FAME Isomer Mix

MIX	COMPONENTS	WT. %	QTY.	CAT. NO.	PRICE
Linoleic Acid Methyl Ester cis/trans Isomer Mix	4 components, 10mg/mL in methylene chloride		1mL	47791	
	trans-9,trans-12-Octadecadienoic acid methyl ester	50			
	cis-9,trans-12-Octadecadienoic acid methyl ester	20			
	trans-9,cis-12-Octadecadienoic acid methyl ester	20			
	cis-9,cis-12-Octadecadienoic acid methyl ester	10			

C18 cis/trans Fatty Acids/FAMEs

10mg/mL in heptane.

DESCRIPTION	CAS NO.	QTY.	CAT. NO.	PRICE
cis-6-Octadecenoic methyl ester (Petroselinic)	2777-58-4	1mL	47198	
trans-6-Octadecenoic methyl ester (Petroselaidic)	-	1mL	47199	
cis-7-Octadecenoic methyl ester	2278-59-3	1mL	46900-U	
cis-9-Octadecenoic methyl ester (Oleic)	112-62-9	1mL	46902-U	
trans-9-Octadecenoic methyl ester (Elaidic)	2462-84-2	1mL	46903	
cis-11-Octadecenoic methyl ester (cis-Vaccenic)	1937-63-9	1mL	46904	
trans-11-Octadecenoic methyl ester (trans-Vaccenic)	6198-58-9	1mL	46905-U	
cis-12-Octadecenoic methyl ester	2733-86-0	1mL	46906	
trans-12-Octadecenoic methyl ester	20221-23-2	1mL	46907-U	
trans-13-Octadecenoic methyl ester	42199-38-2	1mL	46909	
cis-15-Octadecenoic methyl ester	10411-39-9	1mL	46953	
cis-9,12-Octadecadienoic methyl ester	112-63-0	1mL	46950-U	
trans-9,12-Octadecadienoic methyl ester	2566-97-4	1mL	46951-U	
cis-5,8,11,14,17-Eicosapentaenoic methyl ester	2734-47-6	1mL	47571-U	
cis-4,7,10,13,16,19-Docosahexaenoic methyl ester	301-01-9	1mL	47570-U	
cis-7,10,13,16,19-Docosapentaenoic methyl ester	108698-02-8	1mL	47563-U	

Food and Beverage Standards

Lipids

Highly Characterized Reference Oils

We offer highly characterized common reference oil samples for use as controls or check samples for fatty acid methyl ester (FA) analyses. These samples provide an excellent means of standardizing your lipid procedures and comparing your results to others. Certificate of Analysis is provided with each oil sample.

1g, packed in amber ampul under nitrogen.

DESCRIPTION	CAS NO.	WT.	CAT. NO.	PRICE
Canola oil	120962-03-0	1g	46961	
Coconut oil	8001-31-8	1g	46949	
Corn oil	8001-30-7	1g	47112-U	
Cottonseed oil	8001-29-4	1g	47113	
Flaxseed oil	8001-26-1	1g	47559-U	
Lard oil	8016-28-2	1g	47115-U	
Menhaden fish oil	8002-50-4	1g	47116	
Menhaden fish oil, partially hydrogenated (PHMO)	—	1g	47117	
Olive oil – refined	8001-25-0	1g	47118	
Palm oil	8002-75-3	1g	46962	
Peanut oil	8002-03-7	1g	47119	
Safflower oil	8001-23-8	1g	47120-U	
Soybean oil	8001-22-7	1g	47122	
Sunflower seed oil	8001-21-6	1g	47123	

Supelco 37 Component FAME Mix

This fatty acid methyl ester (FAME) reference standard is carefully prepared by weight. The weight percentage of each component indicated. Each ampul contains 10mg/mL of the FAME reference standard mix in methylene chloride.

DESCRIPTION	QTY.	CAT. NO.	PRICE
Supelco 37 Component FAME Mix	1mL	47885-U	
Component FAME 1. Butyric (C4:0) 4 2. Caproic (C6:0) 4 3. Caprylic (C8:0) 4 4. Capric (C10:0) 4 5. Undecanoic (C11:0) 2 6. Lauric (C12:0) 4 7. Tridecanoic (C13:0) 2 8. Myristic (C14:0) 4 9. Myristoleic (C14:1) 2 10. Pentadecanoic (C15:0) 2 11. cis-10-Pentadecenoic (C15:1) 2 12. Palmitic (C16:0) 6 13. Palmitoleic (C16:1) 2 14. Heptadecanoic (C17:0) 2 15. cis-10-Heptadecenoic (C17:1) 2 16. Stearic (C18:0) 4 17. Elaidic (C18:1n9t) 2 18. Oleic (C18:1n9c) 4 19. Linoleaidic (C18:2n6t) 2 20. Linoleic (C18:2n6c) 2 21. Arachidic (C20:0) 4 22. γ -Linolenic (C18:3n6) 2 23. cis-11-Eicosenoic (C20:1) 2 24. Linolenic (C18:3n3) 2 25. Heneicosanoic (C21:0) 2 26. cis-11,14-Eicosadienoic (C20:2) 2 27. Behenic (C22:0) 4 28. cis-8,11,14-Eicosatrienoic (C20:3n6) 2 29. Erucic (C22:1n9) 2 30. cis-11,14,17-Eicosatrienoic (C20:3n3) 2 31. Arachidonate (C20:4n6) 2 32. Tricosanoic (C23:0) 2 33. cis-13,16-Docosadienoic (C22:2) 2 34. Lignoceric (C24:0) 4 35. cis-5,8,11,14,17-Eicosapentaenoic (C20:5n3) 2 36. Nervonic (C24:1) 2 37. cis-4,7,10,13,16,19-Docosahexaenoic (C22:6n3) 2			

Column: SP-2560, 100m x 0.25mm ID, 0.20 μ m film

Cat. No.: 24056

Oven: 140°C (5 min) to 240°C at 4°C/min

Carrier: helium, 20cm/sec

Det.: FID, 260°C

Inj.: 1 μ L Cat. No. 47885-U, split 100:1, 260°C

Order: 1.800.325.3010 Technical Service: 1.800.359.3041 Web: www.sigmapelridg.com/supelco

Chemical Standards

SUPELCO

Food and Beverage Standards

Lipids

AOCS Animal and Vegetable Reference Mixes

The quantitative mixes listed here conform to the requirements of American Oil Chemists' Society (AOCS) Method Ce 1-62. The composition of each mix is similar to the fatty acid distribution of certain oils (see Table 1).

- RM-1 – Corn, cottonseed, kapok, poppyseed, rice, safflower, sesame, soybean, sunflower, and walnut oils
- RM-2 – Hempseed, linseed, perilla, and rubberseed oils
- RM-3 – Mustard seed, peanut, and rapeseed oils
- Rapeseed Oil Reference Mix Modern low erucic acid oils
- RM-4 – Neatsfoot, olive, and teaseed oils
- RM-5 – Babassu, coconut, ouri-curi, and palm kernel oils
- RM-6 – Lard, beef tallow, mutton tallow, and palm oil

NHI/NIH Fatty Acid Methyl Ester Reference Mixes

Quantitative standards, identical in composition to those developed and distributed for several years by the National Institutes of Health (Horning, E.C., et al., J. Lipid Research, 5:20-27, 1964). Each mix (see Table 1) was designed to test part of the chromatographic system to ensure its reliability for quantitative analyses.

GLC Standard Mixes

These quantitative mixes are useful for determining relative retention times and approximating response factors.

Table 1. Specifications for Quantitative Products
Each mix is carefully prepared by weight percent.

MIX	CAT. NO.	METHYL ESTER (% COMPOSITION BY WEIGHT)																		
Qty.: 100mg each, neat Storage Temp.: -0 C																				
PRICE		C8:0 (caprylate)	C9:0 (nonanoate)	C10:0 (caprate)	C11:0 (undecanoate)	C12:0 (laurate)	C13:0 (tridecanoate)	C14:0 (myristate)	C15:0 (pentadecanoate)	C16:0 (palmitate)	C16:1 (palmitoleate)	C17:0 (heptadecanoate)	C18:0 (stearate)	C18:1 (oleate)	C18:2 (linoleate)	C18:3 (linolenate)**	C19:0 (nonadecanoate)	C20:0 (arachidate)	C20:1 (eicosenoate)	
RM-1	O7006-1AMP								6.0			3.0	35.0	50.0	3.0		3.0			
RM-2	O7131-1AMP									7.0			5.0	18.0	36.0	34.0				
RM-3	O7256-1AMP							1.0		4.0			3.0	45.0	15.0	3.0		3.0		
Rapeseed	O7756-1AMP							1.0		4.0			3.0	60.0	12.0	5.0		3.0	1.0	
RM-4	O7381-1AMP								11.0			3.0	80.0	6.0						
RM-5	O7506-1AMP	7.0	5.0	48.0	15.0			7.0			3.0	12.0	3.0							
RM-6	O7631-1AMP					2.0	30.0	3.0		14.0	41.0	7.0	3.0							
PRICE																				
NHI-C	O8256-1AMP	1.5	3.0	6.0	12.0	19.4		24.9								33.2				
NHI-D	O8381-1AMP					11.8	23.6	6.9		13.1	44.6									
NHI-F	O8631-1AMP					2.5	4.2		7.3							13.6		25.4	47.0	
NHI-A	O7881-1AMP					25.0	10.0		65.0											
NHI-B	O8131-1AMP					4.0	40.0		56.0											
NHI-E	O8506-1AMP	6.3	9.1	12.1	23.3	49.2														
PRICE																				
GLC-10	1891-1AMP							20.0		20.0	20.0	20.0								
GLC-20	1892-1AMP								20.0		20.0	20.0	20.0			20.0				
GLC-30	1893-1AMP	20.0	20.0	20.0	20.0	20.0		20.0												
GLC-40	1895-1AMP								25.0		25.0					25.0		25.0	25.0	
GLC-50	1894-1AMP									25.0		25.0					25.0		25.0	25.0
GLC-70	1897-1AMP	20.0	20.0	20.0	20.0	20.0				25.0		25.0								
GLC-80	1898-1AMP							20.0	20.0	20.0	20.0	20.0					20.0		20.0	
GLC-90	1896-1AMP							20.0		20.0			20.0				20.0		20.0	
GLC-100	1899-1AMP												20.0				20.0	20.0	20.0	20.0

**Stability problems arise with international shipment of mixtures containing linolenate. When ordering such mixtures from outside the continental US or central Europe, please check with your local dealer regarding ordering practices.

Food and Beverage Standards

Lipids

Grain Fatty Acid Methyl Ester Mix

This fatty acid methyl ester (FAME) mixture is carefully prepared by weight. The weight percentage of each component is indicated. Each ampul contains 10mg/mL of the FAME reference standard mix in methylene chloride.

DESCRIPTION		QTY.	CAT. NO.	PRICE
Grain Fatty Acid Methyl Ester Mix		1mL	47801	
Component FAME				
1. Caprylic (C8:0)	Wt. %	1.9		
2. Capric (C10:0)		3.2		
3. Lauric (C12:0)		6.4		
4. Tridecanoic (C13:0)		3.2		
5. Myristic (C14:0)		3.2		
6. Myristoleic (C14:1n9c)		1.9		
7. Pentadecanoic (C15:0)		1.9		
8. Palmitic (C16:0)		13.0		
9. Palmitoleic (C16:1n9c)		6.4		
10. Heptadecanoic (C17:0)		3.2		
11. Stearic (C18:0)		6.5		
12. Elaidic (C18:1n9t)		2.6		
13. Oleic (C18:1n9c)		19.6		
14. Linoleic (C18:2n6c)		13.0		
15. Arachidic (C20:0)		1.9		
16. cis-11-Eicosenoic (C20:1)		1.9		
17. Linolenic (C18:3n3)		6.4		
18. Behenic (C22:0)		1.9		
19. Erucic (C22:1n9)		1.9		

797-0004

Column: SP-2560, 100m x 0.25mm ID, 0.20µm film

Cat. No.: 24056

Oven: 140°C (5 min) to 240°C at 4°C/min

Carrier: helium, 20cm/sec

Det.: FID, 260°C

Inj.: 1µL, 260°C, split 100:1

Polyunsaturated Fatty Acid (PUFA) Methyl Esters

These are complex qualitative standard mixtures. Because they are extracted from natural materials, relative peak sizes may vary from lot to lot.

DESCRIPTION		QTY.	CAT. NO.	PRICE
PUFA NO. 1				
Marine Source	Typically contains each FAME listed	100mg	47033	
	C14:0 C18:2 ω6 C16:0 C18:4 ω3 C16:1 ω7 C20:1 ω9 C18:1 ω9 C20:5 ω3 C18:1 ω7 C22:1 ω11			
PUFA NO. 2				
Animal Source	Typically contains each FAME listed	100mg	47015-U	
	C14:0 C18:1 ω7 C16:0 C18:2 ω6 C16:1 ω7 C18:3 ω6 C18:0 C18:3 ω3 C18:1 ω9 C20:3 ω6			
PUFA NO. 3				
From Menhaden Oil	Typically contains each FAME listed	100mg	47085-U	
	C14:0 C18:1 ω7 C16:0 C18:2 ω6 C16:1 ω7 C18:2 ω4 C16:2 ω4 C18:3 ω4 C16:4 ω1 C18:3 ω3 C18:0 C18:4 ω3 C18:1 ω9 C20:1 ω9 C20:4 ω6 C20:5 ω3 C22:4 ω6 C22:5 ω3 C22:6 ω3			

Note: In the ω-x classification, carbon atoms are numbered from the terminal methyl group to the first carbon of the ethylenic bond.

X denotes the position at the double bond closest to the terminal methyl group.

Order: 1.800.325.3010 Technical Service: 1.800.359.3041 Web: www.sigma-aldrich.com/supelco

Chemical Standards

SUPELCO

Food and Beverage Standards

Lipids

Qualitative Methyl Ester Mixes

These mixes are prepared from unsaturated medium-chain fatty acids. They are useful for establishing retention times~~s~~ and for per identification. All components are 99% pure by GLC and/or TLC, unless otherwise stated.

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
SATURATED FAMES				
Fatty Acid Methyl Esters, Saturated Straight Chains Kit	10 individually packaged FAMEs, each 1g Caproic (6:0) Caprylic (8:0) Capric (10:0) Lauric (12:0) Myristic (14:0)	Palmitic (16:0) Stearic (18:0) Arachidic (20:0) Behenic (22:0) Lignoceric (24:0)		ME10-1KT
Fatty Acid Methyl Esters, Saturated Straight Chains Kit	19 individually packaged FAMEs, each 1g Caproic (6:0) Heptanoic (7:0) Caprylic (8:0) Nonanoic (9:0) Capric (10:0) Undecanoic (11:0) Lauric (12:0) Tridecanoic (13:0) Myristic (14:0) Pentadecanoic (15:0)	Palmitic (16:0) Heptadecanoic (17:0) Stearic (18:0) Nonadecanoic (19:0) Arachidic (20:0) Heneicosanoic (21:0) Behenic (22:0) Tricosanoic (23:0) Lignoceric (24:0)		ME19-1KT
Fatty Acid Methyl Esters, Saturated Straight Chains Kit	7 individually packaged FAMEs, each 100mg Pentacosanoic (25:0) Hexacosanoic (26:0) Heptacosanoic (27:0) Octacosanoic (28:0)	Nonacosanoic (29:0) Triacanthanoic (30:0) Hentriacontanoic (31:0), approx. 98% pure		ME7-1KT
UNSATURATED FAMES				
Fatty Acid Methyl Esters, Unsaturated Kit	14 individually packaged FAMEs in the amounts indicated Myristoleic (14:1), 100mg Palmitoleic (16:1), 100mg Petroselinic (18:1), 100mg Oleic (18:1), 1g Elaidic (18:1), 500mg cis-Vaccenic (18:1,cis), 100mg Linoleic (18:2), 1g	Linolealidic (18:2), 100mg Linolenic (18:3, trans), 100mg cis-11-Eicosenoic (20:1), 100mg Arachidonic (20:4), 100mg Erucic (22:1), 100mg cis-4,7,10,13,16,19-Docosahexaenoic (22:6), 100mg Nervonic (24:1), 100mg		ME14-1KT
Fatty Acid Methyl Ester Mix	4 components, approx. 10mg each 11-Eicosenoic (20:1) 11,14-Eicosadienoic (20:2) (98% pure) Arachidonic (20:4) 5,8,11,14,17-Eicosapentaenoic (20:5)		40mg	18912-1AMP
Fatty Acid Methyl Ester Mix	5 components, approx. 10mg each 11-Eicosenoic (20:1) 11,14-Eicosadienoic (20:2) (98% pure) 11,14,17-Eicosatrienoic (20:3) (97-99% pure) Arachidonic (20:4) 5,8,11,14,17-Eicosapentaenoic (20:5)		50mg	18913-1AMP

Food and Beverage Standards

Lipids

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
UNSATURATED FAMES (CONTD.)				
Fatty Acid Methyl Ester Mix	6 components, total of 100mg neat mixture weight percentages indicated Stearic acid m.e. (C18:0), 10% Oleic acid m.e. (C18:1, cis-9), 20% Elaidic acid m.e. (C18:1, trans-9), 20%	100mg	18916-1AMP	
				Linoleic acid m.e. (C18:2 cis-9,12), 20% Linolelaidic acid m.e. (C18:2, trans-9,12), 20% Arachidic acid m.e. (C20:0), 10%
Fatty Acid Methyl Ester Mix	10 components, total of 100mg neat mixture weight percentages indicated Myristic acid m.e. (C14:0), 4% Palmitic acid m.e. (C16:0), 10% Stearic acid m.e. (C18:0), 6% Oleic acid m.e. (C18:1, cis-9), 25% Elaidic acid m.e. (C18:1, trans-9), 10%	100mg	18917-1AMP	
				Linoleic acid m.e. (C18:2 cis-9,12), 34% Linolelaidic acid m.e. (C18:2, trans-9,12), 2% Linolenic acid m.e. (C18:3, cis-9,12,15), 5% Arachidic acid m.e. (C20:0), 2% Behenic acid m.e. (C22:0), 2%
Fatty Acid Methyl Ester Mix R: 36/37/38 S: 26-36	14 components, total of 100mg neat mixture weight percentages indicated Caprylic acid m.e. (C8:0), 8% Capric acid m.e. (C10:0), 8% Lauric acid m.e. (C12:0), 8% Myristic acid m.e. (C14:0), 8% Palmitic acid m.e. (C16:0), 11% Palmitoleic acid m.e. (C16:1, cis-9), 5% Stearic acid m.e. (C18:0), 8%	100mg	18918-1AMP	
				Oleic acid m.e. (C18:1, cis-9), 5% Linoleic acid m.e. (C18:2 cis-9,12), 5% Linolenic acid m.e. (C18:3, cis-9,12,15), 5% Arachidic acid m.e. (C20:0), 8% Behenic acid m.e. (C22:0), 8% Erucic acid m.e. (C22:1, cis-13), 5% Lignoceric acid m.e. (C24:0), 8%
Fatty Acid Methyl Ester Mix R: 36/37/038 S: 26-36	37 components, total of 100mg neat mixture weight percentages indicated Butyric acid m.e. (C4:0), 4% Caproic acid m.e. (C6:0), 4% Caprylic acid m.e. (C8:0), 4% Capric acid m.e. (C10:0), 4% Undecanoic acid m.e. (C11:0), 2% Lauric acid m.e. (C12:0), 4% Tridecanoic acid m.e. (C13:0), 2% Myristic acid m.e. (C14:0), 4% Myristoleic acid m.e. (C14:0, cis-9), 2% Pentadecanoic acid m.e. (C15:0), 2% cis-10 Pentadecenoic acid m.e. (C15:1) (approx. 98%), 2% Palmitic acid m.e. (C16:0), 6% Palmitoleic acid m.e. (C16:1, cis-9), 2% Heptadecanoic acid m.e. (C17:0), 2% cis-10 Heptadecenoic acid m.e. (C17:1), 2% Stearic acid m.e. (C18:0), 4% Oleic acid m.e. (C18:1, cis-9), 4% Elaidic acid m.e. (C18:1, trans-9), 2%	100mg	18919-1AMP	
				Linoleic acid m.e. (C18:2 cis-9,12), 2% Linolelaidic acid m.e. (C18:2, trans-9,12), 2% Linolenic acid m.e. (C18:3, cis-9,12,15), 2% g-Linolenic acid m.e. (C18:3, cis-6,9,12), 2% Arachidic acid m.e. (C20:0), 4% cis-11-Eicosenoic acid m.e. (C20:1), 2% cis-11,14-Eicosadienoic acid m.e. (C20:2), (98%), 2% cis-11,14,17-Eicosatrienoic acid m.e. (C20:3) (97-99%), 2% cis-8,11,14-Eicosatrienoic acid m.e. (C20:3), 2% Arachidonic acid m.e. (C20:4, cis-5,8,11,14), 2% cis-5,8,11,14,17-Eicosapentaenoic acid m.e. (C20:5), 2% Heneicosanoic acid m.e. (C21:0), 2% Behenic acid m.e. (C22:0), 4% Erucic acid m.e. (C22:1, cis-13), 2% cis-13,16-Docosadienoic acid m.e. (C22:2), 2% cis-4,7,10,13,16,19-Docosahexaenoic acid m.e. (C22:6), 2% Tricosanoic acid m.e. (C23:0), 2% Lignoceric acid m.e. (C24:0), 4% Nervonic acid m.e. (C24:1, cis-15), 2%
Fatty Acid Methyl Ester Mix	19 components, total of 100mg neat mixture weight percentages indicated Caprylic acid m.e. (C8:0), 1.9% Capric acid m.e. (C10:0), 3.2% Lauric acid m.e. (C12:0), 6.4% Tridecanoic acid m.e. (C13:0), 3.2% Myristic acid m.e. (C14:0), 3.2% Myristoleic acid m.e. (C14:0, cis-9), 1.9% Pentadecanoic acid m.e. (C15:0), 1.9% Palmitic acid m.e. (C16:0), 13.0% Palmitoleic acid m.e. (C16:1, cis-9), 6.4% Heptadecanoic acid m.e. (C17:0), 3.2%	100mg	18920-1AMP	
				Stearic acid m.e. (C18:0), 6.5% Oleic acid m.e. (C18:1, cis-9), 19.6% Elaidic acid m.e. (C18:1, trans-9), 2.6% Linoleic acid m.e. (C18:2 cis-9,12), 13.0% Linolenic acid m.e. (C18:3, cis-9,12,15), 6.4% Arachidic acid m.e. (C20:0), 1.9% cis-11-Eicosenoic acid m.e. (C20:1), 1.9% Behenic acid m.e. (C22:0), 1.9% Erucic acid m.e. (C22:1, cis-13), 1.9%

Food and Beverage Standards

Lipids

Qualitative Fatty Acid Mixes

These mixes are prepared from unsaturated medium-chain fatty acids. They are useful for establishing retention times and for identification. All components are 99% pure by GLC and/or TLC, unless otherwise stated.

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
SATURATED FATTY ACIDS				
Fatty Acids, Even Carbon Straight Chains Kit	10 individually packaged fatty acids in the amounts indicated Caproic acid (C6:0), 10mL Myristic acid (C14:0), 10g Arachidic acid (C20:0), 10g Caprylic acid (C8:0), 10mL Palmitic acid (C16:0), 10g Behenic acid (C22:0), 10g Capric acid (C10:0), 10g Stearic acid (C18:0), 10g Lignoceric acid (C24:0), 10g Lauric acid (C12:0), 10g		EC10-1KT	
Fatty Acids, Even Carbon Straight Chains Kit	10 individually packaged fatty acids in the amounts indicated Caproic acid (C6:0), 10mL Myristic acid (C14:0), 10g Arachidic acid (C20:0), 5g Caprylic acid (C8:0), 10mL Palmitic acid (C16:0), 10g Behenic acid (C22:0), 5g Capric acid (C10:0), 10g Stearic acid (C18:0), 5g Lignoceric acid (C24:0), 1g Lauric acid (C12:0), 10g		EC10A-1KT	
Fatty Acids, Odd Carbon Straight Chains Kit	9 individually packaged fatty acids, 1g each Heptanoic acid (C7:0) (approx. 98% pure) Nonanoic acid (C9:0) (97-99% pure) Undecanoic acid (C11:0) Tridecanoic acid (C13:0) Pentadecanoic acid (C15:0)		OC9-1KT	
Volatile Acid Standard Mix	10 components, 10mM each in 100mL deionized water Formic acid Butyric acid Isocaprylic acid (4-Methyl-n-valeric acid) Acetic acid Isovaleric acid Hexanoic acid (n-Caprylic acid) Propionic acid n-Valeric acid Heptanoic acid	100mL	46975-U	
UNSATURATED FATTY ACIDS				
Fatty Acid Mix	4 fatty acids, approx. 10mg each 6,9,12,15-Octadecatetraenoic acid (18:4) (90% pure) Arachidonic acid (20:4) 5,8,11,14,17-Eicosapentaenoic acid (20:5) 4,7,10,13,16,19-Docosahexaenoic acid (22:6)	40mg	17824-1AMP	
Fatty Acids, Unsaturated Kit	10 individually packaged fatty acids, 100mg each Palmitoleic acid (16:1) Linoleic acid (18:2) Erucic acid (22:1) Elaidic acid (18:1, trans) Linolenic acid, ~98% (18:3) Docosahexaenoic acid (22:6) Oleic acid (18:1) Arachidonic acid (20:4) Nervonic acid (24:1) Petroselinic acid (18:1)		UN10-1KT	

Mono-, Di-, and Triglycerides

These standards are approximately equal weight mixtures. They are qualitative standards, useful in determining relative retention times and establishing approximate response factors.

Each mix is prepared by weight, and the composition verified by gas and/or thin layer liquid chromatography. The weight percent of each component is indicated. Each product contains 100mg total weight of lipid unless otherwise indicated.

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
QUALITATIVE MIXES				
Triglyceride Mix	5 components, approx. 20mg each Tricaprylin (8:0) Trilaurin (12:0) Tripalmitin (16:0) Tricaprin (10:0) Trimyristin (14:0)	100mg	17811-1AMP	
Triglyceride Mix	3 components, approx. 33 1/3 mg each Triolein (18:1, cis-9) Tripetroselinin (18:1, cis-6) Trielaidin (18:1, trans-9)	100mg	17818-1AMP	
Olive Oil Standard Mix▲	5 components at the concentrations indicated Trilinolein (18:2), 2% 1,2-Dilinoleoyl-3-oleoyl-rac-glycerol (18:2, 18:2, 18:1), 4% Triolein (18:1), 60% 1,2-Dioleoyl-3-palmitoyl-rac-glycerol (18:1, 18:1, 16:0), 30% 1,2-Dioleoyl-3-stearoyl-rac-glycerol (18:1, 18:1, 18:0), 4%	50mg	TRI5-1AMP	
Mono-, Di-, and Triglyceride Mix	4 components, approx. 10mg each Monolein (18:1) 1,3-Diolein (18:1) 1,2-Diolein (18:1) Triolein (18:1)	40mg	1787-1AMP	

▲ Component percentages are target values. Actual composition may vary from lot to lot. Lot specific analysis is available upon request from our Technical Service department.

Food and Beverage Standards

Lipids

609

MIXES AND SOLUTIONS	COMPONENTS	CAT. NO.	PRICE
QUALITATIVE MIXES (CONT'D.)			
Triglycerides Kit	10 individually packaged triglycerides, 50mg each Trilaurin (C12:0) 1,2-Dilauroyl-3-myristoyl-rac-glycerol (C12:0/C12:0/C14:0) 1,2-Dimyristoyl-3-lauroyl-rac-glycerol (C14:0/C14:0/C12:0) Trimyristin (C14:0) 1,2-Dimyristoyl-3-palm itoyl-rac-glycerol (C14:0/C14:0/C16:0)	1,2-Dipalmitoyl-3-myristoyl-rac-glycerol (C16:0/C16:0/C14:0) Tri palmitin (C16:0) 1,2-Distearoyl-3-myristoyl-rac-glycerol (C18:0/C18:0/C14:0) 1,2-Distearoyl-3-palmitoyl-rac-glycerol (C18:0/C18:0/C16:0) Tristearin (C18:0)	TRI10-1KT
Triglycerides Kit	19 individually packaged triglycerides in quantities indicated Shipped in dry ice. Triacetin (C2:0), 100mg Tributyrin (C4:0), 100mg Tricaproin (C6:0), 1mL Tricaprylin (C8:0), 0.5mL Tricaprin (C10:0), 100mg Trilaurin (C12:0), 100mg Trimyristin (C14:0), 1g Tripalmitin (C16:0), 100mg Tripalmitolein (C16:1, cis-9), approx. 98%, 100mg Tripetroselinin (C18:1, cis-6), 100mg	Tristearin (C18:0), 100mg Triolein (C18:1, cis-9), 100mg Trielaidin (C18:1, trans-9), 100mg Trilinolein (C18:2, cis-9,12), 100mg Trilinolenin (C18:3, cis-9,12,15), approx. 98%, 100mg Triarachidin (C20:0), 100mg Tri-cis-11-eicosenoic (C20:1), 100mg Tribehenin (C22:0), 100mg Trierucin (C22:1, cis-13), 100mg	TRI19-1KT
Triglycerides, Saturated, Even Carbon Chains Kit	11 individually packaged triglycerides in quantities indicated Triacetin (2:0), 100mg Tributyrin (4:0), 100mg Tricaproin (6:0), 1mL Tricaprylin (8:0), 1mL Tricaprin (10:0), 100mg Trilaurin (12:0), 100mg	Trimyristin (14:0), 100mg Tripalmitin (16:0), 100mg Tristearin (18:0), 100mg Triarachidin (20:0), 100mg Tribehenin (22:0), 100mg	TRI11-1KT
Mono-, Di-, and Triglycerides Kit	12 individually packaged glycerides, 100mg each 1-Monolauroyl-rac-glycerol (12:0) Dilaurin (12:0) Trilaurin (12:0) 1-Monomyristoyl-rac-glycerol (14:0) Dimyristin (14:0)	Trimyristin (14:0) 1-Monopalmitoyl-rac-glycerol (16:0) Dipalmitin (16:0) Tripalmitin (16:0) 1-Monostearoyl-rac-glycerol (18:0) Distearin (18:0) Tristearin (18:0)	MDT12-1KT
Phospholipid Mixture for HPLC R: 45-46-23/24/25-36/37/38 S: 45-26-36/37/39-23	4 components in 2.0mL of a chloroform solution L- α -Phosphatidylcholine, 1.5mg/mL L- α -Phosphatidylinositol, ammonium salt, 0.9mg/mL L- α -Phosphatidylethanolamine, 1.2mg/mL L- α -Lysophosphatidylcholine, 0.3mg/mL	P3817-1VL	

Sterols

Sterols make up the majority of the unsaponifiable matter in vegetable and animal fats. Animal fats contain mostly cholesterol; most vegetable fats contain only traces of this sterol. Plant sterols are collectively called phytosterols.

These standards are not corrected for purity. Packed in amber ampul under nitrogen.

DESCRIPTION	CAS NO.	CONCENTRATION	QTY.	CAT. NO.	PRICE
5- α -Cholestane	481-21-0	10mg/mL in chloroform	1mL	47124	
Campesterol	474-62-4	100 μ g/mL in chloroform	1mL	47126	
Cholesterol	57-88-5	10mg/mL in chloroform	1mL	47127-U	
Dihydrocholesterol	80-97-7	10mg/mL in chloroform	1mL	47129	
Ergosterol	57-87-4	10mg/mL in chloroform	1mL	47130-U	
Stigmasterol	83-48-7	10mg/mL in chloroform	1mL	47132	
β -Sitosterol	83-46-5	100 μ g/mL in chloroform	1mL	47133	

*Purity approx. 60%.

Order: 1.800.325.3010 Technical Service: 1.800.359.3041 Web: www.sigma-aldrich.com/supelco

Chemical Standards

SUPELCO

Bacterial Identification Standards

Lipids

Standard for Bacteria Identification

This is a qualitative standard of bacterial acid methyl esters in methyl caproate (10mg/mL total concentration).

DESCRIPTION	QTY.	CAT. NO.	PRICE
Bacterial Acid Methyl Ester (BAME)	1mL	47080-U	
1. 11:0 Me. undecanoate 2. 2-OH 10:0 Me. 2-hydroxydecanoate 3. 12:0 Me. dodecanoate 4. 13:0 Me. tridecanoate 5. 2-OH 12:0 Me. 2-hydroxydodecanoate 6. 3-OH 12:0 Me. 3-hydroxydodecanoate 7. 14:0 Me. tetradecanoate 8. i-15:0 Me. 13-methyltetradecanoate 9. a-15:0 Me. 12-methyltetradecanoate 10. 15:0 Me. pentadecanoate 11. 2-OH 14:0 Me. 2-hydroxytetradecanoate 12. 3-OH 14:0 Me. 3-hydroxytetradecanoate 13. i-16:0 Me. 14-methylpentadecanoate 14. 16:1 ⁹ Me. cis-9-hexadecenoate 15. 16:0 Me. hexadecanoate 16. i-17:0 Me. 15-methylhexadecanoate 17. 17:0 ^A Me. cis-9,10-methylenehexadecanoate 18. 17:0 Me. heptadecanoate 19. 2-OH 16:0 Me. 2-hydroxyhexadecanoate 20. 18:2 ^{9,12} Me. cis-9,12-octadecadienoate 21. 18:1 ⁹ Me. cis-9-octadecenoate 22. 18:1 ⁹ Me. trans-9-octadecenoate 23. 18:0 Me. octadecanoate 24. 19:0 ^A Me. cis-9,10-methyleneoctadecanoate 25. 19:0 Me. nonadecanoate 26. 20:0 Me. eicosanoate			

Column: SPB-1, 30m x 0.25mm ID, 0.25µm film
 Cat. No.: 24028
 Oven: 150°C (4 min) to 250°C (5 min) at 4°C/min
 Carrier: helium, 20cm/sec (150°C)
 Det.: FID
 Inj.: 1µL

Water Soluble Fatty Acid (WSFA) Mixes

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
WSFA-2	0.1% of each fatty acid in water (weight percent) Acetic acid (C2) Butyric acid (C4) Valeric acid (C5) Propionic acid (C3) Isobutyric acid (iC4) Isovaleric acid (iC5)	5mL	47056	
WSFA-4	0.1% of each fatty acid in water (weight percent) Acetic acid (C2) Isobutyric acid (iC4) 2-Methylbutyric acid (2-methyl C4) Propionic acid (C3) Valeric acid (C5) Isovaleric acid (iC5) Butyric acid (C4)	5mL	47058	

Other Qualitative Standards

MIXES AND SOLUTIONS	COMPONENTS	QTY.	CAT. NO.	PRICE
Volatile Acid Standard Mix	10 components, 10mM each in deionized water Formic acid n-Valeric acid Acetic acid Isocaprylic acid (4-Methyl-n-valeric acid) Propionic acid Hexanoic acid (n-Caproic acid) Isobutyric acid Isovaleric acid Butyric acid Heptanoic acid	100mL	46975-U	
Nonvolatile Acid Standard Mix	8 components, 1 meq of each in deionized water Pyruvic acid Methyl malonic acid Lactic acid Malonic acid Oxalacetic acid Fumaric acid Oxalic acid Succinic acid	100mL	46985-U	

TABLE OF CONTENTS

TABLE OF CONTENTS	i
Technical Service	iii
Natural Products	iii
Storage	iii
Sphingolipid Structures and Pathways	iii
Package Weight	iii
Sphingoid bases, sphingolipids and glycosphingolipids.....	1
Sphingosines.....	2
Synthetic sphingosines with C18 sphingoid base.....	2
Synthetic sphingosines with sphingoid bases other than C18.....	2
Synthetic dihydroosphingosines	3
3-keto-Dihydroosphingosines	5
Phytosphingosines	5
Other sphingosine derivatives and precursors	6
Ceramides	7
Synthetic ceramides derived from C18-sphingosine	7
2-Hydroxy ceramides	10
Ceramide made from sphingosines with sphingoid bases other than C18	10
Dihydroceramides	11
2-Hydroxy dihydroceramides	11
Ceramides from natural sources.....	12
Phytoceramides	13
Fluorescent ceramides.....	14
Phosphosphingolipids	15
Sphingomyelins	15
Sphingosylphosphorylcholines (SPC).....	17
Sphingosine phosphates.....	18
Fluorescent sphingomyelins	18
Glycosphingolipids	19
Galactosylceramides and glucosylceramides.....	19
Lactosylceramides	23
Ceramide trihexosides	24
Globosides	25
Labeled glycolipids.....	25
Gangliosides	28
Glycosphingolipid reference mixes for TLC	31
Antibodies directed against glycolipids	32
Enzyme Inhibitors	34
Glycerolipids	40
Glycerophospholipids	40
Natural phospholipids	40
Synthetic phospholipids	42
Phosphatidylinositols	45
Bacterial tetraethers.....	47
Glycosyl glycerides.....	48
Fatty acids	48
Simple fatty acids	48
Saturated fatty acids and methyl esters.....	48
Unsaturated fatty acids and methyl esters.....	52
Trans fatty acids and methyl esters	57
Conjugated linoleic acid isomers (CLA)	59
Other CLA products and derivatives	61

Hydroxy fatty acids	61
2-Hydroxy fatty acids and methyl esters	61
3-Hydroxy fatty acids and methyl esters	64
Omega hydroxy fatty acids	66
Other hydroxy fatty acids	68
Branched and cyclic fatty acids	69
iso-Fatty acids and esters	69
anteiso-Fatty acids and esters	70
Methylated fatty acids	70
Cyclopropyl fatty acids and esters	71
Unusual fatty acids and derivatives	71
Other lipids.....	72
Tocopherols	72
Cholestane derivatives.....	73
Plant sterols and steryl glucosides	74
Propyleneglycol Monoesters.....	75
Standards and reference compounds	75
Food industry mixes.....	75
Each methyl ester mix is carefully prepared by weight	75
Polyunsaturated fatty acid methyl esters mixes	76
Carbohydrate mixes.....	76
Other fatty acid methyl ester mixes	77
AOCS animal and vegetable oil reference mixes (RM mixes).....	78
Custom mixes	80
GLC Standard mixes.....	81
Water soluble fatty acid mixes	82
Microbiology standard mixes	83
Biochemical research standard mixes	83
Glycosphingolipid mixtures for TLC	84
Biochemicals and reagents	85
Stable isotope labeled compounds	85
Fluorescent compounds.....	86
Sphingolipid Structures and Pathways Wall Chart.....	95
Cross reference for product numbers and catalog pages.....	96
Product name index	100
International Dealers & Representatives.....	107
Tables	
Table I. AOCS Oil Reference Mixes.....	78
Table II. Standards for GC analysis	81
Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC	90

All chemicals listed in this catalog are for investigational use only. They are not intended for human consumption or to be used in food or food additives. None are for general drug or medicinal use on humans. We believe the information in this catalog, offered in good faith, is accurate.

Limited Warranty: All Matreya Products, except those specifically exempted, are warranted (for 90 days) to be free of defects in materials and workmanship, if properly stored. Any replacements required as a result of such defects will be made without charge provided that such defective products are returned with a written explanation. Please request a Returned Goods Authorization before returning products under this warranty.

Technical Service

Our technical service department may be contacted by telephone at 800.342.3595, or by e-mail to techservice@matreya.com.

Note that there are excellent resources for general information on lipid nomenclature, distribution and biochemistry available on the internet. One such source is www.lipidlibrary.co.uk maintained by W.W. Christie and colleagues. Another source for detailed discussion on isolation and purification protocols is www.cyberlipid.org. This site also maintains discussion groups for lipid researchers.

Natural Products

Some of our glycolipids are extracted from natural sources. These products have a normal heterogeneity in their lipid components, particularly in the fatty acids. Variations include carbon chain length as well as the presence or absence of 2-hydroxy fatty acids. Products based on sphingosine may contain longer chain sphingoid bases as well as chains with multiple double bonds. This heterogeneity may result in additional spots showing on TLC plates or multiple peaks in LC analyses. We have listed the typical fatty acid compositions of our natural products in the appendix.

Storage

Catalog items in unopened containers are stable for at least one year when stored under the conditions indicated in the catalog listing. Items containing unsaturated fatty acids are subject to oxidation and should be stored in solution in organic solvents or under argon. Glycolipids and phospholipids should not be stored in aqueous solutions due to potential hydrolysis.

Sphingolipid Structures and Pathways

In a clear and straightforward manner, this wall chart indicates the structures and relationships between most commonly discussed sphingolipids. A one-page thumbnail version of the chart is shown on page 95. Full size copies (approximately 35 x 26 inches) are available on request to customer service.

Package Weight

Unless otherwise specified, the package will contain at least the indicated amount and usually slightly more. The user is cautioned to always measure the required amount from the container.


Sphingoid bases, sphingolipids and glycosphingolipids.

Sphingoid bases such as sphingosine are the characteristic structural unit of the sphingolipids. The bases are long chain aliphatic amines, containing two or three hydroxyl groups, and typically a *trans*-double bond at C4. In animal tissues most abundant base is sphingosine with a C18 aliphatic chain containing a double bond in position 4. The saturated analogue is dihydroosphingosine or sphinganine. In plants the common long chain base is the 4 hydroxy saturated base phytosphingosine.

Sphingolipids are widely distributed in animal tissues, particularly cell membranes. Sphingoid bases linked to fatty acids via an amide bond at C2 are ceramides and are present in trace amounts in most tissues. Glycosphingolipids (ceramides having various mono- and oligosaccharides on the OH group at C1) are neutral glycosphingolipids (i.e., cerebrosides and globosides). Those with sialic acid derivatized sugars are acidic glycolipids (i.e., gangliosides). They are amphiphilic and can be solubilized in buffers via sonication and micelle formation.


Gangliosides are present in substantial amounts in nerve cell membranes, and together with globosides are found in the membranes of white and red blood cells. These plus the glycosphingolipids of the lacto- and neolacto-series are involved in cell recognition (e.g. blood group determinants). Glycolipid expression on the surface of cells determines their antigenicity as well as their status, i.e. differentiated vs. undifferentiated (embryonic), normal vs. malignant, etc. (1). The ganglioside GM1 stimulates nerve growth (2,3) and has been reported to have a curative effect on experimental Parkinsonism (4). For an overview see (5). Gangliosides are also being investigated as potential anti-tumor vaccines (57). Glycosphingolipids are also essential for the correct functioning of cell surface receptors (6). Matreya is your best source for many sphingolipids. Most of Matreya's sphingosines and ceramides are fully synthetic and as such 98%+ pure. Others, particularly the glycosphingolipids are highly purified natural products (98%+), and can be used either as standards or biochemical reagents without further purification.

Through total synthesis, all four isomers of sphingosine are available as well as a number of sphingosines with other than 18 carbons and a number of ceramides (for details in using ceramides in cell culture see Hauser et al. [9]). Fluorescent labeled ceramides, glycosphingolipids and sphingomyelins are also available for study. D. N. Brindley and his group have been exploring the interaction of ceramides, sphingosine and sphingosine 1-phosphate in regulating DNA synthesis and phospholipase D activity (10). **See Literature References on page 96.**



Sphingosines

Synthetic sphingosines with C18 sphingoid base




1802	D-erythro-Sphingosine Sphingosine, C18 chain C ₁₈ H ₃₇ NO ₂ CAS#: 123-78-4	25 mg
Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: white solid Solubility: alcohols, chloroform, DMSO Storage: -20°C		
Selective inhibitor of phosphokinase C		
References: Y. Hannun et al., <i>Science</i> , 235 , 670, 1987 S. Spiegel et al., <i>Proc. Intern. Conf. Biol. Function Glycosphingolipids</i> , Santa Barbara, CA 1990		
1806	L-threo-Sphingosine L-threo-Sphingosine, C18 chain C ₁₈ H ₃₇ NO ₂	10 mg
Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C		
1826	L-erythro-Sphingosine L-erythro-Sphingosine, C18 chain C ₁₈ H ₃₇ NO ₂ CAS#: 6036-75-5	5 mg
Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C		
1827	D-threo-Sphingosine D-threo-Sphingosine, C18 chain C ₁₈ H ₃₇ NO ₂ CAS#: 6036-85-7	5 mg
Source: synthetic Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C		
1304	Sphingosine D-erythro-Sphingosine C ₁₈ H ₃₇ NO ₂ CAS#: 123-78-4	10 mg
Source: semi-synthetic, bovine Mol. Wt.: 299 Purity: 98+% by TLC, GC Appearance: white solid Solubility: alcohol, chloroform Storage: -20°C		

Synthetic sphingosines with sphingoid bases other than C18

Varying chain lengths allow the study of translocation effects of sphingosines and ceramides into cells.


1833	D-erythro-C14-Sphingosine Sphingosine with C14 chain C ₁₄ H ₂₉ NO ₂	5 mg
Source: synthetic Mol. Wt.: 243 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C		

1835	D-erythro-C16-Sphingosine Sphingosine with C16 chain C ₁₆ H ₃₃ NO ₂	5 mg
	Source: synthetic Mol. Wt.: 271 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1837	D-erythro-C10-Sphingosine Sphingosine with C10 chain C ₁₀ H ₂₁ NO ₂	5 mg/ml, 1 ml
	Source: synthetic Mol. Wt.: 187 Purity: 98+% by TLC, GC Appearance: liquid Solvent: ethanol Solubility: ethanol Storage: -20°C	
1838	D-erythro-C12-Sphingosine Sphingosine with C12 chain C ₁₂ H ₂₅ NO ₂ CAS#: 6918-49-6	5 mg
	Source: synthetic Mol. Wt.: 215 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1840	D-erythro-C20-Sphingosine Sphingosine with C20 chain C ₂₀ H ₄₁ NO ₂	5 mg
	Source: synthetic Mol. Wt.: 328 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	



Synthetic dihydrosphingosines

D,L-*threo*- Dihydrosphingosine has also been found to be a significant inhibitor of sphingosine kinase (11). The D,L-*erythro*-isomer has been used as an inactive control. We now offer all four isomers in pure form making detailed studies possible. Safingol, the L-*threo*-isomer is a potent inhibitor of PKC and as such is capable of reversing multi-drug resistance in cancer cells (12). See Literature References on page 96.



1807	L-threo-Dihydrosphingosine (Safingol)	5 mg
1807-025	L- <i>threo</i> -Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 15639-50-6	25 mg

Source: synthetic **Mol. Wt.:** 301 **Melting Point (°C):** 103-114 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

References:


- C.W. Sachs et al., ibid., **270**, 26639, 1995
G.K. Schwartz et al., J. Natl. Cancer Inst., **87**, 1394, 1995

1831	D-erythro-Dihydrosphingosine	25 mg
1831-1	D-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 764-22-7	1 g
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C Inhibitor of PLA ₂ and PLD	
	References: D.I. Yule et al., ibid, 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, Adv. in Lipid Res., 26 , 59, 1993 C. W. Sachs et al., J. Biol. Chem., 270 , 26639, 1995	
1846	L-erythro-Dihydrosphingosine	1 mg
	L-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂	
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., ibid, 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, Adv. in Lipid Res., 26 , 59, 1993 C. W. Sachs et al., J. Biol. Chem., 270 , 26639, 1995	
1851	D-threo-Dihydrosphingosine	1 mg
	D-threo-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 6036-86-8	
	Source: synthetic Mol. Wt.: 301 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., ibid, 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, Adv. in Lipid Res., 26 , 59, 1993 C. W. Sachs et al., J. Biol. Chem., 270 , 26639, 1995	
1324	D,L-erythro-Dihydrosphingosine	25 mg
	D,L-erythro-Sphinganine, C18 chain C ₁₈ H ₃₉ NO ₂ CAS#: 3102-56-5	
	Source: synthetic Mol. Wt.: 301 Purity: erythro 77%; threo 23% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C Inhibitor of sphingosine kinase	
	References: D.I. Yule et al., ibid, 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, Adv. in Lipid Res., 26 , 59, 1993 C. W. Sachs et al., J. Biol. Chem., 270 , 26639, 1995	
1326	D,L-C16-Dihydrosphingosine (mixed isomers)	10 mg
	D,L-Sphinganine with C16 chain C ₁₆ H ₃₅ NO ₂	
	Source: synthetic Mol. Wt.: 273 Purity: erythro 90%, threo 10% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
	References: D.I. Yule et al., ibid, 268 , 12353, 1993 B. M. Buehrer, R.M. Bell, Adv. in Lipid Res., 26 , 59, 1993 C. W. Sachs et al., J. Biol. Chem., 270 , 26639, 1995	
1845	D-erythro-C20-Dihydrosphingosine	5 mg
	D-erythro-Sphinganine, C20 chain C ₂₀ H ₄₃ NO ₂ CAS#: 24006-62-0	
	Source: synthetic Mol. Wt.: 330 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1, warm ethanol Storage: -20°C	

1839	D,L-erythro-C20-Dihydrosphingosine D,L-erythro-Sphinganine, C20 chain	C ₂₀ H ₄₃ NO ₂	10 mg
------	---	---	--------------

Source: synthetic **Mol. Wt.:** 330 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol 5:1, warm ethanol **Storage:** -20°C

3-keto-Dihydrosphingosines



Catalog number 1876

1876	3-keto-Dihydrosphingosine•HCl 3-keto-Sphinganine hydrochloride	C ₁₈ H ₃₇ NO ₂ •HCl	CAS#: 18944-28-0 10 mg
------	--	--	---

Source: synthetic **Mol. Wt.:** 336 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1891	3-keto-C6-Dihydrosphingosine•HCl 1-Hydroxy-2-amino-3-keto-hexane • HCl	C ₆ H ₁₃ NO ₂ • HCl	10 mg
------	--	--	--------------

Source: synthetic **Mol. Wt.:** 168 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** ethanol, methanol, DI water **Storage:** -20°C


1892	3-keto-C8-Dihydrosphingosine•HCl 1-Hydroxy-2-amino-3-keto-octane • HCl	C ₈ H ₁₇ NO ₂ • HCl	10 mg
------	--	--	--------------

Source: synthetic **Mol. Wt.:** 196 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, DI water **Storage:** -20°C

1893	3-keto-C12-Dihydrosphingosine•HCl 1-Hydroxy-2-amino-3-keto-dodecane • HCl	C ₁₂ H ₂₅ NO ₂ • HCl	10 mg
------	---	---	--------------

Source: synthetic **Mol. Wt.:** 252 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Phytosphingosines




Catalog number 1330

1330	Phytosphingosine 4-Hydroxysphinganine	C ₁₈ H ₃₉ NO ₃	CAS#: 554-62-1 50 mg 1 g
------	---	---	---

Source: natural, yeast (*Pichia ciferri*) **Mol. Wt.:** 318 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** ethanol, methanol, chloroform/methanol 2:1 (warm) **Storage:** -20°C

Reference:
Keliu, Xiping Zhang, Robert L. Lester, and Robert C. Dickson. J. Biol. Chem., **280**, Issue 24, 22679-22687, June 17, 2005

Other sphingosine derivatives and precursors



Catalog number 1896

1320	N,N-Dimethyl-D-erythro-sphingosine C ₂₀ H ₄₁ NO ₂ CAS#: 119567-63-4	5 mg/ml, 1 ml
------	--	----------------------

Source: synthetic **Mol. Wt.:** 328 **Purity:** 98+% by TLC **Appearance:** liquid
Solvent: isopropanol **Solubility:** chloroform, ethanol, isopropanol, methanol
Storage: -20°C

Inhibitor of phosphokinase C

Reference:
B. Felding-Habermann et al., Biochemistry, **29**, 6314, 1990

1896	N,N-Dihexyl-D-erythro-sphingosine Sphingosine with tertiary amine group C ₃₀ H ₆₁ NO ₂	5 mg/ml, 1 ml
------	---	----------------------

Source: synthetic **Mol. Wt.:** 468 **Purity:** 95% by TLC **Appearance:** liquid
Solvent: ethanol **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

1805	N-Palmitoyl serinol C ₁₉ H ₃₉ NO ₃ CAS#: 126127-31-9	10 mg
------	---	--------------

Source: synthetic **Mol. Wt.:** 329 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, methanol, ethanol **Storage:** -20°C

Sphingosine precursor

Ceramides

Ceramide is a fatty acid amide of sphingosine. It may be formed by dehydrogenation of dihydroceramide; by hydrolysis of sphingomyelin or glycosphingolipids; or by acylation of free sphingosine. Ceramide functions as a precursor in the synthesis of sphingomyelin (by an exchange reaction with phosphatidylcholine and phosphatidylethanolamine); of glycosphingolipids (by glycosylation with UDP-hexose); and of free sphingosine and fatty acid by hydrolysis. The sphingosine can be phosphorylated by a kinase to form sphingosine-1-phosphate, which may undergo further hydrolysis or cleavage.

Control of sphingolipid metabolism maintains vital balance points in cell physiology. Two of ceramide's metabolites, sphingosine-1-phosphate and glucosylceramide, produce cell proliferation. Sphingosine-1-phosphate is also a highly active regulator of angiogenesis, vascular maturation, cardiac development, immunity, and directed cell movement. Sphingosine, the free base, is a potent inhibitor of protein kinase C and is involved in intracellular calcium regulation.

Sphingolipid enzymes seem to be particularly active in cancers, so modifying their activities by exogenous action may provide alternatives to chemical therapies. These enzymes are controlled by many known agents, such as 1,25-dihydroxy-vitamin D₃, tumor necrosis factor- α , nerve growth factor, interleukin 1, endothelial growth factor, glutathione, arachidonic acid, dexamethasone, many anticancer drugs, therapeutic radiation, and activators of the FAS receptor.


Ceramide exerts numerous biological effects, including induction of cell maturation, cell cycle arrest, terminal cell differentiation, cell senescence, and cell death. Other effects include producing reactive oxygen in mitochondria (followed by apoptosis) and stimulating phosphorylation of certain proteins (especially mitogen activated protein). It also stimulates some protein phosphatases (especially protein phosphatase 2A). Thus ceramide is an important controller of protein activity.

It is apparent from these relationships that ceramide exists at the crux of several enzyme reaction cycles and that experiments involving sphingolipid function call for control of all of the cycles and their branch-off points. Matreya is the major supplier of these lipids, which can be used as standards for analysis of tissues (a much needed part of modern research) and identification of major sphingolipids.

Ceramides with short side chains have been shown to enter easily into cells where they are biologically active. Ceramides with longer side chains, however, also enter cells if dissolved in dodecane-isopropanol first. Fluorescent labeled ceramides and sphingomyelin made from fluorescent labeled acids instead of plain fatty acids are also available for the study of the localization and metabolism of sphingolipids in the cell. Matreya now offers all four isomers of C2, C4, C6 and C18 ceramides. The corresponding dihydroceramides are being used as inactive controls.

In three major reviews, Radin (54-56) has discussed the biochemistry and chemistry of ceramide and outlined many potential approaches to cancer therapy using ceramides and related compounds as generators of apoptosis.

See Literature References on page 96.



Catalog number 1901

Synthetic ceramides derived from C18-sphingosine

1901	N-Acetyl-D-erythro-sphingosine N-C2:0-D-erythro-Ceramide C ₂₀ H ₃₉ NO ₃ CAS#: 3102-57-6	10 mg
------	---	--------------

Source: synthetic **Mol. Wt.:** 342 **Purity:** 98+ by TLC, GC **Appearance:** white solid
Solubility: chloroform, ethanol, methanol, DMSO, (up to 5 mg/ml) **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1829	N-Acetyl-L-threo-sphingosine N-C2:0-L-threo-Ceramide C ₂₀ H ₃₉ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1847	N-Acetyl-L-erythro-sphingosine N-C2:0-L-erythro-Ceramide C ₂₀ H ₃₉ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1900	N-Hexanoyl-D-erythro-sphingosine N-C6:0-D-erythro-Ceramide C ₂₄ H ₄₇ NO ₃ CAS#: 124753-97-5	10 mg
	Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1828	N-Hexanoyl-L-threo-sphingosine N-C6:0-L-threo-Ceramide C ₂₄ H ₄₇ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1848	N-Hexanoyl-L-erythro-sphingosine N-C6:0-L-erythro-Ceramide C ₂₄ H ₄₇ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC; GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1809	N-Hexanoyl-D-threo-sphingosine N-C6:0-D-threo-Ceramide C ₂₄ H ₄₇ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 398 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol, DMSO (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1903	N-Octanoyl-D-erythro-sphingosine N-C8:0-D-erythro-Ceramide C ₂₆ H ₅₁ NO ₃ CAS#: 74713-59-0	10 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+ by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO (up to 5 mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	

1830	N-Octanoyl-L-threo-sphingosine N-C8:0-L-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1810	N-Octanoyl-D-threo-sphingosine N-C8:0-D-threo-Ceramide C ₂₆ H ₅₁ NO ₃	1 mg
	Source: synthetic Mol. Wt.: 426 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1333	N-Decanoyl-D-erythro-sphingosine N-C10:0-D-erythro-Ceramide C ₂₈ H ₅₅ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 454 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO, (up to 5mg/ml) Storage: -20°C	
2037	N-Pentadecanoyl-D-erythro-sphingosine N-C15:0-D-erythro-Ceramide C ₃₃ H ₆₅ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 524 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
1915	N-Hexadecanoyl-D-erythro-sphingosine N-C16:0-D-erythro-Ceramide C ₃₄ H ₆₇ NO ₃ CAS#: 24696-26-2	10 mg
	Source: synthetic Mol. Wt.: 538 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
	Reference: Yosuke Osawa, Hiroshi Uchinami, Jacek Bielawski, Robert F. Schwabe, Yusuf A. Hannun, and David A. Brenner. J. Biol. Chem., 280 , Issue 30, 27879-27887, July 29, 2005	
2038	N-Heptadecanoyl-D-erythro-sphingosine N-C17:0-D-erythro-Ceramide C ₃₅ H ₆₉ NO ₃ CAS#: 24696-26-2	10 mg
	Source: synthetic Mol. Wt.: 552 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	
1832	N-Octadecanoyl-D-erythro-sphingosine N-C18:0-D-erythro-Ceramide C ₃₆ H ₇₁ NO ₃ CAS#: 2304-81-6	10 mg
	Source: synthetic Mol. Wt.: 566 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, hot ethanol, chloroform/methanol 2:1 (up to 5mg/ml) Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
2039	N-Nonadecanoyl-D-erythro-sphingosine N-C19:0-D-erythro-Ceramide C ₃₇ H ₇₃ NO ₃	10 mg
	Source: synthetic Mol. Wt.: 580 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, warm methanol Storage: -20°C	

1843	N-Octadecanoyl-L-threo-sphingosine N-C18:0-L-threo-Ceramide C ₃₆ H ₇₁ NO ₃	1 mg
Source: synthetic Mol. Wt.: 566 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C		
Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994		
1850	N-Octadecanoyl-L-erythro-sphingosine N-C18:0-L-erythro-Ceramide C ₃₆ H ₇₁ NO ₃	1 mg
Source: synthetic Mol. Wt.: 566 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up tp 5mg/ml) Storage: -20°C		
Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994		
1855	N-Octadecanoyl-D-threo-sphingosine N-C18:0-D-threo-Ceramide C ₃₆ H ₇₁ NO ₃	1 mg
Source: synthetic Mol. Wt.: 566 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5mg/ml) Storage: -20°C		
1916	N-Tetracosanoyl-D-erythro-sphingosine N-C24:0-D-erythro-Ceramide C ₄₂ H ₈₃ NO ₃ CAS#: 34435-05-7	5 mg
Source: synthetic Mol. Wt.: 650 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform Storage: -20°C		


2-Hydroxy ceramides

2042	N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-sphingosine N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-ceramide C ₃₀ H ₅₉ NO ₄	5 mg
Source: synthetic Mol. Wt.: 498 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol, ethanol, DMSO Storage: -20°C		
2044	N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-sphingosine N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-ceramide C ₃₆ H ₇₁ NO ₄	5 mg
Source: synthetic Mol. Wt.: 582 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol/water, 2:1:0.5 Storage: -20°C		

Ceramide made from sphingosines with sphingoid bases other than C18

1842	N-Acetyl-D-erythro-sphingosine (C14 sphingoid base) N-C2:0 Ceramide of D-erythro-C14-sphingosine C ₁₆ H ₃₁ NO ₂	5 mg
Source: synthetic Mol. Wt.: 285 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, DMSO, DMF (up to 5 mg/ml) Storage: -20°C		
Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994		
1856	N-Hexanoyl-D-erythro-sphingosine (C8 sphingoid base) N-C6:0 Ceramide of D-erythro-C8-sphingosine C ₁₄ H ₂₇ NO ₂	1 mg
Source: synthetic Mol. Wt.: 257 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		

Dihydroceramides



1834	N-Acetyl-D-erythro-dihydrosphingosine N-C2:0-D-erythro-Dihydroceramide; N-Acetyl-D-erythro-sphinganine C ₂₀ H ₄₁ NO ₃	1 mg
------	---	------

Source: synthetic **Mol. Wt.:** 344 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1910	N-Hexanoyl-D-erythro-dihydrosphingosine N-C6:0-D-erythro-Dihydroceramide; N-Hexanoyl-D-erythro-sphinganine C ₂₄ H ₄₉ NO ₃	1 mg
------	---	------

Source: synthetic **Mol. Wt.:** 400 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1854	N-Octanoyl-D-erythro-dihydrosphingosine N-C8:0-D-erythro-Dihydroceramide; N-Octanoyl-D-erythro-sphinganine C ₂₆ H ₅₃ NO ₃	1 mg
------	---	------

Source: synthetic **Mol. Wt.:** 428 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, DMSO **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

2041	N-Octadecanoyl-D-erythro-dihydrosphingosine N-C18:0-D-erythro-Dihydroceramide; N-Octadecanoyl-D-erythro-sphinganine C ₃₆ H ₇₃ NO ₃	10 mg
------	--	-------

Source: synthetic **Mol. Wt.:** 568 **Purity:** 98% by TLC **Appearance:** white solid **Solubility:** warm chloroform/methanol, 5:1; hot ethanol, DMSO **Storage:** -20°C

2-Hydroxy dihydroceramides

2043	N-(R,S)-alpha-Hydroxydodecanoyl-D-erythro-dihydrosphingosine N-(R,S)-alpha-Hydroxy-C12:0-D-erythro-dihydroceramide C ₃₀ H ₆₁ NO ₄	5 mg
------	--	------


Source: synthetic **Mol. Wt.:** 500 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol/water, 2:1:0.5 **Storage:** -20°C

2045	N-(R,S)-alpha-Hydroxyoctadecanoyl-D-erythro-dihydrosphingosine N-(R,S)-alpha-Hydroxy-C18:0-D-erythro-dihydroceramide C ₃₆ H ₇₃ NO ₄	5 mg
------	--	------

Source: synthetic **Mol. Wt.:** 584 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform/methanol/water, 2:1:0.5 **Storage:** -20°C

2047	N-(R,S)-alpha-Hydroxyhexadecanoyl-D-erythro-dihydrosphingosine N-(R,S)-alpha-Hydroxy-C16:0-D-erythro-dihydroceramide C ₃₄ H ₆₉ NO ₄	5 mg
	Source: synthetic Mol. Wt.: 556 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol/water, 2:1:0.5 Storage: -20°C	

Ceramides from natural sources




General ceramide structure

1056	Ceramides Ceramides with hydroxy and non-hydroxy acyl groups C ₄₂ H ₈₃ NO ₄ CAS#: 104404-17-13	25 mg
	Source: natural, bovine Mol. Wt.: 666(2-hydroxy-lignoceroyl) Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1322	Ceramides	10 mg
1322-05	Ceramides with mostly non-hydroxy acyl groups C ₃₆ H ₇₁ NO ₃	50 mg
	Source: natural, bovine Mol. Wt.: 566 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, ethanol Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	
1323	Ceramides	10 mg
1323-05	Ceramides with mostly hydroxy acyl groups C ₃₆ H ₇₁ NO ₄	50 mg
	Source: natural, bovine Mol. Wt.: 582 (2-hydroxy-stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
	Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994	

See Table III in Appendix for typical fatty acid content of products prepared from natural sources.


Phytoceramides



Catalog number 1897

1897	N-Acetyl-phytosphingosine N-C2:0-Phytoceramide C ₂₀ H ₄₁ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 360 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm), warm DMSO Storage: -20°C		
1895	N-Hexanoyl-phytosphingosine N-C6:0-Phytoceramide C ₂₄ H ₄₉ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 416 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm) Storage: -20°C		
1894	N-Octanoyl-phytosphingosine N-C8:0-Phytoceramide C ₂₆ H ₅₃ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 444 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol, chloroform/methanol 1:1 (warm) Storage: -20°C		
2035	N-Hexadecanoyl-phytosphingosine N-C16:0-Phytoceramide C ₃₄ H ₆₉ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 556 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		
2034	N-Stearoyl-phytosphingosine N-C18:0-Phytoceramide C ₃₆ H ₇₃ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 584 Purity: 98+% by TLC- MS Appearance: white solid Solubility: chloroform/methanol 1:1 (warm) Storage: -20°C		
2036	N-Tetracosanoyl-phytosphingosine N-C24:0-Phytoceramide C ₄₂ H ₈₅ NO ₄	5 mg
Source: semi-synthetic, yeast (Pichia ciferri) Mol. Wt.: 668 Purity: 98+% by TLC- MS Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		

Fluorescent ceramides



1841	N-Hexanoyl-NBD-D-erythro-sphingosine	100 µg
1841-001	N-C6:0-NBD-ceramide; N-C6:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-sphingosine C ₃₀ H ₄₉ N ₅ O ₆ CAS#: 86701-10-2	1 mg
Source: synthetic Mol. Wt.: 575 Melting Point (°C): 85.7-87.9 Purity: 98+% by TLC Appearance: orange film, vacuum dried Solubility: chloroform, ethanol, methanol Storage: -20°C		
Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994		
1618	N-Dodecanoyl-NBD-D-erythro-sphingosine	100 µg
1618-001	N-C12:0-NBD ceramide; N-C12:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-sphingosine C ₃₆ H ₆₁ N ₅ O ₆	1 mg
Source: synthetic Mol. Wt.: 660 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C		
1857	N-Hexanoyl-NBD-L-threo-sphingosine	100 µg
1857-001	N-C6:0-NBD-ceramide; N-C6:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminocaproyl)-L-threo-sphingosine C ₃₀ H ₄₉ N ₅ O ₆	1 mg
Source: synthetic Mol. Wt.: 575 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
Reference: J. M. L. Hauser et al., J. Biol. Chem. 269 , 6803, 1994		
1620	N-Dodecanoyl-NBD-L-threo-sphingosine	100 µg
1620-001	N-C12:0-NBD-ceramide; N-C12:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-sphingosine C ₃₆ H ₆₁ N ₅ O ₆	1 mg
Source: synthetic Mol. Wt.: 660 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C		
1624	N-Hexanoyl-NBD-L-threo-dihydroceramide	100 µg
1624-001	N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-L-threo-dihydroceramide, fluorescent; N-(NBD-aminocaproyl)-L-threo-dihydroceramide C ₃₀ H ₅₁ N ₅ O ₆	1 mg
Source: synthetic Mol. Wt.: 578 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C		
1623	N-Dodecanoyl-NBD-L-threo-dihydroceramide	100 µg
1623-001	N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-L-threo-dihydroceramide, fluorescent; N-(NBD-aminolauroyl)-L-threo-dihydroceramide C ₃₆ H ₆₃ N ₅ O ₆	1 mg
Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C		


1626	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine	100 µg
1626-001	N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-dihydrosphingosine C ₃₀ H ₅₁ N ₅ O ₆	1 mg
	Source: synthetic Mol. Wt.: 578 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1625	N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine	100 µg
1625-001	N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-dihydrosphingosine C ₃₆ H ₆₃ N ₅ O ₆	1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1628	N-Hexanoyl-NBD-phytosphingosine	100 µg
1628-001	N-C6:0-NBD-phytoceramide; N-C6:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminocaproyl)-phytosphingosine C ₃₀ H ₅₁ N ₅ O ₇	1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 594 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1627	N-Dodecanoyl-NBD-phytosphingosine	100 µg
1627-001	N-C12:0-NBD-phytoceramide; N-C12:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminolauroyl)-phytosphingosine C ₃₆ H ₆₃ N ₅ O ₇	1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 678 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information.

Phosphosphingolipids

Sphingomyelins




Catalog number 1051

1051	Sphingomyelin	25 mg
1051-1	SPM; ceramide-1-phosphorylcholine C ₄₁ H ₈₃ N ₂ O ₆ P CAS#: 85187-10-6	1 g

Source: natural, bovine **Mol. Wt.:** 731 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Predominately C18:0 and C24:1 fatty acids


1328	Sphingomyelin SPM; ceramide-1-phosphorylcholine C ₄₇ H ₉₅ N ₂ O ₆ P CAS#: 85187-10-6	25 mg
	Source: natural, porcine Mol. Wt.: 815 (lignoceroyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol Storage: -20°C	
	Predominately C16:0 and C24:0 fatty acids	
1329	Sphingomyelin SPM; ceramide-1-phosphorylcholine C ₄₆ H ₉₃ N ₂ O ₆ P CAS#: 85187-10-6	25 mg
	Source: natural, bovine buttermilk Mol. Wt.: 801 (tricosanoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform, ethanol Storage: -20°C	
	Approximately equal amounts of C16:0, C22:0, C23:0, and C24:0 fatty acids	
1332 1332-1	Sphingomyelin Ceramide-1-phosphorylcholine C ₃₉ H ₇₉ N ₂ O ₆ P	25 mg 1 gram
	Source: natural, egg yolk, chicken Mol. Wt.: 703 (palmitate) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, methanol, warm ethanol Storage: -20°C	
1907	N-Acetyl-sphingosylphosphorylcholine Sphingomyelin with C2:0 fatty acid C ₂₅ H ₅₁ N ₂ O ₆ P	5 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 506 Purity: 98+% by TLC Appearance: vacuum dried Solubility: ethanol, chloroform/methanol 2:1 Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1909	N-Hexanoyl-sphingosylphosphorylcholine Sphingomyelin with C6:0 fatty acid C ₂₉ H ₅₉ N ₂ O ₆ P	5 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 563 Purity: 98+% by TLC Appearance: solid, vacuum dried Solubility: ethanol, chloroform/methanol 2:1 Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1911	N-Octadecanoyl-sphingosylphosphorylcholine Sphingomyelin with C18:0 fatty acid C ₄₁ H ₈₃ N ₂ O ₆ P	5 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 731 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1890	N-Heptadecanoyl-sphingosylphosphorylcholine Sphingomyelin with C17:0 fatty acid C ₄₀ H ₈₁ N ₂ O ₆ P	5 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 717 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1917	N-Eicosanoyl-D-erythro-sphingosylphosphorylcholine Sphingomyelin with C20:0 fatty acid C ₄₃ H ₈₇ N ₂ O ₆ P	0.5 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 759 Purity: 98+% by TLC Appearance: solid, vacuum dried Solubility: chloroform/methanol 14:1, ethanol, methanol Storage: -20°C	



Catalog number 1918

1918	N-Docosanoyl-D-erythro-sphingosylphosphorylcholine Sphingomyelin with C22:0 fatty acid	0.5 mg $C_{45}H_{91}N_2O_6P$
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 787 Purity: 98+% by TLC Appearance: solid, vacuum dried Solubility: chloroform/methanol 14:1, ethanol, methanol Storage: -20°C	
2200	N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine D-erythro-Sphingomyelin with 1- ¹³ C-palmitic acid; SPM with ¹³ C labeled fatty acid	1mg $^{12}C_{38}^{13}CH_{79}N_2O_6P$
	Source: semi-synthetic, bovine Mol. Wt.: 703 Purity: 98+% by TLC Appearance: waxy solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1327	N-Acetyl-sphingosylphosphorylethanolamine Sphingosylphosphorylethanolamine with C2:0 fatty acid side chain (D-erythro)	5 mg $C_{38}H_{77}N_2O_6P$
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 689 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

Sphingosylphosphorylcholines (SPC)



Catalog number 1318


1318	D-erythro-Sphingosylphosphorylcholine D-erythro-SPC	5 mg $C_{23}H_{49}N_2O_5P$
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 464 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1319	L-threo-Sphingosylphosphorylcholine L-threo-SPC	5 mg $C_{23}H_{49}N_2O_5P$
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 464 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1321 1321-05	Sphingosylphosphorylcholine lyso-Sphingomyelin; SPC (mixture of D-erythro and L-threo isomers) $C_{23}H_{49}N_2O_5P$ CAS#: 82970-80-7	10 mg 50 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 466 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

1913	lyso-Dihydrosphingomyelin Dihydrosphingosylphosphorylcholine (mixture of D-erythro and L-threo isomers) C ₂₃ H ₅₂ N ₂ O ₅ P	1 mg
Source: semi-synthetic, bovine buttermilk Mol. Wt.: 485 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C		

Sphingosine phosphates

1803	D-erythro-Sphingosine-1-phosphate S-1-P, S-P-A C ₁₈ H ₃₈ NO ₅ P CAS#: 26993-30-6	5 mg
Source: synthetic Mol. Wt.: 380 Purity: 98+% by TLC Appearance: white solid Solubility: warm acetic acid Storage: -20°C		
References: T.K. Ghosh et al., J. Biol. Chem., 269 , 22628, 1994 L. Kindman et al., ibid, 269 , 13088, 1994 A. Olivera et al., ibid, 269 , 17924, 1994 M. Mattie, et. al., J. Biol. Chem. 269 :3181, 1994 Yosuke Osawa, Hiroshi Uchinami, Jacek Bielawski, Robert F. Schwabe, Yusuf A. Hannun, and David A. Brenner. J. Biol. Chem., 280 , Issue 30, 27879-27887, July 29, 2005		
1852	D-erythro-Dihydrosphingosine-1-phosphate C ₁₈ H ₄₀ NO ₅ P CAS#: 19794-97-9	5 mg
Source: synthetic Mol. Wt.: 382 Purity: 98+% by TLC Appearance: white solid Solubility: warm acetic acid Storage: -20°C		
2046	N-Hexadecanoyl-D-erythro-sphingosine-1-phosphate, NH₄⁺ salt N-C16:0-Ceramide-1-phosphate C ₃₄ H ₆₈ NO ₆ P	5 mg
Source: synthetic Mol. Wt.: 618 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/acetic acid, 60:15:25 Storage: -20°C		

Fluorescent sphingomyelins



1912	N-Hexanoyl-NBD-sphingosylphosphorylcholine N-C6:0-NBD-sphingomyelin, fluorescent; N-C6:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminocaproyl)-sphingomyelin C ₃₅ H ₆₁ N ₆ O ₉ P CAS#: 94885-04-8	100 µg 1 mg
Source: semi-synthetic, bovine buttermilk Mol. Wt.: 740 Purity: 98+% by TLC Appearance: red-brown solid Solubility: chloroform, ethanol, methanol Storage: -20°C		

Mixture of D-erythro and L-threo isomers

1619 1619-001	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	100 µg 1 mg
------------------	---	------------------------

N-C12:0-NBD-sphingomyelin, fluorescent; N-C12:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminolauroyl)-sphingomyelin C₄₁H₇₃N₆O₄P

Source: semi-synthetic, bovine buttermilk **Mol. Wt.:** 825 **Purity:** 98+% by TLC

Appearance: orange solid **Solubility:** chloroform/methanol 2:1, methanol

Storage: -20°C

Mixture of D-erythro and L-threo isomers

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical Service for more information.


Glycosphingolipids

Glycosphingolipids are widely distributed in animal and plant tissues. They consist of a ceramide (Cer) bound in glycosidic linkage through the primary hydroxyl to a mono- or oligosaccharide which may contain substituents such as a sulfate, acetate, or phosphate group. They are amphiphilic and the less glycosylated compounds can be dispersed in buffers by dissolving them in a detergent or organic solvent (EtOH, DMSO, isoPrOH) and mixing by sonication.


Galactosphingolipids, mainly GalCer (cerebrosides) and its sulfate ester, occur in large amounts in the nervous system. Glucosphingolipids, the simplest of which is GlcCer (glucocerebrosides), are very widely distributed, particularly in nerve cell membranes. GlcCer is isolated from a variety of natural sources including human, bovine, and plant. Each of these sources has a heterogeneity in the fatty acid content of the ceramide as well as an occasional variation in the sphingoid chain. Globosides (containing both glucose and galactose) are a prominent group of glycosphingolipids, they contain an α -linked galactose moiety and are typically located in blood cell membranes. Gangliosides are another prominent group of glycosphingolipids; they are acidic because of substitution with sialic (neuraminic) acid. The glycosphingolipids function in a wide range of enzyme and structural interactions, such as immunological or membrane recognition phenomena, binding of microbial pathogens, hormone and growth factor actions, cancer cell growth and malignancy, atherosclerosis, genetic disease errors, blood group determinants, etc. Tissues change in glycosphingolipid composition during embryogenesis, maturation, aging, and other vital physiological processes. Some glycosphingolipids stimulate cell proliferation, others induce apoptosis, effects of great significance to cancer therapy and maturational development. Marked differences in glycosphingolipid composition are seen in normal and cancerous cells. See references 41-53.

See Literature References on page 96.


Galactosylceramides and glucosylceramides



Galactosylceramide



Glucosylceramide

1050	Cerebrosides Galactosylceramide, ceramide beta-D-galactoside $C_{48}H_{93}NO_8$ CAS#: 85305-88-0	50 mg
	Source: natural, bovine Mol. Wt.: 812 (lignoceryl form) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Contains both hydroxy and non-hydroxy fatty acid side chains	
1066	Cerebroside, Kerasin (top spot) Galactosylceramide with mostly non-hydroxy fatty acid side chain $C_{42}H_{81}NO_8$ CAS#: 536-13-0	10 mg
	Source: natural, bovine Mol. Wt.: 728 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.5 Storage: -20°C	
1138	Cerebroside, Phrenosin (bottom spot) Galactosylceramide with mostly 2-hydroxy fatty acid side chains $C_{42}H_{81}NO_9$ CAS#: 37211-11-3	10 mg
	Source: natural, bovine Mol. Wt.: 743 (2-hydroxystearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.5 Storage: -20°C	
1305	Psychosine, (in free amine form) lyso-Cerebroside; 1-beta-D-galactosylsphingosine $C_{24}H_{47}NO_7$ CAS#: 2238-90-6	10 mg
	Source: semi-synthetic, bovine Mol. Wt.: 461 Purity: 98+% by TLC Appearance: off-white solid Solubility: ethanol, chloroform/methanol 5:1 Storage: -20°C	
		
	Catalog number 1325	
1325	N-Acetyl-psychosine N-C2:0-Cerebroside; cerebroside with C2:0 fatty acid $C_{26}H_{49}NO_8$	10 mg
	Source: semi-synthetic, bovine Mol. Wt.: 503 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1335	N-Pentadecanoyl-psychosine N-C15:0-Cerebroside $C_{39}H_{75}NO_8$	5 mg
	Source: semi-synthetic, bovine Mol. Wt.: 685 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/ methanol, 2:1 Storage: -20°C	
1334 1334-50	N-Octanoyl-β-D-galactosylceramide N-C8:0-Galactosylceramide $C_{32}H_{61}NO_8$	10 mg 50 mg
	Source: semi-synthetic, bovine Mol. Wt.: 588 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/ methanol, 9:1, ethanol, methanol Storage: -20°C	

1621	N-Hexanoyl-NBD-galactosylceramide	100 µg
1621-001	N-C6:0-NBD-beta-D-galactosylsphingosine; N-C6:0-NBD-cerebroside; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-galactosylsphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg

Source: semi-synthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/ methanol, 5:1, methanol **Storage:** -20°C

1057	Glucocerebrosides	5 mg
	Glucosylceramide; ceramide beta-D-glucoside C ₄₆ H ₉₃ NO ₈ CAS#: 85305-87-9	


Source: natural, human **Mol. Wt.:** 812 (lignoceroyl) **Purity:** 98+% by TLC
Appearance: white to an off white solid **Solubility:** chloroform/methanol 2:1
Storage: -20°C

Contains 24:1 fatty acid side chain. See Table III (p 90-94) for other fatty acid content

1521	Glucocerebrosides	5 mg
1521-50	Glucosylceramide; ceramide beta-D-glucoside C ₄₆ H ₈₉ NO ₈	50 mg

Source: natural, bovine buttermilk **Mol. Wt.:** 784 (docosanoyl) **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

See Table III (p 90-94) for side chain variants



Catalog number 1522

1522	Glucocerebrosides, plant	5 mg
1522-100	Glucosylceramide; ceramide beta-D-glucoside C ₄₀ H ₇₅ NO ₉	100 mg

Source: natural, plant **Mol. Wt.:** 714 (2-hydroxyhexadecanoyl) **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C


Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4,8) and most of the fatty acids are of the 2-hydroxy type. See Table III page 90-94.

1622	N-Hexanoyl-NBD-glucosylceramide	100 µg
1622-001	N-C6:0-NBD-beta-D-glucosylsphingosine; N-C6:0-NBD-glucosylceramide, fluorescent C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg

Source: semi-synthetic, bovine **Mol. Wt.:** 738 **Purity:** 98+% by TLC **Appearance:** orange solid **Solubility:** chloroform/methanol 5:1, methanol **Storage:** -20°C

1306	Glucopsychosine	5 mg
	Glucosylsphingosine; lyso-glucocerebroside; 1-beta-D-glucosylsphingosine C ₂₄ H ₄₇ NO ₇ CAS#: 52050-17-6	

Source: semi-synthetic, bovine **Mol. Wt.:** 461 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** ethanol, methanol, chloroform/methanol 2:1 **Storage:** -20°C



Catalog number 1310

1310 Glucosylsphingosine 5 mg

Glucosylsphingosine; lyso-glucocerebroside; 1-beta-D-glucosylsphingosine
 $C_{24}H_{45}NO_7$ CAS#: 52050-17-6


Source: natural, plant **Mol. Wt.:** 459 **Purity:** 98+% by TLC
Appearance: off white solid **Solubility:** chloroform/methanol 4:1 **Storage:** -20°C

Sphingoid backbone is >95% 4,8-sphingadiene (C18:2 t,t-4, 8)

1531 N-Docosanoyl-glucosylsphingosine 1 mg

Glucocerebroside with C22:0 fatty acid side chain; N-Docosanoyl- β -glucosylsphingosine $C_{46}H_{89}NO_8$

Source: semi-synthetic, bovine **Mol. Wt.:** 784 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform **Storage:** -20°C



Catalog number 1049

1049 Sulfatides 50 mg

Ceramide-galactoside-3-sulfate; cerebroside sulfate $C_{42}H_{81}NO_{11}S$
 CAS#: 85496-63-5

Source: natural, bovine **Mol. Wt.:** 830 (stearoyl) Na Salt **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1 (if needed, a few drops of acetic acid) **Storage:** -20°C

1904 lyso-Sulfatide (NH_4^+ salt) 1 mg

Sphingosine-1-galactoside-3-sulfate $C_{24}H_{47}NO_{10}S$ CAS#: 38621-58-8

Source: semi-synthetic, bovine **Mol. Wt.:** 542 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

2076 N-Acetyl-sulfatide 1 mg

N-C2:0-sulfatide; N-acetyl-sphingosyl-beta-D-galactoside-3-sulfatide
 $C_{26}H_{49}NO_{11}S$

Source: semi-synthetic, bovine **Mol. Wt.:** 584 **Purity:** 98+% by TLC **Appearance:** solid **Solubility:** chloroform/methanol 1:1, ethanol, methanol **Storage:** -20°C


1875 N-Palmitoyl-sulfatide 1 mg

Sulfatide with C16:0 fatty acid side chain; N-palmitoyl-sphingosyl-beta-D-galactoside-3-sulfate $C_{40}H_{76}NO_{11}S$

Source: semi-synthetic, bovine **Mol. Wt.:** 780 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

1888	N-Tetracosanoyl-sulfatide N-C24:0-Sulfatide; N-tetracosanoyl-sphingosyl-beta-D-galactoside-3-sulfate $C_{48}H_{93}NO_{11}S$	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 892 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide N-C18:0-D ₃ -Sulfatide $C_{42}H_{78}D_3NO_{11}S$	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 833 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/DI water 2:1:0.1 Storage: -20°C	
1632	N-Dodecanoyl-NBD-sulfatide	100 µg
1632-001	N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate $C_{42}H_{71}N_5O_{14}S$	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 901 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	


Lactosylceramides



1500	Lactosylceramide LC, lactocerebrosides; CDH, ceramide beta-lactoside $C_{48}H_{91}NO_{13}$ CAS#: 4682-48-8	1 mg
	Source: natural, porcine Mol. Wt.: 890 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1, DMSO Storage: -20°C	
	Contains 2-hydroxy fatty acids (See Table I)	
1507	Lactosylceramide	5 mg
1507-50	LC; lactocerebrosides; CDH, ceramide beta-lactoside $C_{53}H_{101}NO_{13}$ CAS#: 4682-48-8	50 mg
	Source: natural, bovine buttermilk Mol. Wt.: 960 (tricosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
1517	lyso-Lactosylceramide Lactosylsphingosine; lyso-LC $C_{30}H_{57}NO_{12}$	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 623 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
1532	N-Palmitoyl-lactosylceramide Lactosylceramide with C16:0 fatty acid side chain $C_{46}H_{87}NO_{13}$	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 862 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	

1629	N-Hexanoyl-NBD-lactosylceramide	50 µg
1629-001	N-Hexanoyl-NBD-beta-D-lactosylphingosine; N-C6:0-NBD-beta-D-lactosylphingosine; N-C6:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminocaproyl)-lactosylphingosine C ₄₂ H ₆₉ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1630	N-Dodecanoyl-NBD-lactosylceramide	50 µg
1630-001	N-Dodecanoyl-NBD-beta-D-lactosylphingosine; N-C12:0-NBD-beta-D-lactosylphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-lactosylphingosine C ₄₈ H ₈₁ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	


Ceramide trihexosides



1067	Ceramide trihexosides	1 mg
1067-10	CTH; Gb3; globotriaosylceramide C ₆₀ H ₁₀₃ NO ₁₈ CAS#: 71965-57-6	10 mg
	Source: natural, porcine Mol. Wt.: 1126 (tetracosanoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform/methanol 2:1, DMSO, hot methanol Storage: -20°C	
	Contains hydroxy and non-hydroxy fatty acid side chains	
1513	Ceramide trihexosides (top spot)	0.5 mg
	CTH with non-hydroxy fatty acid side chain C ₅₄ H ₁₀₁ NO ₁₈	
	Source: natural, porcine Mol. Wt.: 1052 (stearoyl) Purity: 98+% by TLC Appearance: off white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1514	Ceramide trihexosides (bottom spot)	0.5 mg
	CTH with hydroxy fatty acid side chain C ₅₄ H ₁₀₁ NO ₁₉	
	Source: natural, porcine Mol. Wt.: 1068 (hydroxystearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 1:1, DMSO, hot methanol Storage: -20°C	
	References: S. Ashkenazi, T. C. Cleary, J. Clin. Microbiol., 27 , 1145, 1989 J. Ghislain et al., J. of Immunol., 153 , 3655, 1995	
1520	lyso-Ceramide trihexoside	1 mg
	lyso-CTH; lyso-globotriosylphingosine C ₃₆ H ₆₇ NO ₁₇ CAS# 126550-86-5	
	Source: semi-synthetic, porcine Mol. Wt.: 786 Purity: 98+% by TLC Appearance: film, vacuum dried Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
1523	N-Heptadecanoyl ceramide trihexoside	0.5 mg
	N-C17:0-Ceramide trihexoside; N-heptadecanoyl globotriaosylceramide C ₅₅ H ₉₉ NO ₁₈	
	Source: semi-synthetic, porcine Mol. Wt.: 1038 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 2:1:, DMSO, hot methanol Storage: -20°C	

1524	N-Tricosanoyl ceramide trihexoside N-C23:0-Ceramide trihexoside; N-tricosanoyl globotriaosylceramide $C_{59}H_{111}NO_{18}$	0.5 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1122 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 2:1; DMSO, hot methanol Storage: -20°C	
1631	N-Dodecanoyl-NBD-ceramide trihexoside N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl) ceramide trihexoside $C_{54}H_{91}N_5O_{21}$	100 µg
1631-001	Source: semi-synthetic, porcine Mol. Wt.: 1145 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1; DMSO; hot methanol Storage: -20°C	1 mg

Globosides



1068	Globosides Gb4; globotetrahexosylceramide $C_{62}H_{114}N_2O_{23}$ CAS#: 11034-93-8	5 mg
	Source: natural, porcine Mol. Wt.: 1254 (tetracosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1; DMSO, hot methanol Storage: -20°C	

Labeled glycolipids

Stable isotopes

1914	N-Stearoyl-D₃₅-psychosine, perdeuterated Cerebrosides with N-C18:0-D ₃₅ fatty acid side chain $C_{42}H_{46}D_{35}NO_8$	5 mg
	Source: semi-synthetic, bovine Mol. Wt.: 762 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, hot ethanol, chloroform/methanol 2:1 Storage: -20°C	
	Deuterium labeled stearoyl-sidechain	
1533	N-Palmitoyl-D₃-glucopsychosine, deuterated N-C16:0-D ₃ -Glucopsychosine; glucocerebroside with C16:0-D ₃ fatty acid side chain $C_{40}H_{74}D_3NO_8$	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 703 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

1534	N-Palmitoyl-D₃-lactosylceramide, deuterated N-C16:0-D ₃ -Lactosylceramide; lactosylceramide with C16:0-D ₃ fatty acid side chain C ₄₆ H ₈₄ D ₃ NO ₁₃	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 864 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide N-C18:0-D ₃ -Sulfatide C ₄₂ H ₇₈ D ₃ NO ₁₁ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 833 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/DI water 2:1:0.1 Storage: -20°C	
1537	N-Octadecanoyl-D₃-ceramide trihexoside C18:0-D ₃ -CTH; C18:0-D ₃ -Gb3; N-Octadecanoyl-D ₃ -globotriaosylceramide C ₅₄ H ₉₈ D ₃ NO ₁₈	0.5 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1059 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol 2:1:, DMSO Storage: -20°C	

Fluorescent compounds

1621	N-Hexanoyl-NBD-galactosylceramide	100 µg
1621-001	N-C6:0-NBD-beta-D-galactosylphingosine; N-C6:0-NBD-cerebroside; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-galactosylphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1622	N-Hexanoyl-NBD-glucosylceramide	100 µg
1622-001	N-C6:0-NBD-beta-D-glucosylphingosine; N-C6:0-NBD-glucosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-glucosylphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1629	N-Hexanoyl-NBD-lactosylceramide	50 ug
1629-001	N-Hexanoyl-NBD-beta-D-lactosylphingosine; N-C6:0-NBD-beta-D-lactosylphingosine; N-C6:0-NBD-lactosylceramide, fluorescent: fluorescent LC; N-(NBD-aminocaproyl)-beta-D-lactosylphingosine C ₄₂ H ₆₉ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1630	N-Dodecanoyl-NBD-lactosylceramide	50 µg
1630-001	N-Dodecanoyl-NBD-beta-D-lactosylphingosine; N-C12:0-NBD-beta-D-lactosylphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-beta-D-lactosylphingosine C ₄₈ H ₈₁ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	


1631	N-Dodecanoyl-NBD-ceramide trihexoside	100 µg
1631-001	N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl) ceramide trihexoside C ₅₄ H ₉₁ N ₅ O ₂₁	1 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1145 Purity: 98+% by TLC	
	Appearance: red-orange solid Solubility: chloroform/methanol 2:1; DMSO; hot methanol Storage: -20°C	
1632	N-Dodecanoyl-NBD-sulfatide	100 µg
1632-001	N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate; N-(NBD-aminolauroyl) sulfatide C ₄₂ H ₇₁ N ₅ O ₁₄ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 901 Purity: 98+% by TLC	
	Appearance: red-orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

See Biochemicals and Reagents section (page 85) for additional fluorescent labeled products.

Compounds with fluorescent labels other than NBD are available on custom basis. Contact Technical service for more information.

Gangliosides

The diagram below can be used with the general formulas given in the ganglioside descriptions to construct the individual structures.



1064

Gangliotetraosylceramide

Asialo GM₁; Gg4 C₆₂H₁₁₆N₂O₂₃ CAS#: 71012-19-6

1 mg

Source: semi-synthetic, bovine **Mol. Wt.:** 1240 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3,4

1512

Gangliotriosylceramide

Asialo-GM₂; Gg3 C₅₆H₁₀₄N₂O₁₈

100 µg

Source: semi-synthetic, human **Mol. Wt.:** 1078 (stearoyl) **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.1, forms micellar solution in water **Storage:** -20°C

General formula: 1,2,3

1061	Monosialoganglioside GM₁ (NH₄⁺ salt)	5 mg
1061-50	GM₁ C₇₃H₁₃₁N₃O₃₁ CAS#: 37758-47-7	50 mg
Source: natural, bovine Mol. Wt.: 1545 (stearoyl) Purity: 98+% by TLC Appearance: white solid, Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C		
General formula: 1,2,3,4,5		
References: Qing Zhang, Keiko Furukawa, Ho-Hsiang Chen, Takumi Sakakibara, Takeshi, and Koichi Furukawa. J. Biol. Chem., 281 , Issue 26, 18145-18155, <i>June 30, 2006</i> Birkles, Zeng G, Gaol, Yu R.K., Aubry J. Biochimie, 85 :455-63, 2003 Inokuchi J, Momosaki K., Shimeno H., Nagamatsu A., Radin NS. J. Cell Physiol, 141 : 573-83, 1989		
2050	N-Octadecanoyl-D₃-monosialoganglioside GM₁	0.5 mg
	N-D₃-Stearoyl-GM₁ C₇₃H₁₂₈N₃O₃₁D₃	
Source: semi-synthetic, bovine Mol. Wt.: 1548 Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C		
Reference: Qing Zhang, Keiko Furukawa, Ho-Hsiang Chen, Takumi Sakakibara, Takeshi, and Koichi Furukawa. J. Biol. Chem., 281 , Issue 26, 18145-18155, <i>June 30, 2006</i>		
1502	Monosialoganglioside GM₂ (NH₄⁺ salt)	500 µg
	GM₂ C₆₇H₁₂₁N₃O₂₆ CAS#: 19600-01-02	
Source: natural, human Mol. Wt.: 1383 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C		
General formula: 1,2,3,5		
1503	Monosialoganglioside GM₃ (NH₄⁺ salt)	500 µg
	GM₃ C₆₄H₁₁₈N₂O₂₁ CAS#: 54827-14-4	
Source: natural, bovine buttermilk Mol. Wt.: 1250 (tricosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, forms micellar solution in water Storage: -20°C		
General formula: 1,2,5		
References: Inokuchi J, Momosaki K., Shimeno H., Nagamatsu A., Radin NS. J. Cell Physiol, 141 : 573-83, 1989 Lovat P.E., Corazzari M., Disano F., Piacentini M., Redfern C.P. Cancer Lett., 228 : 105-110, 2005 Malisan R., Testi R., IUBMB Life, 57 :477-482, 2005		
1535	Monosialoganglioside GM₄, egg (NH₄⁺ salt)	0.5 mg
	GM₄ C₅₃H₈₈N₂O₁₆	
Source: natural, egg, chicken Mol. Wt.: 1008 (oleoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, forms micellar solution in water Storage: -20°C		
General formula: 1,5		
References: Ledeon, R.W., R.K. Yu, and L.F. Eng. J. Neurochem, 21 :829-839, 1978 Li Y., E. Sugiyama, T. Ariga, J. Nakayama, M. Hayama, Y. Hama, H. Nakagawa, T. Tai, S. Li, and T. Ksama. J. Lipid Res., 43 :1019-1025, 2002		

1062	Disialoganglioside GD_{1a} (NH₄⁺ salt) GD _{1a} C ₈₄ H ₁₄₈ N ₄ O ₄₀ CAS#: 12707-58-3	5 mg
	Source: natural, bovine Mol. Wt.: 1852 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C	
	General formula: 1,2,3,4,5,7	
1501	Disialoganglioside GD_{1b} (NH₄⁺ salt) GD _{1b} C ₈₄ H ₁₄₈ N ₄ O ₄₀ CAS#: 19553-76-5	1 mg
	Source: natural, bovine Mol. Wt.: 1852 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	General formula: 1,2,3,4,5,6	
1504	Disialoganglioside GD₃ (NH₄⁺ salt) GD ₃ C ₇₅ H ₁₂₅ N ₃ O ₂₉ CAS#: 62010-37-1	1 mg
	Source: natural, bovine buttermilk Mol. Wt.: 1541 (tricosanoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1, forms micellar solution in water Storage: -20°C	
	General formula: 1,2,5,6	
	References: Lovat P.E., Corazzari M., Disano F., Piacentini M., Redfern C.P. <i>Cancer Lett.</i> , 228 : 105-110, 2005 Malisan R., Testi R., IUBMB Life, 57 :477-482, 2005 Reimer A.B., Forster-Waldl E., Bramswig K.H., Pollak A., Zielinski C.C., Pehamberger H., Lode H.N., Scheiner O., Jensen-Jarolim E., <i>Eur. J. Immunol.</i> , 36 :1267-1270, 2006	
1063	Trisialoganglioside GT_{1b} (NH₄⁺ salt) GT _{1b} C ₉₅ H ₁₆₅ N ₅ O ₄₈ CAS#: 59247-13-1	5 mg
	Source: natural, bovine Mol. Wt.: 2144 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C	
	General Formula: 1,2,3,4,5,6,7	
1516	Tetrasialoganglioside GQ_{1b} (NH₄⁺ salt) GQ _{1b} C ₁₀₆ H ₁₈₂ N ₆ O ₅₆ CAS#: 68652-37-9	100 µg
	Source: natural, bovine Mol. Wt.: 2435 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C	
	General formula: 1,2,3,4,5,6,7,8	
	References: Birkles. Zeng G, Gaol, Yu R.K., Aubry J. <i>Biochimie</i> , 85 :455-63, 2003 Overell J.R., Willison H.J., <i>Curr. Opin. Neurol.</i> , 18 :562-566, 2005	
1526	Fucosylated monosialoganglioside GM₁ (NH₄⁺ salt) Fucosyl-GM ₁ C ₇₉ H ₁₄₁ N ₃ O ₃₅	500 µg
	Source: natural, porcine Mol. Wt.: 1691 (stearoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C	

1518 lyso-Monosialoganglioside GM₁ (NH₄⁺ salt) **500 µg**
lyso-GM₁ C₅₅H₉₇N₃O₃₀ CAS#: 171483-40-2

Source: semi-synthetic, bovine **Mol. Wt.:** 1279 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform/methanol/water 2:1:0.2
Storage: -20°C

1065 Purified mixed gangliosides, bovine (NH₄⁺ salt) **25 mg**
Mixed gangliosides

Source: natural, bovine **Purity:** 98+% by TLC **Appearance:** off white solid
Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water
Storage: -20°C

Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides

1525 Purified mixed gangliosides, porcine, (NH₄⁺ salt) **25 mg**

Source: natural, porcine **Purity:** 98+% by TLC **Appearance:** off-white solid
Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water
Storage: -20°C

Approximately 98% GM₁, GD_{1a}, GD_{1b} and GT_{1b}, remaining 2% other gangliosides

Glycosphingolipid reference mixes for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505 Neutral glycosphingolipid qualmix **1 mg/ml, 1 ml**
Glycosylceramides, qualitative mix

Source: natural, bovine and porcine **Appearance:** liquid **Solvent:** chloroform/methanol 2:1 **Solubility:** chloroform/methanol 2:1 **Storage:** -20°C

Contains: cerebrosides, lactosylceramide, ceramide trihexoside, globoside

1508 Monosialoganglioside mix **0.5 mg/ml, 1 ml**
GM₃, GM₂, GM₁ qualitative mix

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water 2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: GM₃, GM₂, GM₁

1509 Disialoganglioside mix **0.5 mg/ml, 1 ml**
GD₃, GD_{1a}, GD_{1b}, qualitative mix

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water 2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: GD₃, GD_{1a}, GD_{1b}

1510 Lactosylceramide and sialosyl derivatives mix **0.5 mg/ml, 1 ml**
LC, GM₃, GD₃ qualitative mix

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water 2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: LC, GM₃, GD₃

1511	Gangliotetraosylceramide and sialosyl derivatives mix asialo-GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b} qualitative mix	0.5 mg/ml, 1 ml
Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C		
Contains: asialo-GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b}		

Antibodies directed against glycolipids

These monoclonal and polyclonal antibodies are directed against the carbohydrate chains of Matreya's glycolipids. The same carbohydrate moieties are found on many glycoproteins. The antibodies are for use in ELISA or TLC immunoblotting applications (12). All antibodies are quality tested by actual performance in ELISA and TLC immunoblotting. The antibodies contain no preservatives and are shipped on dry ice.

See Literature References on page 96.

1977	Anti-ganglioside GD₃ Monoclonal antibody to GD ₃ , isotype IgG	50 µl
Source: natural, mouse hybridoma R-24 cell line Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies		
Suitable for TLC immunoblotting, ELISA		
References: Kusunoki, A. et al., Neurology, 37 :1795 1987 Pukel, C. S. et al., J. Exptl. Med., 155 :1137, 1982 Ren, S. et al., Cancer Res., 49 :7051, 1989		
1950	Anti-ganglioside asialo GM₁ Polyclonal antibody to asialo-GM ₁ , isotype IgG	100 µl
Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies		
Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to GM ₁		
References: Kusunoki, A. et al., Neurology, 37 :1795 1987 Yoshida, H. et al., J. Neurochemistry, 61 :658, 1993		
1951	Anti-ganglioside asialo-GM₂ Polyclonal antibody to asialo-GM ₂ , isotype IgG, IgM	50 µl
Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies		
Suitable for ELISA, TLC-immunoblotting		
References: Kusunoki, A. et al., Neurology, 37 :1795 1987 Yoshida, H. et al., J. Neurochemistry, 61 :658, 1993		
1954	Anti-ganglioside GM₁ Polyclonal antibody to GM ₁ , isotype IgG	100 µl
Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies		
Suitable for ELISA, TLC-immunoblotting. Slight cross reaction to asialo-GM ₁		
References: Kusunoki, A. et al., Neurology, 37 :1795 1987 Yoshida, H. et al., J. Neurochemistry, 61 :658, 1993		

1961	Anti-ganglioside GM₂ (NANA) Polyclonal antibody to GM ₂ (NANA), isotype IgG, IgM Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies Suitable for ELISA, TLC-immunoblotting	50 µl
1962	Anti-ganglioside GM₂ (NGNA) Polyclonal antibody to GM ₂ (NGNA), isotype IgG, IgM Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies Suitable for TLC immunoblotting, ELISA	50 µl
1957	Anti-ganglioside GM₄ Polyclonal antibody to GM ₄ , isotype IgG Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies Suitable for ELISA, TLC-immunoblotting	50 µl
1960	Anti-globoside GL-4 Polyclonal antibody to GL-4, isotype IgG, IgM Source: natural, rabbit Appearance: liquid Solubility: water Storage: -20°C Dry Ice Charge Applies Suitable for ELISA, TLC-immunoblotting	50 µl

Enzyme Inhibitors


Ceramide: UDPglucose Transferase. PDMP (D,L-threo-1-phenyl-2-decanoylamino-3-morpholino-1-propanol-HCl) closely resembles the natural sphingolipid substrate of brain glucosyl transferase and is a very potent and competitive inhibitor of the enzyme (13). It has been shown to block outgrowth of neurites in cultured retina and to block glucolipid synthesis in cultured 3T3 cells (14). N.S. Radin and co-workers have shown (15) that PPMP has activity equivalent to that of PDMP when cell homogenates and brain and liver microsomes are used, but it is about 20 times more potent when used with intact cells. In another paper (16), Radin's group has shown that PDMP has substantial activity against Ehrlich ascites tumors in mice. Recent publications from the laboratory of Myles Cabot (17, 18) show that PPMP can reverse multi-drug resistance in cancer cells by causing a build-up of ceramide and preventing the synthesis of glycosylated ceramides. See Literature References on page 96.

Matreya also offers the resolved D- and L-threo-isomers of PDMP and PPMP.

Protein Kinase C Inhibitor. Sphingosine is a potent and reversible inhibitor of protein kinase C (19); it also has been shown at low concentrations to stimulate DNA synthesis and act synergistically with known growth factors (20). Note that Safingol (our L-threo-dihydrosphingosine) has also been shown to partially reverse multi-drug resistance in cancer cells (18) via inhibition of protein kinase C.

Dihydroceramide desaturase Inhibitor. Cyclopropenylceramide is the first known inhibitor of this enzyme and may allow significant studies on the role of ceramide in apoptosis. Matreya is the only source for this inhibitor. (58)

Ceramidase Inhibitors. N-Oleoylethanolamine has been shown to be an efficacious inhibitor of the ceramidase found in human kidney and cerebellum (21). It is specifically an inhibitor of acid ceramidase (22) with an IC₅₀ of ca. 500 µM. N-Hexadecanoylethanolamine can be used as an inactive control. D-MAPP is a potent (IC₅₀ approximately 5 µM) inhibitor of alkaline ceramidase. Its enantiomer L-MAPP is inactive as an inhibitor and acts as a substrate for this enzyme (22,23). See Literature References on page 96.



Catalog number 1751

1751	N-Oleoylethanolamine NOE C ₂₀ H ₃₉ NO ₂ CAS#: 111-58-0	100 mg
------	---	---------------

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC,
GC Appearance: white solid **Solubility:** chloroform, ethanol, methanol, ethyl ether,
Storage: -20°C

Activity: acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. J. Lipid. Res., **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymnowska D, Wojtczak L. BBA **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. BBRC, **255**: 456-459, 1999

1786	N-Hexadecanoylethanolamine C ₁₈ H ₃₇ NO ₂ CAS# 544-31-0	100 mg
------	--	---------------


Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: inactive as acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. J. Lipid. Res., **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymnowska D, Wojtczak L. BBA **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. BBRC, **255**: 456-459, 1999

1757	Anandamide Arachidonylethanolamide; 5,8,11,14(Z,Z,Z,Z)-Eicosatetraenoyl 2'-hydroxyethyl-amide C ₂₂ H ₃₇ NO ₂ CAS#: 94421-68-8	10 mg/ml, 1 ml
	Source: synthetic Mol. Wt.: 347 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C	
	Induces apoptosis, endocannabinoid	
	References: M. van der Stelt and V. DiMarzo; Prostaglandins Other Lipid Mediat. 77 , 111, 2005 Wasilewski M., Wieckowski M.R., Dymnowska D., Wojtczak L. BBA 1657 : 151-163, 2004 C. Grimaldi, et al.; Exp. Cell Res. 312 , 363, 2006	
1807 1807-025	L-threo-Dihydrosphingosine (Safingol) L-threo-Sphinganine, C ₁₈ chain C ₁₈ H ₃₉ NO ₂ CAS#: 15639-50-6	5 mg 25 mg
	Source: synthetic Mol. Wt.: 301 Melting Point (°C): 103-114 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	References: C.W. Sachs et al., ibid., 270 , 26639, 1995 G.K. Schwartz et al., J. Natl. Cancer Inst., 87 , 1394, 1995	
	<p>Catalog number 1859</p>	
1859	D-MAPP D-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol C ₂₃ H ₃₉ NO ₂ CAS#: 143492-39-1	100 mg
	Source: synthetic Mol. Wt.: 361 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol Storage: -20°C	
	Activity: alkaline ceramidase inhibitor	
1860	L-MAPP L-erythro-2-Tetradecanoylamino-1-phenyl-1-propanol C ₂₃ H ₃₉ NO ₂ CAS#: 143492-38-0	100 mg
	Source: synthetic Mol. Wt.: 361 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol Storage: -20°C	
	Activity: inactive as alkaline ceramidase inhibitor	



Catalog number: 1886

**1886
1886-005**

N-C8:0-Cyclopropenylceramide

N-C8:0-CPPC; N-[(1R, 2S)-2-hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] octanamide; GT₁₁ C₂₇H₅₁NO₃

**1 mg
5 mg**

Source: synthetic **Mol. Wt.:** 437 **Melting Point (°C):** 69-70 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** off white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Dihydroceramide desaturase inhibitor

References:

- Jacqueline M. Kravka, Li Li, Zdzislaw M. Szulc, Jacek Bielawski, Besium Ogretmen, Yusuf A. Hannun, Lina M. Obeid, and Alicja Bielawska. *J. Biol. Chem.*, **10**, 1074/jbc.M700647200, February 5, 2007
 G. Triola, G. Fabrias, and A. Liebaria. *Agnew. Chem. Int. Ed.*, **40**, No. 10, 1960-1962, 2001
 Triola G., Fabrias G., Casas J., and Liebaria A. *J. Org. Chem.* **68** (26), 9924-9932, 2003
 Bedia C., Triola G., Casas J., Liebaria A., Fabrias G. *Or. Biomol. Chem.* **3** (20), 3707-3712, 2005

**1887
1887-005**

N-C16:0-Cyclopropenylceramide

N-C16:0-CPPC; N-[(1R, 2S)-2-hydroxy-1-hydroxymethyl-2-(2-tridecyl-1-cyclopropenyl) ethyl] hexadecamide C₃₅H₆₇NO₃


**1 mg
5 mg**

Source: synthetic **Mol. Wt.:** 550 **Melting Point (°C):** 156-157 **Purity:** 98+% by ¹H NMR; HPLC **Appearance:** off white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Dihydroceramide desaturase inhibitor

References:

- G. Triola, G. Fabrias, and A. Liebaria. *Agnew. Chem. Int. Ed.*, **40**, No. 10, 1960-1962, 2001
 Triola G., Fabrias G., Casas J., and Liebaria A. *J. Org. Chem.* **68** (26), 9924-9932, 2003
 Bedia C., Triola G., Casas J., Liebaria A., Fabrias G. *Or. Biomol. Chem.* **3** (20), 3707-3712, 2005



Catalog number 1889

1889

Conduritol B epoxide


C₆H₁₀O₅ CAS#: 6090-95-5


25 mg


Source: synthetic **Mol. Wt.:** 162 **Melting Point (°C):** 164-166 **Purity:** 98+% by TLC, NMR **Appearance:** solid **Solubility:** water, DMSO, methanol (slightly) **Storage:** -20°C

Inhibits α-glucosidase activity; specific inhibitor of glucocerebrosidase in cultured cells.

1719	D,L-threo-PDMP D,L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl C ₂₃ H ₃₈ N ₂ O ₃ •HCl CAS#: 80938-69-8	100 mg
	Source: synthetic Mol. Wt.: 427 Melting Point (°C): 158-161 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol, methanol, chloroform, DMSO Storage: -20°C	
	Activity: glucosyl ceramide synthase inhibitor	
1720	D,L-threo-PPMP D,L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl C ₂₉ H ₅₀ N ₂ O ₃ •HCl CAS#: 149022-18-4	100 mg
	Source: synthetic Mol. Wt.: 511 Purity: 98+% by TLC Appearance: white solid Solubility: alcohols, chloroform Storage: -20°C	
	Activity: glucosyl ceramide synthase inhibitor	
1749	L-threo-PDMP L-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl C ₂₃ H ₃₈ N ₂ O ₃ •HCl CAS#: 109836-81-9	10 mg
	Source: synthetic Mol. Wt.: 427 Melting Point (°C): 89-92 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	
1753	D,L-erythro-PPMP D,L-erythro-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl C ₂₉ H ₅₀ N ₂ O ₃ •HCl	100 mg
	Source: synthetic Mol. Wt.: 511 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	
1755	D,L-erythro-PDMP D,L-erythro-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl C ₂₃ H ₃₈ N ₂ O ₃ •HCl CAS#: 109760-77-2	100 mg
	Source: synthetic Mol. Wt.: 427 Melting Point (°C): 112-115 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethanol, methanol, DMSO Storage: -20°C	



1756	D-threo-PDMP D-threo-1-Phenyl-2-decanoylamino-3-morpholino-1-propanol•HCl $C_{23}H_{38}N_2O_3 \cdot HCl$ CAS#: 109836-82-0	10 mg
	Source: synthetic Mol. Wt.: 427 Melting Point (°C): 94-97 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	
	Activity: glucosyl ceramide synthase inhibitor	
	References: Nicholson K.M., Quinn D.M., Kellett G. L., Warr J.R. Br. J. Cancer 81: 423-430, 1999 Sietsma H., Veldman R.J., Vander Kolk D., Ausema B., Nijhof W., Kamps W., Vellenga E., Kok J.W. Clin. Cancer Res. 6: 942-948, 2000 Basu S., Ma R., Mikulla B., Bradley M., Moulton C., Basu M., Banerjee S., Inokuchi J. J. Glycoconj. 20: 157-168, 2003 Radin N.S. Biochem Pharmacol 57: 589-595, 1999	
		
	Catalog number 1865	
1865	D-threo-PPMP D-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl $C_{29}H_{50}N_2O_2 \cdot HCl$	10 mg
	Source: synthetic Mol. Wt.: 511 Melting Point (°C): 94-98 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	
	Activity: glucosyl ceramide synthase inhibitor	
	References: Abe A., Inokuchi J., Jimbo M., Shimeno H., Nagamatsu A., Shayman J.A., Shukla G.S., Radin N.S.J. Biochem (Tokyo) 111: 191-196, 1992 Maurer B.J., Melton L., Billups C., Cabot M.C., Reynolds C.P. J. Natl. Cancer Inst. 92: 1897-1909, 2000 Puri A., Hug P., Munoz-Barroso I., Blumenthal R. Biochem. Biophys. Res. Commun 242: 219-225, 1998 Couto A.S., Caffaro C., Uhrig M.L., Kimura E., Peres V.J., Merino E.F., Katzin A.M., Nishioka M., Nonami H., Era-Balsells R. Eur. J. Biochem. 271: 2204-2214, 2004 Morjani H., Aouali N., Belhoussine R., Veldman R.J., Levade T., Manfait M. Int. J. Cancer 94: 157-165, 2001	
1868	L-threo-PPMP L-threo-1-Phenyl-2-hexadecanoylamino-3-morpholino-1-propanol•HCl $C_{29}H_{50}N_2O_2 \cdot HCl$	10 mg
	Source: synthetic Mol. Wt.: 511 Melting Point (°C): 89-94 Purity: 98+% by TLC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C	
1800	Castanospermine 1,6,7,8-tetrahydroxyoctahydroindolizine $C_8H_{15}NO_4$ CAS#: 79831-76-8	25 mg
	Source: natural, plant Mol. Wt.: 189 Melting Point (°C): 210-215 Purity: 98+%	
	by TLC, NMR Appearance: solid Solubility: water, methanol/water, 90:10	
	Storage: -20°C	



Catalog number 1717

1717 **2-Fluoropalmitic acid** **25 mg**
 $C_{16}H_{31}FO_2$ CAS#: 89270-22-4

Source: synthetic **Mol. Wt.:** 274 **Melting Point (°C):** 83-85 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: Acyl-CoA synthase inhibitor

1718 **Methyl 2-fluoropalmitate** **10 mg**
 $C_{17}H_{33}FO_2$

Source: synthetic **Mol. Wt.:** 288 **Melting Point (°C):** 36-38 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:** -20°C

Activity: inactive ester of 2-fluoropalmitic acid


1750 **2,2-Difluoropalmitic acid** **25 mg**
 $C_{16}H_{30}F_2O_2$

Source: synthetic **Mol. Wt.:** 292 **Melting Point (°C):** 50.8-53 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C

1858 **2-Acetyl-4-(1R, 2S, 3R, 4-tetrahydroxybutyl)-imidazole** **1 mg**
THI $C_9H_{14}N_2O_5$ CAS#: 94944-70-4

Source: synthetic **Mol. Wt.:** 230 **Melting Point (°C):** n/a **Purity:** 99% by HPLC, MS, NMR **Appearance:** white solid **Solubility:** water **Storage:** -20°C

Reference:
S.R. Schwab, J.P. Pereira, M. Matloubian, Y. Xu, Y. Huang, and J.G. Cyster. Science **309**: 1735, 2005



Catalog number 1752

1752 **EST** **5 mg**
E-64-d; Loxastatin $C_{17}H_{30}N_2O_5$ CAS#: 88321-09-9

Source: synthetic **Mol. Wt.:** 342 **Melting Point (°C):** 125-127 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol
Storage: -20°C


Activity: cysteine protease inhibitor

Reference:
S. Mehdi, TIBS, **16**, April 1991


Glycerolipids

Glycerophospholipids

Natural phospholipids



Catalog number 1044



Catalog number 1046


1044	Lecithin Phosphatidylcholine; PC C ₄₄ H ₈₄ NO ₈ P CAS#: 8002-43-5 Source: natural, egg Mol. Wt.: 787 (oleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethyl ether, ethanol Storage: -20°C See Table III page 90-94 for fatty acid content	50 mg/ml, 1 ml
1070	Lecithin Phosphatidylcholine; PC C ₄₄ H ₈₄ NO ₈ P CAS#: 8002-43-5 Source: natural, bovine Mol. Wt.: 787 (oleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethyl ether Storage: -20°C See Table III page 90-94 for fatty acid content	50 mg/ml, 1 ml
1302	Lecithin Phosphatidylcholine; PC C ₄₄ H ₈₀ NO ₃ P CAS#: 8002-43-5 Source: natural, plant Mol. Wt.: 783 (linoleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethyl ether Storage: -20°C See Table III page 90-94 for fatty acid content	50 mg/ml, 1 ml
1046	lyso-Lecithin lyso-Phosphatidylcholine C ₂₄ H ₅₂ NO ₇ P CAS#: 9008-30-4 Source: natural, egg Mol. Wt.: 496 (palmitoyl) Purity: 98+% by TLC Appearance: solid Solubility: chloroform/methanol 2:1 Storage: -20°C See Table III page 90-94 for fatty acid content	50 mg
1047	Phosphatidylserine PS C ₄₂ H ₇₈ NO ₁₀ P Source: natural, bovine Mol. Wt.: 788 (oleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, toluene Storage: -20°C See Table III page 90-94 for fatty acid content	50 mg/ml, 1 ml

1048	Phosphatidylinositol (Na⁺ salt) PI C ₄₅ H ₇₈ O ₁₃ P Na CAS# 383907-36-6	10 mg/ml, 1 ml
	Source: natural, plant Mol. Wt.: 880 (linoleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethyl ether Storage: -20°C	
	See Table III page 90-94 for fatty acid content	
1336	Phosphatidylinositol, plant, soy, (K⁺ salt) C ₄₅ H ₇₈ O ₁₃ P K CAS# 383907-36-6	50 mg/ml, 1ml
	Source: natural, plant, soy Mol. Wt.: 873 (linoleoyl and pamitoyl) Purity: 98+% by TLC Appearance: tinted liquid Solvent: chloroform Solubility: chloroform, ethyl ether Storage: -20°C	
1053	Phosphatidic acid (NH₄⁺ salt) PA C ₃₉ H ₇₂ O ₈ P NH ₄	50 mg
	Source: semi-synthetic, egg Mol. Wt.: 744 (oleoyl) Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C	
	See Table III page 90-94 for fatty acid content	
1045	Phosphatidylethanolamine PE C ₄₁ H ₇₈ NO ₈ P CAS#: 39382-08-6	50 mg/ml, 1 ml
	Source: natural, egg Mol. Wt.: 744 (oleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	See Table III page 90-94 for fatty acid content	
1069	Phosphatidylethanolamine PE C ₄₁ H ₇₈ NO ₈ P CAS#: 90989-93-8	50 mg/ml, 1 ml
	Source: natural, bovine Mol. Wt.: 744 (oleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
1301	Phosphatidylethanolamine PE C ₄₁ H ₇₄ NO ₃ P CAS#: 90989-93-8	50 mg/ml, 1 ml
	Source: natural, plant Mol. Wt.: 740 (linoleoyl) Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	See Table III page 90-94 for fatty acid content	
1052	Phosphoglycerides kit	1 each
	Source: natural, egg, bovine, plant Purity: 98+% by TLC Appearance: liquid/solid Solvent: various Storage: -20°C	
	Individually packed in ampules and vials (Purity 98+%) : Phosphatidic acid NH ₄ ⁺ salt 10mg; Phosphatidylethanolamine, egg (in 1 ml CHCl ₃) 10mg; Sphingomyelin, bovine 10mg; Phosphatidylserine, bovine (in 1 ml CHCl ₃) 10 mg; Lecithin, egg (in 1 ml CHCl ₃) 10 mg; lyso-Lecithin, egg 10 mg; Cerebroside, bovine 10mg; Sulfatides, bovine 10mg; Phosphatidylinositol, Na ⁺ salt, plant (in 1 ml CHCl ₃) 3mg	

Synthetic phospholipids

These phospholipids have 98+% chemical purity except where stated and 99% fatty acid chain purity. Store at -20° C.
Solubility: see individual entries

Phosphatidic acid derivatives



Catalog number 1428

1428	1,2-Dimyristoyl-sn-glycero-3-phosphatidic acid DMPA C ₃₁ H ₆₀ O ₈ P•Na CAS#: 80724-31-8	100 mg
------	--	--------

Source: synthetic Mol. Wt.: 615 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C


1429	1,2-Dipalmitoyl-sn-glycero-3-phosphatidic acid DPPA C ₃₅ H ₆₈ O ₈ P•Na CAS#: 70240-64-1	100 mg
------	--	--------

Source: synthetic Mol. Wt.: 671 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C

1430	1,2-Distearoyl-sn-glycero-3-phosphatidic acid DSPA C ₃₉ H ₇₆ O ₈ P•Na CAS#: 108321-18-2	100 mg
------	--	--------

Source: synthetic Mol. Wt.: 727 Purity: 98+% by TLC Appearance: white solid
Solubility: chloroform/methanol/acetic acid, 4:1:0.1 Storage: -20°C

Phosphatidylcholines



Catalog number 1425

1442	1,2-Dilauroyl-sn-glycero-3-phosphorylcholine DLPC C ₃₂ H ₆₄ NO ₈ P CAS#: 18194-25-7	100 mg
------	--	--------


Source: synthetic Mol. Wt.: 622 Purity: 98+% by TLC Appearance: white solid
Solubility: methylene chloride, methanol Storage: -20°C

1425	1,2-Dimyristoyl-sn-glycero-3-phosphorylcholine DMPC C ₃₆ H ₇₂ NO ₈ P CAS#: 18194-24-6	100 mg
------	--	--------

Source: synthetic Mol. Wt.: 678 Purity: 98+% by TLC Appearance: white solid
Melting Point: 130-139°C Solubility: methylene chloride, methanol Storage: -20°C

1426	1,2-Dipalmitoyl-sn-glycero-3-phosphorylcholine DPPC C ₄₀ H ₈₀ NO ₈ P CAS#: 63-89-8	100 mg
	Source: synthetic Mol. Wt.: 734 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1400	1,2-Diheptadecanoyl-sn-glycero-3-phosphorylcholine DHDP C ₄₂ H ₈₄ NO ₈ P CAS#: 70897-27-7	50 mg
	Source: synthetic Mol. Wt.: 762 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1427	1,2-Distearoyl-sn-glycero-3-phosphorylcholine DSPC C ₄₄ H ₈₈ NO ₈ P CAS#: 816-94-4	100 mg
	Source: synthetic Mol. Wt.: 790 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1437	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylcholine POPC C ₄₂ H ₈₄ NO ₈ P CAS#: 26853-31-6	100 mg
	Source: synthetic Mol. Wt.: 760 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	
1445	1-Palmitoyl-sn-glycero-3-phosphorylcholine lyso-PPC C ₂₄ H ₅₀ NO ₇ P CAS#: 17364-16-8	100 mg
	Source: synthetic Mol. Wt.: 496 Purity: 98+% by TLC Appearance: white solid Solubility: methylene chloride, methanol Storage: -20°C	

Phosphatidylglycerols




Catalog number 1443

1443	1,2-Dilauroyl-sn-glycero-3-phosphorylglycerol DLPG C ₃₀ H ₅₈ O ₁₀ P•Na CAS#: 73548-69-3	100 mg
	Source: synthetic Mol. Wt.: 632 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1431	1,2-Dimyristoyl-sn-glycero-3-phosphorylglycerol DMPG C ₃₄ H ₆₆ O ₁₀ P•Na CAS#: 67232-80-8	100 mg
	Source: synthetic Mol. Wt.: 689 Purity: 98+% by TLC Appearance: white solid Melting Point: 120-129°C Solubility: chloroform/methanol, 5:1 Storage: -20°C	
1432	1,2-Dipalmitoyl-sn-glycero-3-phosphorylglycerol DPPG C ₃₈ H ₇₄ O ₁₀ P•Na CAS#: 67232-81-9	100 mg
	Source: synthetic Mol. Wt.: 745 Purity: 98+% by TLC Appearance: white solid Melting Point: 122-127°C Solubility: chloroform/methanol, 5:1 Storage: -20°C	

1433	1,2-Distearoyl-sn-glycero-3-phosphorylglycerol DSPG C ₄₂ H ₈₂ O ₁₀ P•Na CAS#: 4537-78-4	100 mg
Source: synthetic Mol. Wt.: 801 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		
1438	1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphorylglycerol POPG C ₄₀ H ₇₆ O ₁₀ P•Na CAS#: 81490-05-3	100 mg
Source: synthetic Mol. Wt.: 771 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol, 5:1 Storage: -20°C		

Phosphatidylethanolamines




1444	1,2-Dilauroyl-sn-glycero-3-phosphorylethanolamine DLPE C ₂₉ H ₅₈ NO ₈ P CAS#: 59752-57-7	100 mg
Source: synthetic Mol. Wt.: 579 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform + methanol mixture Storage: -20°C		
1434	1,2-Dimyristoyl-sn-glycero-3-phosphorylethanolamine DMPE C ₃₃ H ₆₆ NO ₈ P CAS# 998-07-2	100 mg
Source: synthetic Mol. Wt.: 636 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/facetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 Storage: -20°C		
1435	1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine DPPE C ₃₇ H ₇₄ NO ₈ P CAS#: 923-61-5	100 mg
Source: synthetic Mol. Wt.: 692 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/acetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 Storage: -20°C		
1436	1,2-Distearoyl-sn-glycero-3-phosphorylethanolamine DSPE C ₄₁ H ₈₂ NO ₈ P CAS#: 1069-79-0	100 mg
Source: synthetic Mol. Wt.: 748 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/acetic acid 95:5; chloroform/methanol/water/acetic acid 100:30:10:2.5 Storage: -20°C		
1439	1,2-Distearoyl-phosphatidylethanolamine-methyl-polyethyleneglycol conjugate-2000 (Na⁺ salt) DSPE-MPEG-2000 CAS#: 147867-65-0	100 mg
Source: synthetic Purity: 98+% by TLC Appearance: white solid Solubility: chloroform Storage: -20°C		

Phosphatidylinositols

The metabolism of inositol lipids is involved in the signal transduction of many hormones, neurotransmitters and growth factors (1,2). In the classical pathway, phosphatidylinositol-specific phospholipase C (PI-PLC) hydrolyzes phosphatidyl 4,5-biphosphate (PIP₂) to yield 1,2-diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃). The role of IP₃ and DAG as second messengers is well recognized.

In a second, more recently discovered pathway, the activation of phosphoinositide (PI) 3-kinase results in the formation of three novel phosphatidyl (PI) lipids phosphorylated at the D3 position of the inositol ring: PI-3-P, PI-3,4-P₂ and PI-3,4,5-P₃ (3). These D3 lipids are not known substrates for any of the phospholipase C enzymes and function as second messengers. PI 3-kinase activity is correlated with many cellular processes, including the regulation of cell growth, oncogenic transformation, chemotaxis and receptor down-regulation among others (4,5,6). The recent paper on the effect of PI3,4-P₂ on the *Akt* proto-oncogene product (9) also contains protocols for applying PIP's to cell cultures. Matreya's synthetic phosphatidylinositols and inositol phosphates are excellent tools for investigating these second messengers, understanding the enzyme mechanisms involved in phosphoinositide metabolism (7,8) and for designing therapeutic pharmacological agents. The compounds are evaluated by ¹H and ³¹P NMR to guarantee enantiomeric purity of >98%. See Literature References on page 96.

Phosphatidylinositols



Catalog number 1779

1779	Phosphatidylinositol, dipalmitoyl, (NH₄⁺ salt)	0.5 mg
1779-1	PI; DPPI (NH₄⁺ salt) C₄₁H₇₈O₁₃P•NH₄	1 mg

Source: synthetic **Mol. Wt.:** 828 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: white solid **Solubility:** chloroform/methanol/water 5:1:0.1
Storage: -20°C

1773	Phosphatidylinositol 3-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg
1773-1	DPPI-3-P; PI-3-P dipalmitoyl (NH₄⁺ salt) C₄₁H₇₇O₁₆P₂•3NH₄	1 mg
1773-5		5 mg

Source: synthetic **Mol. Wt.:** 942 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: white solid **Solubility:** chloroform/methanol/water 1:1:0.3
Storage: -20°C

References:

- B.A. Fenderson, E.M. Eddy, S. Hakomori, *BioEssays* **12**, 173, 1990
R. T. Dobrowsky et al., *ibid*, **268**, 15523, 1993
Berridge, M. J., *Nature* **361**:315, 1993
Bhamare, N. et al., 1996 *Phosphorus, Sulfur and Silicon XXX*, Overseas Publishers Association, Amsterdam B.V. 109-110:317

1780	Phosphatidylinositol 3-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg
1780-1	DPPI-3-P; PI-3-P dipalmitoyl (Na⁺ salt) C₄₁H₇₇O₁₆P₂•3Na	1 mg
1780-5		5 mg

Source: synthetic **Mol. Wt.:** 957 **Purity:** 98+% by ¹H NMR, ³¹P NMR
Appearance: white solid **Solubility:** water **Storage:** -20°C

1922	Phosphatidylinositol 4-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	
1922-1	DOPI-4-P; PI-4-P dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₅ O ₁₆ P ₂ •3NH ₄	1 mg	
1922-5		5 mg	
	Source: synthetic Mol. Wt.: 718 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: soluble in water; slightly soluble in methanol; slightly soluble in chloroform/methanol/DI water, 1:1:0.3 Storage: -20°C		
1919	Phosphatidylinositol 4-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1919-1	DPPI-4-P; PI-4-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	1 mg	
1919-5		5 mg	
	Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: methanol, chloroform/methanol/water 1:1:0.3, slightly soluble in water Storage: -20°C		
1923	Phosphatidylinositol 5-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	0
1923-1	DOPI-5-P; PI-5-P dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₅ O ₁₆ P ₂ •3NH ₄	1 mg	
1923-5		5 mg	
	Source: synthetic Mol. Wt.: 718 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: soluble in water; slightly soluble in methanol; slightly soluble in chloroform/methanol/DI water, 1:1:0.3 Storage: -20°C		
1920	Phosphatidylinositol 5-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1920-1	DPPI-5-P; PI-5-P dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₇ O ₁₆ P ₂ •3NH ₄	1 mg	
1920-5		5 mg	
	Source: synthetic Mol. Wt.: 942 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: methanol, chloroform/methanol/water 1:1:0.3, slightly soluble in water Storage: -20°C		
1781	Phosphatidylinositol bis-3,4-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1781-1	DPPI-3,4-P2; PI-3,4-P2 dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5NH ₄	1 mg	
1781-5		5 mg	
	Source: synthetic Mol. Wt.: 1056 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		
1774	Phosphatidylinositol bis-3,4-phosphate, dipalmitoyl, (Na⁺ salt)	100 µg	
1774-1	DPPI-3,4-P2; PI-3,4-P2 dipalmitoyl (Na ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5Na	1 mg	
1774-5		5 mg	
	Source: synthetic Mol. Wt.: 1081 Purity: 98+% by ¹ H NMR, ³¹ P NMR, Appearance: white solid Solubility: water Storage: -20°C		
	References: Carpenter, C. L. and L.C. Cantley, Curr. Opin. Cell Biol. 8:153, 1996 Ireton, K. et al., Science 274:80, 1996		
1784	Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (NH₄⁺ salt)	100 µg	
1784-1	DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (NH ₄ ⁺ salt) C ₂₅ H ₄₉ O ₁₉ P ₃ •5NH ₄	1 mg	
1784-5		5 mg	
	Source: synthetic Mol. Wt.: 831 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		
1778	Phosphatidylinositol bis-4,5-phosphate, dioctanoyl, (Na⁺ salt)	100 µg	
1778-1	DOPI-4,5-P2; PI-4,5-P2 dioctanoyl (Na ⁺ salt) C ₂₅ H ₄₄ O ₁₉ P ₃ •5Na	1 mg	
1778-5		5 mg	
	Source: synthetic Mol. Wt.: 856 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: water Storage: -20°C		
1777	Phosphatidylinositol bis-4,5-phosphate, dipalmitoyl, (NH₄⁺ salt)	100 µg	
1777-1	DPPI-4,5-P2; PI-4,5-P2 dipalmitoyl (NH ₄ ⁺ salt) C ₄₁ H ₇₆ O ₁₉ P ₃ •5NH ₄	1 mg	
1777-5		5 mg	
	Source: synthetic Mol. Wt.: 1056 Purity: 98+% by ¹ H NMR, ³¹ P NMR Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3 Storage: -20°C		

1782	Phosphatidylinositol bis-4,5-phosphate, dipalmitoyl, (Na^+ salt)	100 µg
1782-1	DPPI-4,5-P2; PI-4,5-P2 dipalmitoyl (Na^+ salt) $\text{C}_{41}\text{H}_{76}\text{O}_{19}\text{P}_3 \cdot 5\text{Na}$	1 mg
1782-5		5 mg
	Source: synthetic Mol. Wt.: 1081 Purity: 98+% by ^1H NMR, ^{31}P NMR	
	Appearance: white solid Solubility: water Storage: -20°C	
1921	Phosphatidylinositol tris-3,4,5-phosphate, dioctanoyl, (Na^+ salt)	100 µg
1921-1	DOPI-3,4,5-P3; PI-3,4,5-P3 dioctanoyl (Na^+ salt) $\text{C}_{25}\text{H}_{43}\text{O}_{22}\text{P}_4 \cdot 7\text{Na}$	1 mg
1921-5		5 mg
	Source: synthetic Mol. Wt.: 980 Purity: 98+% by ^1H NMR, ^{31}P NMR	
	Appearance: white solid Solubility: water Storage: -20°C	
1783	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (NH_4^+ salt)	100 µg
1783-1	DPPI-3,4,5-P3; PI-3,4,5-P3 dipalmitoyl (NH_4^+ salt) $\text{C}_{41}\text{H}_{75}\text{O}_{22}\text{P}_4 \cdot 7\text{NH}_4$	1 mg
1783-5		5 mg
	Source: synthetic Mol. Wt.: 1170 Purity: 98+% by ^1H NMR, ^{31}P NMR	
	Appearance: white solid Solubility: chloroform/methanol/water 1:1:0.3	
	Storage: -20°C	
1775	Phosphatidylinositol tris-3,4,5-phosphate, dipalmitoyl, (Na^+ salt)	100 µg
1775-1	DPPI-3,4,5-P3; PI-3,4,5-P3, dipalmitoyl (Na^+ salt) $\text{C}_{41}\text{H}_{75}\text{O}_{22}\text{P}_4 \cdot 7\text{Na}$	1 mg
1775-5		5 mg
	Source: synthetic Mol. Wt.: 1205 Purity: 98+% by ^1H NMR, ^{31}P NMR	
	Appearance: white solid Solubility: water Storage: -20°C	


References:

Carpenter, C. L. and L.C. Cantley, *Curr. Opin. Cell Biol.* **8**:153, 1996
Ireton, K. et al., *Science* **274**:80, 1996

Bacterial tetraethers

1303	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>95% pure)	5 mg
	Purified MPL of <i>Thermoplasma acidophilum</i> (>95% pure) $\text{C}_{95}\text{H}_{188}\text{O}_{16}\text{P}$	
	Source: natural, Archaeabacteria Mol. Wt.: 1618 Purity: >95% by TLC, HPLC	
	Appearance: yellow solid Solubility: chloroform/methanol 2:1, hexane/2-propanol/DI water 30:40:5 Storage: 4-8°C	
	References:	
	H.-J. Freisleben et al., <i>J. Liposome Res.</i> 3 (3), 817, 1993	
	H.-J. Freisleben et al., <i>ibid.</i> , 5 (1), 215, 1995	
	H.-J. Freisleben et al., <i>Chem. Phys. Lipids</i> 78 , 137, 1995	
	H.-J. Freisleben et al., <i>Archives Biochem. Biophys.</i> 294 (2), 418, 1992	
1303-2	Main phospholipid (MPL) of <i>Thermoplasma acidophilum</i>, (>50% pure)	50 mg
	MPL of <i>Thermoplasma acidophilum</i> (>50% pure) $\text{C}_{95}\text{H}_{188}\text{O}_{16}\text{P}$	
	Source: natural, Archaeabacteria Mol. Wt.: 1618 Purity: >50% by TLC	
	Appearance: brown viscous liquid Solubility: chloroform/methanol 2:1, hexane/2-propanol/DI water 30:40:5 Storage: 4-8°C highly hygroscopic	
	References:	
	H.-J. Freisleben et al., <i>J. Liposome Res.</i> 3 (3), 817, 1993	
	H.-J. Freisleben et al., <i>ibid.</i> , 5 (1), 215, 1995	
	H.-J. Freisleben et al., <i>Chem. Phys. Lipids</i> 78 , 137, 1995	
	H.-J. Freisleben et al., <i>Archives Biochem. Biophys.</i> 294 (2), 418, 1992	

Glycosyl glycerides



Catalog number 1058

1058	Monogalactosyldiglyceride MGDG (hydrogenated) C ₄₅ H ₈₆ O ₁₀ CAS#: 41670-62-6	10 mg
-------------	--	--------------

Source: natural, plant **Mol. Wt.:** 787 (stearoyl) **Purity:** 98+% by TLC **Appearance:** off white solid **Solubility:** chloroform/methanol/water 4:1:0.1 **Storage:** -20°C

1059	Digalactosyldiglyceride DGDG (hydrogenated) C ₅₁ H ₉₆ O ₁₅ CAS#: 92457-02-8	5 mg
-------------	--	-------------

Source: natural, plant **Mol. Wt.:** 949 (stearoyl) **Purity:** 98+% by TLC **Appearance:** off white solid **Solubility:** chloroform/methanol/water 4:1:0.1 **Storage:** -20°C

Fatty acids

Simple fatty acids

Saturated fatty acids and methyl esters

These products are 99% pure by GC. They are stable at room temperature and are supplied neat.

1200	Methyl hexanoate Methyl caproate; C6:0 methyl ester C ₇ H ₁₄ O ₂ CAS#: 106-70-7	1 g
-------------	--	------------

Source: natural, plant **Mol. Wt.:** 130 **Purity:** 99% by TLC, GC **Appearance:** liquid **Solubility:** chloroform, ethanol, ethyl ether **Storage:** room temperature

1196	Heptanoic acid C7:0 fatty acid C ₇ H ₁₄ O ₂ CAS#: 111-14-8	1 g
-------------	---	------------

Source: natural, plant **Mol. Wt.:** 130 **Purity:** 99% by TLC, GC **Appearance:** liquid **Solubility:** chloroform, ethanol, ethyl ether **Storage:** room temperature

1197	Methyl heptanoate C7:0 fatty acid methyl ester C ₈ H ₁₆ O ₂ CAS#: 106-73-0	1 g
-------------	---	------------

Source: natural, plant **Mol. Wt.:** 144 **Purity:** 99% by TLC, GC **Appearance:** liquid **Solubility:** chloroform, ethanol, ethyl ether **Storage:** room temperature

1198	Octanoic acid Caprylic acid;C8:0 acid C ₈ H ₁₆ O ₂ CAS#: 124-07-2	1 g
-------------	--	------------

Source: natural, plant **Mol. Wt.:** 144 **Purity:** 99% by TLC, GC **Appearance:** liquid **Solubility:** chloroform, ethanol, ethyl ether **Storage:** room temperature

1199	Methyl octanoate Methyl caprylate; C8:0 methyl ester C ₉ H ₁₈ O ₂ CAS#: 111-11-5	1 g
	Source: natural, plant Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1163	Nonanoic acid C9:0 fatty acid; pelargonic acid C ₉ H ₁₈ O ₂ CAS#: 112-05-0	100 mg
	Source: synthetic Mol. Wt.: 158 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1164	Methyl nonanoate C9:0 methyl ester C ₁₀ H ₂₀ O ₂ CAS#: 1731-84-6	100 mg
	Source: synthetic Mol. Wt.: 172 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1261	Methyl decanoate Methyl caprate; C10:0 methyl ester C ₁₁ H ₂₂ O ₂ CAS#: 110-42-9	500 mg
	Source: natural, plant Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane Storage: room temperature	
1165	Undecanoic acid C11:0 fatty acid C ₁₁ H ₂₂ O ₂ CAS#: 112-37-8	100 mg
	Source: synthetic Mol. Wt.: 186 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1166	Methyl undecanoate C11:0 methyl ester C ₁₂ H ₂₄ O ₂ CAS#: 1731-86-8	100 mg
	Source: synthetic Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1008	Dodecanoic acid Lauric acid; C12:0 acid C ₁₂ H ₂₄ O ₂ CAS#: 143-07-7	1 g
	Source: natural, plant Mol. Wt.: 200 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1009	Methyl dodecanoate Methyl laurate; C12:0 methyl ester C ₁₃ H ₂₆ O ₂ CAS#: 111-82-0	1 g
	Source: natural, plant Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1161	Tridecanoic acid C13:0 fatty acid C ₁₃ H ₂₆ O ₂ CAS#: 638-53-9	100 mg
	Source: synthetic Mol. Wt.: 214 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1162	Methyl tridecanoate C13:0 methyl ester C ₁₄ H ₂₈ O ₂ CAS#: 1731-88-0	100 mg
	Source: synthetic Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	

1010	Tetradecanoic acid Myristic acid; C14:0 acid C ₁₄ H ₂₈ O ₂ CAS#: 544-63-8	1 g
	Source: natural, plant Mol. Wt.: 228 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1011	Methyl tetradecanoate Methyl myristate; C14:0 methyl ester C ₁₅ H ₃₀ O ₂ CAS#: 124-10-7	1 g
	Source: natural, plant Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1012	Pentadecanoic acid C15:0 fatty acid C ₁₅ H ₃₀ O ₂ CAS#: 1002-84-2	1 g
	Source: synthetic Mol. Wt.: 242 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1013	Methyl pentadecanoate C15:0 methyl ester C ₁₆ H ₃₂ O ₂ CAS#: 7132-64-1	1 g
	Source: synthetic Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1014	Hexadecanoic acid Palmitic acid; C16:0 fatty acid C ₁₆ H ₃₂ O ₂ CAS#: 57-10-3	1 g
	Source: natural, plant Mol. Wt.: 256 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1015	Methyl hexadecanoate Methyl palmitate; C16:0 methyl ester C ₁₇ H ₃₄ O ₂ CAS#: 112-39-0	1 g
	Source: natural, plant Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1018	Heptadecanoic acid Margaric acid; C17:0 fatty acid C ₁₇ H ₃₄ O ₂ CAS#: 506-12-7	1 g
	Source: synthetic Mol. Wt.: 270 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1019	Methyl heptadecanoate Methyl margarate; C17:0 methyl ester C ₁₈ H ₃₆ O ₂ CAS#: 1731-92-6	1 g
	Source: synthetic Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1020	Octadecanoic acid Stearic acid; C18:0 fatty acid C ₁₈ H ₃₆ O ₂ CAS#: 57-11-4	1 g
	Source: natural, plant Mol. Wt.: 284 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	
1021	Methyl octadecanoate Methyl stearate; C18:0 methyl ester C ₁₉ H ₃₈ O ₂ CAS#: 112-61-8	1 g
	Source: natural, plant Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, ethyl ether Storage: room temperature	

1028	Nonadecanoic acid C19:0 fatty acid C ₁₉ H ₃₈ O ₂ CAS#: 646-30-0	100 mg
	Source: synthetic Mol. Wt.: 298 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1029	Methyl nonadecanoate C19:0 methyl ester C ₂₀ H ₄₀ O ₂ CAS#: 1731-94-8	100 mg
	Source: synthetic Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1030	Eicosanoic acid Arachidic acid; C20:0 fatty acid C ₂₀ H ₄₀ O ₂ CAS#: 506-30-9	500 mg
	Source: natural, plant Mol. Wt.: 312 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1031	Methyl eicosanoate Methyl arachidate; C20:0 methyl ester C ₂₁ H ₄₂ O ₂ CAS#: 1120-28-1	500 mg
	Source: natural, plant Mol. Wt.: 326 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1241	Heneicosanoic acid C21:0 fatty acid C ₂₁ H ₄₂ O ₂ CAS#: 2363-71-5	100 mg
	Source: synthetic Mol. Wt.: 326 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1242	Methyl heneicosanoate C21:0 methyl ester C ₂₂ H ₄₄ O ₂ CAS#: 6064-90-0	100 mg
	Source: synthetic Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1035	Docosanoic acid Behenic acid; C22:0 fatty acid C ₂₂ H ₄₄ O ₂ CAS#: 112-85-6	500 mg
	Source: natural, plant Mol. Wt.: 341 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1036	Methyl docosanoate Methyl behenate; C22:0 methyl ester C ₂₃ H ₄₆ O ₂ CAS#: 929-77-1	500 mg
	Source: natural, plant Mol. Wt.: 354 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1186	Tricosanoic acid C23:0 fatty acid C ₂₃ H ₄₆ O ₂ CAS#: 2433-96-7	100 mg
	Source: synthetic Mol. Wt.: 355 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1187	Methyl tricosanoate C23:0 methyl ester C ₂₄ H ₄₈ O ₂ CAS#: 2433-97-8	100 mg
	Source: synthetic Mol. Wt.: 368 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	

1037	Tetracosanoic acid Lignoceric acid; C24:0 fatty acid C ₂₄ H ₄₈ O ₂ CAS#: 557-59-5	100 mg
	Source: synthetic Mol. Wt.: 369 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1038	Methyl tetracosanoate Methyl lignocerate; C24:0 methyl ester C ₂₅ H ₅₀ O ₂ CAS#: 2442-49-1	100 mg
	Source: synthetic Mol. Wt.: 382 Purity: 99% by GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1251	Hexacosanoic acid Cerotic acid; C26:0 acid C ₂₆ H ₅₂ O ₂ CAS#: 506-46-7	25 mg
	Source: synthetic Mol. Wt.: 370 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	
1252	Methyl hexacosanoate Methyl cerotate; C26:0 methyl ester C ₂₇ H ₅₄ O ₂ CAS#: 5802-85-4	25 mg
	Source: synthetic Mol. Wt.: 411 Purity: 99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: room temperature	

Unsaturated fatty acids and methyl esters

Unsaturated fatty acids are easily oxidized. Flush open containers with argon or nitrogen and store at -20°C, in dark.

1157	Myristoleic acid C14:1 (cis-9) fatty acid C ₁₄ H ₂₆ O ₂ CAS#: 544-64-9	100 mg
	Source: natural, plant Mol. Wt.: 226 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1040	Methyl myristoleate C14:1 (cis-9) methyl ester C ₁₅ H ₂₈ O ₂ CAS#: 56219-06-8	100 mg
	Source: natural, plant Mol. Wt.: 240 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1243	cis-6-Hexadecenoic acid Sapienic acid C ₁₆ H ₃₀ O ₂	25 mg
	Source: synthetic Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethanol, methanol, chloroform, ethyl ether Storage: -20°C	
1016	Palmitoleic acid C16:1 (cis-9) fatty acid C ₁₆ H ₃₀ O ₂ CAS#: 373-49-9	100 mg
	Source: natural, plant Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1017	Methyl palmitoleate C16:1 (cis-9) methyl ester C ₁₇ H ₃₂ O ₂ CAS#: 1120-25-8	100 mg
	Source: natural, plant Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1147	Palmitelaidic acid C16:1 (trans-9) acid C ₁₆ H ₃₀ O ₂ CAS#: 10030-73-6	100 mg
	Source: synthetic Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1148	Methyl palmitelaidate C16:1 (trans-9) methyl ester C ₁₇ H ₃₂ O ₂ CAS#: 10030-74-7	100 mg
	Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1208	11-Hexadecenoic acid, (92% cis, 8% trans) C16:1 (cis-11) acid C ₁₆ H ₃₀ O ₂	50 mg
	Source: synthetic Mol. Wt.: 254 Purity: >98%, by TLC Appearance: liquid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C 92% cis, 8% trans by GC	
1204	Heptadecenoic acid C17:1 (cis-10) acid C ₁₇ H ₃₂ O ₂ CAS#: 29743-97-3	100 mg
	Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1203	Methyl heptadecenoate C17:1 (cis-10) methyl ester C ₁₈ H ₃₄ O ₂ CAS#: 75190-82-8	100 mg
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1022	Oleic acid C18:1 (cis-9) acid C ₁₈ H ₃₄ O ₂ CAS#: 112-80-1	1 g
	Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1023	Methyl oleate C18:1 (cis-9) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 112-62-9	1 g
	Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1149	Elaidic acid C18:1 (trans-9) acid C ₁₈ H ₃₄ O ₂ CAS#: 112-79-8	1 g
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1150	Methyl elaidate C18:1 (trans-9) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 1937-62-8	1 g
	Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1262	trans 11-Octadecenoic acid C18:1 (trans-11) acid, trans vaccenic acid C ₁₈ H ₃₄ O ₂ CAS#: 693-72-1	100 mg
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1263	Methyl trans 11-octadecenoate Methyl trans vaccenate; C18:1 (trans-11) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 6198-58-9	100 mg
	Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1024	Linoleic acid C18:2 (cis,cis-9,12) acid C ₁₈ H ₃₂ O ₂ CAS#: 60-33-3	1 g
	Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: ethyl ether, ethanol, hexane Storage: -20°C	
1025	Methyl linoleate C18:2 (cis,cis-9,12) methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 112-63-0	1 g
	Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1151	Linoelaidic acid C18:2 (trans, trans-9, 12) acid C ₁₈ H ₃₂ O ₂ CAS#: 506-21-8	100 mg
	Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1152	Methyl linoelaidate C18:2 (trans, trans-9,12) methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 2566-97-4	100 mg
	Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1266	cis-11-Octadecenoic acid cis-vaccenic acid; C18:1(cis-11) acid C ₁₈ H ₃₄ O ₂ CAS#: 506-17-2	100 mg
	Source: natural, plant Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1267	Methyl cis-11-octadecenoate Methyl cis-vaccenate; C18:1(cis-11) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 1937-63-9	100 mg
	Source: semi-synthetic, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1026	Linolenic acid C18:3 (all cis-9,12,15) acid C ₁₈ H ₃₀ O ₂ CAS#: 463-40-1	500 mg
	Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1027	Methyl linolenate C18:3 (all cis-9,12,15) methyl ester C ₁₉ H ₃₂ O ₂ CAS#: 301-00-8	500 mg
	Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1153	gamma-Linolenic acid C18:3 (all cis-6,9,12) acid C ₁₈ H ₃₀ O ₂ CAS#: 506-26-3	100 mg
	Source: natural, plant Mol. Wt.: 278 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1154	Methyl gamma-linolenate C18:3 (all cis-6,9,12) methyl ester C ₁₉ H ₃₂ O ₂ CAS#: 16326-32-2	100 mg
	Source: natural, plant Mol. Wt.: 292 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1205	Nonadecenoic acid C19:1 (cis-10) acid C ₁₉ H ₃₂ O ₂ CAS#: 73033-09-7	100 mg
	Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1206	Methyl nonadecenoate C19:1 (cis-10) methyl ester C ₂₀ H ₃₄ O ₂ CAS#: 19788-74-0	100 mg
	Source: synthetic Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1032	Eicosenoic acid C20:1 (cis-11) acid C ₂₀ H ₃₈ O ₂ CAS#: 5561-99-9	100 mg
	Source: natural, plant Mol. Wt.: 310 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1033	Methyl eicosenoate C20:1 (cis-11) methyl ester C ₂₁ H ₄₀ O ₂ CAS#: 2390-09-2	100 mg
	Source: natural, plant Mol. Wt.: 324 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1192	Eicosadienoic acid C20:2 (cis, cis-11, 14) acid C ₂₀ H ₃₆ O ₂ CAS#: 2091-39-6	100 mg
	Source: synthetic Mol. Wt.: 322 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1193	Methyl eicosadienoate C20:2 (cis, cis-11, 14) methyl ester C ₂₁ H ₃₈ O ₂ CAS#: 2463-02-7	100 mg
	Source: synthetic Mol. Wt.: 322 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1179	Methyl 5,8,11-eicosatrienoate C20:3 (all cis-5,8,11) methyl ester, mead acid methyl ester C ₂₁ H ₃₆ O ₂ CAS#: 14602-39-2	1 mg/ml, 1 ml
	Source: natural, plant Mol. Wt.: 320 Purity: 90% by TLC, GC Appearance: liquid Solvent: hexane Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1269	Methyl homogamma linolenate, C20:3n6 Methyl 8,11,14-eicosatrienoate C ₂₁ H ₃₆ O ₂	50 mg
	Source: semi-synthetic, plant Mol. Wt.: 320 Purity: 99% by TLC, GC Appearance: liquid Solubility: hexane, ethyl ether, chloroform Storage: -20°C	
1042	Arachidonic acid C20:4 (all cis-5,8,11,14) acid C ₂₀ H ₃₂ O ₂ CAS#: 506-32-1	100 mg
	Source: natural, plant Mol. Wt.: 304 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride Storage: -20°C Dry Ice Charge Applies	
1034	Methyl arachidonate C20:4 (all cis-5,8,11,14) methyl ester C ₂₁ H ₃₄ O ₂ CAS#: 2566-89-4	100 mg
	Source: natural, plant Mol. Wt.: 318 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	

1167	Eicosapentaenoic acid EPA ; omega-3 fatty acid; C20:5 (all cis-5,8,11,14,17) acid C ₂₀ H ₃₀ O ₂ CAS#: 10417-94-4	25 mg
	Source: natural, fish oil Mol. Wt.: 302 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies	
	Anti-hyperlipoproteinemic agent; 5-LOX inhibitor	
1194	Methyl eicosapentaenoate Methyl ester of omega-3 fatty acid; C20:5 (all cis-5,8,11,14,17) methyl ester C ₂₁ H ₃₂ O ₂ CAS#: 2734-47-6	25 mg
	Source: natural, fish oil Mol. Wt.: 316 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies	
1264	Docosenoic acid C22:1 (cis-13), erucic acid C ₂₂ H ₄₂ O ₂ CAS#: 112-86-7	100 mg
	Source: natural, plant Mol. Wt.: 339 Purity: >99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, hexane Storage: -20°C	
1265	Methyl docosenoate C22:1 (cis-13) methyl ester; methyl erucate C ₂₃ H ₄₄ O ₂ CAS#: 1120-34-9	100 mg
	Source: plant Mol. Wt.: 352 Purity: >99% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, hexane Storage: -20°C	
1175	Docosapentaenoic acid C22:5 (all cis-7,10,13,16,19) acid C ₂₂ H ₃₄ O ₂ CAS#: 24880-45-3	25 mg
	Source: semi-synthetic Mol. Wt.: 330 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether, hexane Storage: -20°C Dry Ice Charge Applies	
1244	Methyl docosapentaenoate C22:5 (all cis-7,10,13,16,19) methyl ester C ₂₃ H ₃₆ O ₂ CAS#: 108698-02-8	25 mg
	Source: semi-synthetic Mol. Wt.: 344 Purity: 98+% by TLC, GC Appearance: liquid Solubility: ethyl ether, ethanol, hexane, Storage: -20°C Dry Ice Charge Applies	
1136	Docosahexaenoic acid DHA; C22:6, (all cis-4,7,10,13,16,19) omega-3 fatty acid C ₂₂ H ₃₂ O ₂ CAS#: 6217-54-5	100 mg
	Source: natural, plant Mol. Wt.: 328 Purity: 99% by TLC, GC Appearance: liquid Solubility: ethyl ether, hexane, methylene chloride, ethanol, DMSO, DMF Storage: -20°C Dry Ice Charge Applies	
1041	Methyl docosahexaenoate C22:6 (all cis-4,7,10,13,16,19) methyl ester; methyl ester of omega-3 fatty acid C ₂₃ H ₃₄ O ₂ CAS#: 2566-90-7	100 mg
	Source: natural, plant Mol. Wt.: 342 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C Dry Ice Charge Applies	
1155	Nervonic acid C24:1 (cis-15) acid C ₂₄ H ₄₆ O ₂ CAS#: 506-37-6	100 mg
	Source: synthetic Mol. Wt.: 367 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1156	Methyl nervonate C24:1 (cis-15) methyl ester C ₂₅ H ₄₈ O ₂ CAS#: 2733-88-2	100 mg
	Source: synthetic Mol. Wt.: 381 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
Trans fatty acids and methyl esters		
1147	Palmitelaidic acid C16:1 (trans-9) acid C ₁₆ H ₃₀ O ₂ CAS#: 10030-73-6	100 mg
	Source: synthetic Mol. Wt.: 254 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1148	Methyl palmitelaidate C16:1 (trans-9) methyl ester C ₁₇ H ₃₂ O ₂ CAS#: 10030-74-7	100 mg
	Source: synthetic Mol. Wt.: 268 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1149	Elaidic acid C18:1 (trans-9) acid C ₁₈ H ₃₄ O ₂ CAS#: 112-79-8	1 g
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1150	Methyl elaidate C18:1 (trans-9) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 1937-62-8	1 g
	Source: natural, plant Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1262	trans 11-Octadecenoic acid C18:1 (trans-11) acid; trans vaccenic acid C ₁₈ H ₃₄ O ₂ CAS#: 693-72-1	100 mg
	Source: synthetic Mol. Wt.: 282 Purity: 99% by TLC, GC Appearance: solid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1263	Methyl trans 11-octadecenoate Methyl trans vaccenate; C18:1 (trans-11) methyl ester C ₁₉ H ₃₆ O ₂ CAS#: 6198-58-9	100 mg
	Source: synthetic Mol. Wt.: 296 Purity: 99% by TLC, GC Appearance: clear liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1151	Linoelaidic acid C18:2 (trans, trans-9, 12) acid C ₁₈ H ₃₂ O ₂ CAS#: 506-21-8	100 mg
	Source: natural, plant Mol. Wt.: 280 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	
1152	Methyl linoelaidate C18:2 (trans, trans-9,12) methyl ester C ₁₉ H ₃₄ O ₂ CAS#: 2566-97-4	100 mg
	Source: natural, plant Mol. Wt.: 294 Purity: 99% by TLC, GC Appearance: liquid Solubility: chloroform, hexane, ethyl ether Storage: -20°C	

1131 Cis-trans Isomer standard **5 mg/ml, 5 ml**
Qualitative mix

Source: margarine **Appearance:** liquid **Solvent:** 5ml methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Analysis of positional cis-trans fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of cis-trans fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 trans isomers (4 peaks), C18:1 cis & trans isomers (2 peaks), C18:1 cis isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

1181 9(E),11(E)-Octadecadienoic acid **25 mg**
9-trans, 11-trans CLA C₁₈H₃₂O₂ CAS#: 544-71-8

Source: synthetic **Mol. Wt.:** 280 **Melting Point (°C):** 55-57 **Purity:** 98+% by TLC,
GC Appearance: off-white solid **Solubility:** chloroform, ethanol, hexane, methanol
Storage: -20°C


Conjugated linoleic acid isomers (CLA)

Linoleic acid is an essential fatty acid (18:2 ω 6) of which several naturally occurring conjugated derivatives have been identified. These derivatives, called "conjugated linoleic acid" or CLA can have the two double bonds mainly in the 9 and 11 or in the 10 and 12 positions, resulting in eight possible geometric isomers. CLA occurs in meat (24) and dairy products (25,35). In both cases, the 9(Z),11(E)-isomer is predominant and is thought to be the biologically active form. CLA assimilated through the diet of animals is found in the intestinal musosa, liver and adipose tissue (26). See also review article by Parodi (35). CLA has several biological properties. Its anti-carcinogenic activity has been demonstrated by its ability to inhibit chemically induced tumor formation in animal models of carcinogenesis (24,27,28,29). The addition of CLA to culture medium suppresses the *in vitro* growth of human melanoma, colorectal and breast cancer cells (30). CLA also exhibits anti-atherogenic activity. Addition of CLA to a controlled atherogenic diet significantly reduced the development of atherosclerosis in hamsters and rabbits (31,32). Animals fed a diet containing CLA also had lower levels of low-density-lipoprotein (LDL) cholesterol. CLA may be involved in regulating fat and protein metabolism (33,34). Several species of animals fed CLA-supplemented diets showed improved feed efficiency. Lean body mass increased while body fat was reduced. This seems to be due, mainly or exclusively, to the 10(E),12(Z)-isomer (catalog # 1249, see below). CLA competes with linoleate for $\Delta 6$ desaturase (36). Dietary CLA normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat (40) via activation of PPAR γ , a result which bears on the possible ameliorization or prevention of NIDDM. The 11(Z),13(E)-isomer (catalog # 1259) has been shown to be concentrated in the heart and in mitochondria. **See Literature References on page 96.**

CLA Research is Being Redone With Our Highly Pure Isomers

Most studies to date have utilized a mixture of CLA isomers containing less than 30% of the presumed active 9(Z),11(E)-isomer (37,38). In addition to the 9,11- and 10,12-isomers, 8,10- and 11,13-isomers have recently been identified in the widely used mixture (38,39). Matreya offers a highly pure CLA which is 98+% the active 9,11-“cis, trans” isomer. The corresponding “trans,trans” and “cis,cis” isomers are also available. In addition, we now offer the pure 10(E),12(Z)-isomer, which has been widely sought for comparison studies.


See Literature References on page 96.



1245	9(Z),11(E)-Octadecadienoic acid	25 mg
1245-1	9-cis, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	1 g
1245-10		10 g
	Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: tinted oil	
	Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1255	Methyl 9(Z), 11(E)-octadecadienoate	25 mg
	Methyl ester of CLA (9-cis, 11-trans) C ₁₉ H ₃₄ O ₂	
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil	
	Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1181	9(E),11(E)-Octadecadienoic acid	25 mg
	9-trans, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-71-8	
	Source: synthetic Mol. Wt.: 280 Melting Point (°C): 55-57 Purity: 98+% by TLC, GC Appearance: off-white solid Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1257	Methyl 9(E),11(E)-octadecadienoate	25 mg
	Methyl ester of CLA (9-trans, 11-trans) C ₁₉ H ₃₄ O ₂	
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil	
	Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	

1248	9(Z),11(Z)-Octadecadienoic acid	25 mg
1248-1	9-cis, 11-cis CLA C ₁₈ H ₃₂ O ₂ CAS#: 544-40-7	1 g
	Source: synthetic Mol. Wt.: 280 Melting Point (°C): 40-42 Purity: 96+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol, ethyl ether Storage: -20°C	
1256	Methyl 9(Z), 11(Z)-octadecadienoate	25 mg
	Methyl ester of CLA (9-cis, 11-cis) C ₁₉ H ₃₄ O ₂	
	Source: synthetic Mol. Wt.: 294 Purity: 96+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1249	10(E),12(Z)-Octadecadienoic acid	25 mg
1249-1	10-trans, 12-cis CLA C ₁₈ H ₃₂ O ₂ CAS#: 2420-44-2	1 g
1249-10	Source: synthetic Mol. Wt.: 280 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	10 g
1254	Methyl 10(E), 12(Z)-octadecadienoate	25 mg
	Methyl ester of CLA (10-trans, 12-cis) C ₁₉ H ₃₄ O ₂	
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1259	11(Z), 13(E)-Octadecadienoic acid	25 mg
	11-cis, 13-trans CLA C ₁₈ H ₃₂ O ₂	
	Source: synthetic Mol. Wt.: 280 Purity: 77% cis, trans; 2 % cis, cis; 6% trans, trans by TLC, GC Appearance: tinted oil Solubility: chloroform, ethanol, hexane, methanol Storage: -20°C	
1247-1	9(Z),11(E)-Octadecadienoic acid	1 g
1247-10	9-cis, 11-trans CLA C ₁₈ H ₃₂ O ₂ CAS#: 2540-56-9	10 g
	Source: synthetic Mol. Wt.: 280 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1%(E),(E) by TLC, GC Appearance: tinted oil Solubility: ethanol, ethyl ether, hexane Storage: -20°C	
1258	Methyl 9(Z),11(E)-octadecadienoate	25 mg
	Methyl ester of CLA (9-cis, 11-trans) C ₁₉ H ₃₄ O ₂	
	Source: synthetic Mol. Wt.: 294 Purity: 74% 9(Z),11(E); 17%(Z),(Z); 1% (E),(E) by TLC, GC Appearance: clear oil Solubility: chloroform, ethyl ether, hexane Storage: -20°C	

Other CLA products and derivatives




Catalog number 1793

1793	8-(5-Hexyl-2-furyl)-octanoic acid Furan fatty acid; 9,12-epoxy-9,11-octadecadienoic acid CAS#: 4179-44-6	25 mg
	Source: synthetic Mol. Wt.: 294 Purity: 98+% by TLC, GC Appearance: oil Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1409	1-Stearoyl-2-linoleoyl-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
	Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C	
1410	1-Stearoyl-2-[9(Z),11(E)-octadecadienoyl]-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
	Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C	
1411	1-Stearoyl-2-[10(E),12(Z)-octadecadienoyl]-sn-glycero-3-phosphorylcholine C ₄₄ H ₈₄ NO ₈ P	25 mg/ml, 1ml
	Source: synthetic Mol. Wt.: 786 Purity: 98+% by TLC Appearance: liquid Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C	
1794	Methyl 8-(5-hexyl-2-furyl)-octanoate Methyl ester of furan fatty acid C ₁₉ H ₃₂ O ₃ CAS#: 10038-16-1	25 mg
	Source: synthetic Mol. Wt.: 308 Purity: 98+% by TLC, GC Appearance: oil Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	

Hydroxy fatty acids

2-Hydroxy fatty acids and methyl esters

These products are racemic and 98+% pure by GC and TLC. The 2-hydroxy fatty acids are components of glycosphingolipids and are involved in fatty acid degradation. They are stable and are supplied neat in vials.




Catalog number 1758

1758	2-Hydroxydecanoic acid	50 mg
1758-1	2-Hydroxy C10:0 acid C ₁₀ H ₂₀ O ₃ CAS#: 5393-81-7	1 g
	Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1759	Methyl 2-hydroxydecanoate	50 mg
1759-1	2-Hydroxy C10:0 methyl ester C ₁₁ H ₂₂ O ₃ CAS#: 71271-24-4	1 g
	Source: synthetic Mol. Wt.: 202 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1701	2-Hydroxydodecanoic acid	50 mg
1701-1	2-Hydroxy C12:0 acid C ₁₂ H ₂₄ O ₃ CAS#: 2984-55-6	1 g
	Source: synthetic Mol. Wt.: 216 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1702	Methyl 2-hydroxydodecanoate	50 mg
1702-1	2-Hydroxy C12:0 methyl ester C ₁₃ H ₂₆ O ₃ CAS#: 51067-85-7	1 g
	Source: synthetic Mol. Wt.: 230 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1703	2-Hydroxytetradecanoic acid	50 mg
1703-1	2-Hydroxy C14:0 acid C ₁₄ H ₂₈ O ₃ CAS#: 2507-55-3	1 g
	Source: synthetic Mol. Wt.: 244 Melting Point (°C): 81-82 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, methanol Storage: -20°C	
1704	Methyl 2-hydroxytetradecanoate	50 mg
1704-1	2-Hydroxy C14:0 methyl ester C ₁₅ H ₃₀ O ₃ CAS#: 56009-40-6	1 g
	Source: synthetic Mol. Wt.: 258 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1705	2-Hydroxyhexadecanoic acid	50 mg
1705-1	2-Hydroxy C16:0 acid C ₁₆ H ₃₂ O ₃ CAS#: 764-67-0	1 g
	Source: synthetic Mol. Wt.: 272 Melting Point (°C): 86-87 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1706	Methyl 2-hydroxyhexadecanoate	50 mg
1706-1	2-Hydroxy C16:0 methyl ester C ₁₇ H ₃₄ O ₃ CAS#: 16742-51-1	1 g
	Source: synthetic Mol. Wt.: 286 Melting Point (°C): 59-60 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	
1707	2-Hydroxyoctadecanoic acid	50 mg
1707-1	2-Hydroxy C18:0 acid C ₁₈ H ₃₆ O ₃ CAS#: 629-22-1	1 g
	Source: synthetic Mol. Wt.: 300 Melting Point (°C): 92-93 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C	
1708	Methyl 2-hydroxyoctadecanoate	50 mg
1708-1	2-Hydroxy C18:0 methyl ester C ₁₉ H ₃₈ O ₃ CAS#: 2420-35-1	1 g
	Source: synthetic Mol. Wt.: 315 Melting Point (°C): 64-66 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C	

1709	2-Hydroxyeicosanoic acid	25 mg
1709-0.5	2-Hydroxy C20:0 acid C ₂₀ H ₄₀ O ₃ CAS#: 16742-48-6	0.5 g
Source: synthetic Mol. Wt.: 329 Melting Point (°C): 91-92 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		
1710	Methyl 2-hydroxyeicosanoate	25 mg
1710-0.5	2-Hydroxy C20:0 methyl ester C ₂₁ H ₄₂ O ₃ CAS#: 16742-49-7	0.5 g
Source: synthetic Mol. Wt.: 343 Melting Point (°C): 62-64 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C		
1711	2-Hydroxydocosanoic acid	25 mg
1711-0.5	2-Hydroxy C22:0 acid C ₂₂ H ₄₄ O ₃ CAS#: 13980-14-8	0.5 g
Source: synthetic Mol. Wt.: 366 Melting Point (°C): 96-97 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		
1712	Methyl 2-hydroxydocosanoate	25 mg
1712-0.5	2-Hydroxy C22:0 methyl ester C ₂₃ H ₄₆ O ₃ CAS#: 13980-17-1	0.5 g
Source: synthetic Mol. Wt.: 371 Melting Point (°C): 72-73 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C		
1713	2-Hydroxytricosanoic acid	10 mg
	2-Hydroxy C23:0 acid C ₂₃ H ₄₆ O ₃ CAS#: 2718-37-8	
Source: synthetic Mol. Wt.: 371 Melting Point (°C): 98-99 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		
1714	Methyl 2-hydroxytricosanoate	10 mg
	2-Hydroxy C23:0 methyl ester C ₂₄ H ₄₈ O ₃ CAS#: 118745-41-8	
Source: synthetic Mol. Wt.: 385 Melting Point (°C): 68-70 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C		
1715	2-Hydroxytetracosanoic acid	5 mg
	2-Hydroxy C24:0 acid; cerebronic acid C ₂₄ H ₄₈ O ₃ CAS#: 544-57-0	
Source: synthetic Mol. Wt.: 385 Melting Point (°C): 101-104 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform/methanol 5:1 Storage: -20°C		
1716	Methyl 2-hydroxytetracosanoate	5 mg
	2-Hydroxy C24:0 methyl ester C ₂₅ H ₅₀ O ₃ CAS#: 2433-95-6	
Source: synthetic Mol. Wt.: 399 Melting Point (°C): 64-65 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether Storage: -20°C		
1722	2-Hydroxy methyl ester mix	10 mg/ml, 1 ml
	Quantitative mixture	
Source: synthetic Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C		
Contains: 2-OH C14:0, 20%; 2-OH C16:0, 20%; 2-OH C18:0, 15%; 2-OH C20:0, 15.0%; 2-OH C22:0, 10%; 2-OH C23:0, 10%; 2-OH C24:0, 10%		

3-Hydroxy fatty acids and methyl esters

These products are racemic and 98+% pure by GC and TLC. 3-Hydroxy fatty acids occur in the lipid fraction of many microorganisms and are useful in the typing of microbial isolates. They are stable and are supplied neat in vials.




Catalog number 1747

1747	3-Hydroxyhexanoic acid	25 mg
1747-0.5	3-Hydroxy C6:0 acid C ₆ H ₁₂ O ₃ CAS#: 10191-24-9	0.5 g
	Source: synthetic Mol. Wt.: 132 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1748	Methyl 3-hydroxyhexanoate	25 mg
1748-0.5	3-Hydroxy C6:0 methyl ester C ₇ H ₁₄ O ₃ CAS#: 21188-58-9	0.5 g
	Source: synthetic Mol. Wt.: 146 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1745	3-Hydroxyoctanoic acid	25 mg
1745-0.5	3-Hydroxy C8:0 acid C ₈ H ₁₆ O ₃ CAS#: 88930-08-9	0.5 g
	Source: synthetic Mol. Wt.: 160 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1746	Methyl 3-hydroxyoctanoate	25 mg
1746-0.5	3-Hydroxy C8:0 methyl ester C ₉ H ₁₈ O ₃ CAS#: 85549-54-8	0.5 g
	Source: synthetic Mol. Wt.: 174 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1725	3-Hydroxynonanoic acid	25 mg
1725-0.5	3-Hydroxy C9:0 acid C ₉ H ₁₈ O ₃ CAS#: 88930-09-0	0.5 g
	Source: synthetic Mol. Wt.: 174 Melting Point (°C): 60-62 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1726	Methyl 3-hydroxynonanoate	25 mg
1726-0.5	3-Hydroxy C9:0 methyl ester C ₁₀ H ₂₀ O ₃ CAS#: 83968-06-3	0.5 g
	Source: synthetic Mol. Wt.: 188 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
1727	3-Hydroxydecanoic acid	25 mg
1727-0.5	3-Hydroxy C10:0 acid C ₁₀ H ₂₀ O ₃ CAS#: 5561-87-5	0.5 g
	Source: synthetic Mol. Wt.: 188 Melting Point (°C): 57-60 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
1728	Methyl 3-hydroxydecanoate	25 mg
1728-0.5	3-Hydroxy C10:0 methyl ester C ₁₁ H ₂₂ O ₃ CAS#: 62675-82-5	0.5 g
	Source: synthetic Mol. Wt.: 202 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C	

1729	3-Hydroxyundecanoic acid	25 mg
1729-0.5	3-Hydroxy C11:0 acid C ₁₁ H ₂₂ O ₃ CAS#: 40165-88-6	0.5 g
Source: synthetic Mol. Wt.: 202 Melting Point (°C): 74-76 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1730	Methyl 3-hydroxyundecanoate	25 mg
1730-0.5	3-Hydroxy C11:0 methyl ester C ₁₂ H ₂₄ O ₃ CAS#: 127593-21-9	0.5 g
Source: synthetic Mol. Wt.: 216 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1731	3-Hydroxydodecanoic acid	25 mg
1731-0.5	3-Hydroxy C12:0 acid C ₁₂ H ₂₄ O ₃ CAS#: 8355-89-3	0.5 g
Source: synthetic Mol. Wt.: 216 Melting Point (°C): 71-72 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		
1732	Methyl 3-hydroxydodecanoate	25 mg
1732-0.5	3-Hydroxy C12:0 methyl ester C ₁₃ H ₂₆ O ₃ CAS#: 85464-97-7	0.5 g
Source: synthetic Mol. Wt.: 230 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethanol, ethyl ether Storage: -20°C		
1733	3-Hydroxytridecanoic acid	25 mg
1733-0.5	3-Hydroxy C13:0 acid C ₁₃ H ₂₆ O ₃ CAS#: 32602-69-0	0.5 g
Source: synthetic Mol. Wt.: 230 Melting Point (°C): 80-83 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1734	Methyl 3-hydroxytridecanoate	25 mg
1734-0.5	3-Hydroxy C13:0 methyl ester C ₁₄ H ₂₈ O ₃	0.5 g
Source: synthetic Mol. Wt.: 244 Purity: 98+% by TLC, GC Appearance: liquid Solubility: chloroform, ethyl ether Storage: -20°C		
1735	3-Hydroxytetradecanoic acid	25 mg
1735-0.5	3-Hydroxy C14:0 acid C ₁₄ H ₂₈ O ₃ CAS#: 3422-31-9	0.5 g
Source: synthetic Mol. Wt.: 244 Melting Point (°C): 80-81 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1736	Methyl 3-hydroxytetradecanoate	25 mg
1736-0.5	3-Hydroxy C14:0 methyl ester C ₁₅ H ₃₀ O ₃ CAS#: 55682-83-2	0.5 g
Source: synthetic Mol. Wt.: 258 Melting Point (°C): 36-37 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethyl ether, methanol Storage: -20°C		
1739	3-Hydroxyhexadecanoic acid	25 mg
1739-0.5	3-Hydroxy C16:0 acid C ₁₆ H ₃₂ O ₃ . CAS#: 928-17-6	0.5 g
Source: synthetic Mol. Wt.: 272 Melting Point (°C): 85-86 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1740	Methyl 3-hydroxyhexadecanoate	25 mg
1740-0.5	3-Hydroxy C16:0 methyl ester C ₁₇ H ₃₄ O ₃ CAS#: 51883-36-4	0.5 g
Source: synthetic Mol. Wt.: 286 Melting Point (°C): 43-45 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		

1741	3-Hydroxyheptadecanoic acid	25 mg
1741-0.5	3-Hydroxy C17:0 acid C ₁₇ H ₃₄ O ₃	0.5 g
Source: synthetic Mol. Wt.: 286 Melting Point (°C): 93-95 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		
1742	Methyl 3-hydroxyheptadecanoate	25 mg
1742-0.5	3-Hydroxy C17:0 methyl ester C ₁₈ H ₃₆ O ₃	0.5 g
Source: synthetic Mol. Wt.: 300 Melting Point (°C): 53-55 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		
1743	3-Hydroxyoctadecanoic acid	25 mg
1743-0.5	3-Hydroxy C18:0 acid C ₁₈ H ₃₆ O ₃ CAS#: 45261-96-9	0.5 g
Source: synthetic Mol. Wt.: 300 Melting Point (°C): 52-54 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		
1744	Methyl 3-hydroxyoctadecanoate	25 mg
1744-0.5	3-Hydroxy C18:0 methyl ester C ₁₉ H ₃₈ O ₃ CAS#: 14531-40-9	0.5 g
Source: synthetic Mol. Wt.: 315 Melting Point (°C): 52-54 Purity: 98+% by TLC, GC Appearance: white solid Solubility: ethanol, methanol Storage: -20°C		

Omega hydroxy fatty acids



Catalog number 1754

1754	Royal Jelly acid	50 mg
1754-0.5	10-Hydroxy-2-(E)-decenoic acid; omega-hydroxy C10:1 (2-trans) C ₁₀ H ₁₈ O ₃ CAS#: 14113-05-4	0.5 g
Source: synthetic Mol. Wt.: 186 Melting Point (°C): 63-65 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: -20°C		
1881	15-Hydroxypentadecanoic acid	25 mg
	omega-Hydroxy C15:0 C ₁₅ H ₃₀ O ₃ CAS#: 4617-33-8	
Source: synthetic Mol. Wt.: 258 Melting Point (°C): 84-86 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, ethanol, methanol Storage: room temperature		
1882	Methyl 15-hydroxypentadecanoate	25 mg
	omega-Hydroxy C15:0 fatty acid methyl ester C ₁₆ H ₃₂ O ₃ CAS#: 76529-42-5	
Source: synthetic Mol. Wt.: 272 Melting Point (°C): 50-52 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature		
1760	17-Hydroxyheptadecanoic acid	25 mg
	omega-Hydroxy C17:0 fatty acid C ₁₇ H ₃₄ O ₃ CAS#: 13099-34-8	
Source: synthetic Mol. Wt.: 286 Melting Point (°C): 93-95 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature		

1761	Methyl 17-hydroxyheptadecanoate omega-Hydroxy C17:0 fatty acid methyl ester C ₁₈ H ₃₆ O ₃ CAS#: 94036-00-7	25 mg
	Source: synthetic Mol. Wt.: 300 Melting Point (°C): 59-63 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1877	20-Hydroxyeicosanoic acid omega-Hydroxy C20:0 fatty acid C ₂₀ H ₄₀ O ₃	25 mg
	Source: synthetic Mol. Wt.: 328 Melting Point (°C): 96-98 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature	
1878	Methyl 20-hydroxyeicosanoate omega-Hydroxy C20:0 fatty acid methyl ester C ₂₁ H ₄₂ O ₃	25 mg
	Source: synthetic Mol. Wt.: 342 Melting Point (°C): 69-71 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1879	21-Hydroxyheneicosanoic acid omega-Hydroxy C21:0 fatty acid C ₂₁ H ₄₂ O ₃	25 mg
	Source: synthetic Mol. Wt.: 342 Melting Point (°C): 72-75 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature	
1880	Methyl 21-hydroxyheneicosanoate omega-Hydroxy C21:0 fatty acid methyl ester C ₂₂ H ₄₄ O ₃	25 mg
	Source: synthetic Mol. Wt.: 356 Melting Point (°C): 73-76 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1818	22-Hydroxydocosanoic acid Phellonic acid; omega-hydroxy C22:0 fatty acid C ₂₂ H ₄₄ O ₃	25 mg
	Source: synthetic Mol. Wt.: 356 Melting Point (°C): 100-102 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol Storage: room temperature	
1819	Methyl 22-hydroxydocosanoate omega-Hydroxy C22:0 fatty acid methyl ester C ₂₃ H ₄₆ O ₃	25 mg
	Source: synthetic Mol. Wt.: 370 Melting Point (°C): 73-75 Purity: 98+% by TLC, GC Appearance: white solid Solubility: chloroform, warm ethanol, ethyl ether Storage: room temperature	
1883	Methyl 27-hydroxyheptacosanoate omega-Hydroxy C27:0 fatty acid methyl ester C ₂₈ H ₅₆ O ₃	25 mg
	Source: synthetic Mol. Wt.: 440 Melting Point (°C): 85-89 Purity: 97+% by TLC, GC Appearance: white solid Solubility: chloroform Storage: room temperature	
1884	Methyl 30-hydroxytriacontanoate omega-Hydroxy C30:0 fatty acid methyl ester C ₃₁ H ₆₂ O ₃	25 mg
	Source: synthetic Mol. Wt.: 482 Melting Point (°C): 88-91 Purity: 97+% by TLC, GC Appearance: white solid Solubility: chloroform Storage: room temperature	

Other hydroxy fatty acids

1815	Methyl threo-2,3-dihydroxypalmitate 2,3-Dihydroxy C16:0 fatty acid methyl ester	10 mg <chem>C17H34O4</chem>
-------------	---	---------------------------------------

Source: synthetic **Mol. Wt.:** 302 **Melting Point (°C):** 77-79 **Purity:** 98+% by TLC,
GC Appearance: white solid **Solubility:** chloroform, ethanol, methanol **Storage:**
 room temperature

1182	Ricinelaidic acid 12-Hydroxy C18:1 (9-trans) fatty acid	100 mg <chem>C18H34O3</chem> CAS#: 82188-83-8
-------------	---	--



Source: synthetic **Mol. Wt.:** 298 **Melting Point (°C):** 50-53 **Purity:** 98+% by TLC,
GC Appearance: white solid **Solubility:** ethanol, methanol **Storage:** -20°C

1183	Methyl ricinelaivate 12-Hydroxy C18:1 (9-trans) methyl ester	100 mg <chem>C19H36O3</chem> CAS#: 7706-01-6
-------------	--	---

Source: synthetic **Mol. Wt.:** 312 **Purity:** 98+% by TLC, GC **Appearance:** white
 solid **Solubility:** ethanol, methanol **Storage:** -20°C

1766	6-Hydroxyoctadecanoic acid 6-Hydroxy C18:0 fatty acid	10 mg <chem>C18H36O3</chem>
-------------	---	---------------------------------------

Source: synthetic **Mol. Wt.:** 300 **Melting Point (°C):** 80-82 **Purity:** 98+% by TLC,
GC Appearance: white solid **Solubility:** ethanol, methanol **Storage:** room
 temperature



1898	Methyl D, L-threo-corynomycolate Hydroxy fatty acid with long branched chain	25 mg <chem>C33H66O3</chem>
-------------	--	---------------------------------------


Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 70 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform **Storage:** room temperature

1899	Methyl D,L-erythro-corynomycolate Hydroxy fatty acid with long branched chain	25 mg <chem>C33H66O3</chem>
-------------	---	---------------------------------------

Source: synthetic **Mol. Wt.:** 511 **Melting Point (°C):** 58 **Purity:** 98+% by TLC
Appearance: white solid **Solubility:** chloroform **Storage:** room temperature

Branched and cyclic fatty acids


iso-Fatty acids and esters



Catalog number 1656

1656	Methyl 11-methyldodecanoate iso-Tridecanoic methyl ester; iso C13 methyl ester C ₁₄ H ₂₈ O ₂ CAS#: 5129-57-7	20 mg
	Source: synthetic Mol. Wt.: 228 Purity: 98+% by GC Appearance: liquid Solubility: hexane, ethyl ether, methylene chloride Storage: -20°C	
1657	Methyl 12-methyltridecanoate iso-Tetradecanoic methyl ester; iso C14 methyl ester C ₁₅ H ₃₀ O ₂ CAS#: 5129-58-8	20 mg
	Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1605	13-Methyltetradecanoic acid iso-Pentadecanoic acid; iso C15 acid C ₁₅ H ₃₀ O ₂	20 mg
	Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: white solid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1600	Methyl 13-methyltetradecanoate iso-Pentadecanoic methyl ester; iso C15 methyl ester C ₁₆ H ₃₂ O ₂ CAS#: 5129-59-9	20 mg
	Source: synthetic Mol. Wt.: 256 Purity: 98+% by GC Appearance: clear liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1601	Methyl 14-methylpentadecanoate iso-Palmitic methyl ester; iso C16 methyl ester C ₁₇ H ₃₄ O ₂ CAS#: 5129-60-2	20 mg
	Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: clear liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1606	15-Methylhexadecanoic acid iso-Heptadecanoic acid; iso C17 acid C ₁₇ H ₃₄ O ₂	20 mg
	Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: white solid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1602	Methyl 15-methylhexadecanoate iso-Heptadecanoic methyl ester; iso C17 methyl ester C ₁₈ H ₃₆ O ₂ CAS#: 6929-04-0	20 mg
	Source: synthetic Mol. Wt.: 284 Purity: 98+% by GC Appearance: clear liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1603	Methyl 17-methyloctadecanoate iso-Nonadecanoic methyl ester; iso C19 methyl ester C ₂₀ H ₄₀ O ₂	20 mg
	Source: synthetic Mol. Wt.: 312 Purity: 98+% by GC Appearance: clear liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	

anteiso-Fatty acids and esters



1615	12-Methyltetradecanoic acid anteiso-Pentadecanoic acid; anteiso C15 acid C ₁₅ H ₃₀ O ₂ CAS#: 5502-94-3	20 mg
	Source: synthetic Mol. Wt.: 242 Purity: 98+% by GC Appearance: white solid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1612	Methyl 12-methyltetradecanoate anteiso-Pentadecanoic methyl ester; anteiso C15 methyl ester C ₁₆ H ₃₂ O ₂ CAS#: 5129-66-8	20 mg
	Source: synthetic Mol. Wt.: 256 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1613	Methyl 13-methylpentadecanoate anteiso-Palmitic methyl ester; anteiso C16 methyl ester C ₁₇ H ₃₄ O ₂ CAS#: 5487-50-3	20 mg
	Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1616	14-Methylhexadecanoic acid anteiso-Heptadecanoic acid; anteiso C17 acid C ₁₇ H ₃₄ O ₂ CAS#: 5918-29-6	20 mg
	Source: synthetic Mol. Wt.: 270 Purity: 98+% by GC Appearance: white solid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	
1614	Methyl 14-methylhexadecanoate anteiso-Heptadecanoic methyl ester; anteiso C17 methyl ester C ₁₈ H ₃₆ O ₂ CAS#: 2490-49-5	20 mg
	Source: synthetic Mol. Wt.: 284 Purity: 98+% by GC Appearance: liquid Solubility: chloroform, ethyl ether, ethanol Storage: -20°C	

Methylated fatty acids

1207	D,L-2,6-Dimethylheptanoic acid 2,6-Dimethyl C7:0 fatty acid C ₉ H ₁₈ O ₂	50 mg
	Source: synthetic Mol. Wt.: 158 Purity: 98+% by TLC, GC Appearance: clear oil Solubility: chloroform Storage: room temperature	
1791	10-Methylhexadecanoic acid 10-Methyl C16:0 fatty acid C ₁₇ H ₃₄ O ₂	25 mg
	Source: synthetic Mol. Wt.: 270 Purity: 98+% by TLC, GC Appearance: clear oil Solubility: chloroform Storage: room temperature	
1792	Methyl 10-methylhexadecanoate 10-Methyl C16:0 fatty acid methyl ester C ₁₈ H ₃₆ O ₂	25 mg
	Source: synthetic Mol. Wt.: 284 Purity: 98+% by TLC, GC Appearance: clear oil Solubility: chloroform Storage: room temperature	

1195 Phytanic acid 3,7,11,15-Tetramethylhexadecanoic acid C₂₀H₄₀O₂ CAS#: 14721-66-5 **25 mg**

Source: semi-synthetic **Mol. Wt.:** 312 **Purity:** 97+% by GC **Appearance:** white solid **Solubility:** chloroform, methanol **Storage:** -20°C

Cyclopropyl fatty acids and esters

1822 Dihydrosterculic acid cis-9,10-Methyleneoctadecanoic acid C₁₉H₃₆O₂ CAS#: 4675-61-0 **25 mg**

Source: synthetic **Mol. Wt.:** 296 **Melting Point (°C):** 38-42 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, hexane **Storage:** -20°C

1823 Methyl dihydrosterculate Methyl cis-9,10-methyleneoctadecanoate C₂₀H₃₈O₂ CAS#: 3971-54-8 **25 mg**

Source: synthetic **Mol. Wt.:** 310 **Purity:** 98+% by TLC, GC **Appearance:** oil **Solubility:** chloroform, ethanol, methanol, hexane **Storage:** -20°C

Unusual fatty acids and derivatives

1751 N-Oleylethanolamine NOE C₂₀H₃₉NO₂ CAS#: 111-58-0 **100 mg**

Source: synthetic **Mol. Wt.:** 326 **Melting Point (°C):** 63-66 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, ethyl ether, DMSO **Storage:** -20°C

Activity: acid ceramidase inhibitor

References:

C. J. Hillard and W.B. Campbell. J. Lipid. Res., **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymnowska D., Wojtczak L. BBA **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. BBRC, **255**: 456-459, 1999

1786 N-Hexadecanoylethanolamine C₁₈H₃₇NO₂ CAS# 544-31-0 **100 mg**

Source: synthetic **Mol. Wt.:** 299 **Melting Point (°C):** 99-102 **Purity:** 98+% by TLC **Appearance:** white solid **Solubility:** chloroform, ethanol, methanol, Storage: -20°C

Activity: inactive as acid ceramidase inhibitor


References:

C. J. Hillard and W.B. Campbell. J. Lipid. Res., **38**: 2383-2398, 1997
Wasilewski M., Wieckowski M.R., Dymnowska D., Wojtczak L. BBA **1657**: 151-163, 2004
Spinedi A., DiBartolomeo S., and Piacentini M. BBRC, **255**: 456-459, 1999

1757	Anandamide Arachidonylethanolamide; 5,8,11,14(Z,Z,Z,Z)-eicosatetraenoyl 2-hydroxy-ethyl-amide C ₂₂ H ₃₇ NO ₂ CAS#: 94421-68-8	10 mg/ml, 1 ml
	Source: synthetic Mol. Wt.: 347 Purity: 98+% by TLC Appearance: liquid	
	Solvent: chloroform Solubility: chloroform, ethanol Storage: -20°C	
	Induces apoptosis, endocannabinoid	
	References: Wasilewski M., Wieckowski M.R., Dymnowska D, Wojtczak L. BBA 1657 : 151-163, 2004 M. van der Stelt and V. Di Marzo; Prostaglandins Other Lipid Mediat. 77 , 2005 C. Grimaldi, et al.: Exp. Cell Res. 312 , 363, 2006	


Other lipids

Tocopherols




1072 **rac-alpha-Tocopherol** **5,7,8-Trimethyltocol** **C₂₉H₅₀O₂** **CAS#:** **59-02-9** **50 mg/ml, 1 ml**

Source: synthetic **Mol. Wt.:** 431 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C




1071 rac-beta-Tocopherol 50 mg/ml, 1 ml
5,8-Dimethyltocol C₂₈H₄₈O₂ CAS#: 148-03-8

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C




1073 rac-gamma-Tocopherol 50 mg/ml, 1 ml
7,8-Dimethyltocol C₂₈H₄₈O₂ CAS#: 73980-80-0

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 97% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C




1074 rac-5,7-Dimethyltocol
C₂₈H₄₈O₂ CAS#: 493-35-6

Source: synthetic **Mol. Wt.:** 417 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, ethyl ether, chloroform, alcohols **Storage:** -20°C



1790 (+)-delta-Tocopherol
8-Methyltocol C₂₇H₄₆O₂ CAS#: 119-13-1


Source: natural, plant **Mol. Wt.:** 403 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C



1797 Tocol
rac-Tocol C₂₆H₄₄O₂

Source: synthetic **Mol. Wt.:** 389 **Purity:** 95% by TLC, 98% by GC
Appearance: liquid **Solvent:** hexane **Solubility:** hexane, methanol, ethanol **Storage:** -20°C

Cholestane derivatives



Catalog number 1006

1006 Cholesterol
C₂₇H₄₆O CAS#: 57-88-5

Source: natural, ovine **Mol. Wt.:** 386 **Melting Point (°C):** 147-148 **Purity:** 98+% by GC **Appearance:** white solid **Solubility:** chloroform, ethanol **Storage:** -20°C

1115	5-alpha-Cholestane C ₂₇ H ₄₈ O CAS#: 481-21-0	100 mg
-------------	---	---------------

Source: synthetic **Mol. Wt.:** 373 **Purity:** 98+% by GC **Appearance:** white solid
Solubility: chloroform, ethyl ether, hexane **Storage:** -20°C

1116	Coprostanol 5-beta-Cholestane-3-beta-ol C ₂₇ H ₄₈ O CAS#: 360-68-9	25 mg
-------------	--	--------------

Source: semi-synthetic **Mol. Wt.:** 390 **Melting Point (°C):** 101-103 **Purity:** 98+% by GC **Appearance:** white solid **Solubility:** chloroform, ethyl ether, warm methanol **Storage:** -20°C

Plant sterols and steryl glucosides

1119	Plant sterol mix Sterol mixture, qualitative	25 mg/ml, 1 ml
-------------	--	-----------------------

Source: natural, plant **Appearance:** liquid **Solvent:** chloroform
Solubility: chloroform **Storage:** -20°C

Contains: Brassicasterol (13%), campesterol (26%), stigmasterol (7%), β-sitosterol (53%) in order of elution

1123	Plant sterols kit Sterols kit	1 kit
-------------	---	--------------

Source: synthetic or plant **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

Contains in individual packages: steryl glucoside 25 mg, esterified steryl glucoside 10 mg, plant sterol mixture 25 mg, β-sitosterol (55%) 100 mg, desmosterol (85%) 2 mg, lanosterol (55%) 100 mg, stigmasterol 25 mg, ergosterol 25 mg, coprostanol 5 mg, cholestanol 100 mg

1113	β-Sitostanol Stigmastanol C ₂₉ H ₅₂ O CAS#: 19466-47-8	50 mg
-------------	--	--------------

Source: synthetic **Mol. Wt.:** 417 **Melting Point (°C):** 127-132 **Purity:** 98+% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1120	Lanosterol C ₃₀ H ₅₀ O CAS#: 79-63-0	500 mg
-------------	--	---------------

Source: synthetic or plant **Mol. Wt.:** 427 **Purity:** 55% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1121	Stigmasterol 5,22-cholestadien-24-beta-ethyl-3-beta-ol C ₂₉ H ₄₈ O CAS#: 83-48-7	100 mg
-------------	--	---------------

Source: synthetic **Mol. Wt.:** 413 **Melting Point (°C):** 170 **Purity:** 95% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1122	Ergosterol C ₂₈ H ₄₄ O CAS#: 57-87-4	100 mg
-------------	--	---------------

Source: synthetic or plant **Mol. Wt.:** 397 **Melting Point (°C):** 156 **Purity:** 95% by TLC, GC **Appearance:** white solid **Solubility:** chloroform **Storage:** -20°C

1117	Steryl glucoside C ₃₅ H ₆₀ O ₆	25 mg
	Source: natural, plant Mol. Wt.: 576 Melting Point (°C): 283-287 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1118	Esterified steryl glucoside 1:1:1, sterol:glucose:fatty acid C ₅₁ H ₉₀ O ₇	10 mg
	Source: natural, plant Mol. Wt.: 814 Purity: 98+% by TLC Appearance: solid film Solubility: chloroform, ethyl ether, pyridine Storage: -20°C Sterol, glucose and fatty acid in a molar ratio 1:1:1. Mol. Wt. based on β-sitosterol glucoside palmitate.	

Propyleneglycol Monoesters

1862	2-Hydroxypropyl hexadecanoate Propyleneglycol monopalmitate C ₁₉ H ₃₈ O ₃	100 mg
	Source: synthetic Mol. Wt.: 314 Purity: 98+% by TLC, GC Appearance: white solid Solubility: hexane, ethyl ether, alcohol, chloroform, Storage: -20°C	
1863	2-Hydroxypropyl octadecanoate Propyleneglycol monostearate C ₂₁ H ₄₂ O ₃	100 mg
	Source: synthetic Mol. Wt.: 342 Purity: 98+% by TLC, GC Appearance: white solid Solubility: hexane, ethyl ether, alcohol, chloroform, Storage: -20°C	

Standards and reference compounds

Food industry mixes

Each methyl ester mix is carefully prepared by weight.

4210	KEL-FIM-FAME-5 mix Methyl ester mix	15.5 mg/ml 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: heptane Solubility: heptane Storage: -20°C	
	Contains the methyl esters of the following fatty acids (mg/ml in brackets): C8:0 (0.3), C10:0 (0.5), C12:0 (1.0), C13:0 (0.5), C14:0 (0.5), C14:1 (0.3), C15:0 (0.3), C16:0 (2.0), C16:1 (1.0), C17:0 (0.5), C18:0 (1.0), C18:1tr (0.4), C18:1c (3.0), C18:2 (2.0), C20:0 (0.3), C18:3 (1.0), C20:1 (0.3), C22:0 (0.3), C22:1 (0.3), listed in order of their elution.	
2009	FIM-FAME-6 mix Methyl ester mix	33 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: heptane Solubility: Storage: -20°C	
	Contains the methyl esters of these fatty acids. Each methyl ester is 3.03% of the mixture except C16:0 which is 6.06%. C4:0 , C6:0 , C8:0, C10:0 , C11:0, C12:0, C13:0 , C14:0, C14:1(cis-9), C15:0, C15:1(cis-10), C16:0, C16:1(cis-9), C17:0, C17:1(cis-10), C18:0, C18:1(trans-9), C18:1(cis-9), C18:2(all-cis-9,12), C20:0, C18:3(all-cis 6,9,12), C20:1(cis-11), C18:3(all-cis 9,12,15), C20:2(all-cis 11,14), C22:0, C20:3(all-cis 8,11,14), C22:1(cis 13), C20:3(all-cis 11,14,17), C20:4(all-cis 5,8,11,14), C22:2(all-cis 13,16), C24:1(cis-15), C22:6(all-cis 4,7,10,13,16,19), listed in order of their elution.	

2010	FIM-FAME-7 mix Methyl ester mix	30 mg/ml, 1 ml
-------------	---	-----------------------

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Contains the methyl esters of these fatty acids (weight percent in [brackets]): C4:0 [4.0], C6:0 [4.0], C8:0 [4.0], C10:0 [4.0], C11:0 [2.0], C12:0 [4.0], C13:0 [2.0], C14:0 [4.0], C14:1(cis-9) [2.0], C15:0 [2.0], C15:1(cis-10) [2.0], C16:0 [6.0], C16:1(cis-9) [2.0], C17:0 [2.0], C17:1(cis-10) [2.0], C18:0 [4.0], C18:1(trans-9) [2.0], C18:1(cis-9) [4.0], C18:2(all-trans-9,12) [2.0], C18:2(all-cis-9,12) [2.0], C18:3(all-cis 6,9,12) [2.0], C20:0 [4.0], C20:1(cis-11) [2.0], C18:3(all-cis 9,12,15) [2.0], C21:0 [2.0], C20:2(all-cis 11,14) [2.0], C20:3 (all-cis 8,11,14) [2.0], C22:0 [4.0], C22:1(cis-13) [2.0], C20:3(all-cis 11,14,17) [2.0], C20:4(all-cis 5,8,11,14) [2.0], C23:0 [2.0], C22:2(all-cis 13,16) [2.0], C20:5(all-cis 5,8,11,14,17) [2.0], C24:0 [4.0], C24:1(cis-15) [2.0], 22:6(all-cis 4,7,10,13,16,19) [2.0}, listed in order of their elution.

Polyunsaturated fatty acid methyl esters mixes

These are complex qualitative standard mixtures of polyunsaturated fatty acid methyl esters. Because they are extracted from natural materials, relative peak sizes may vary from lot to lot.

1093	PUFA-1 Qualitative mix	100 mg
-------------	----------------------------------	---------------

Source: natural, fish oil **Appearance:** yellow oil **Solubility:** chloroform, ethanol, hexane, methanol **Storage:** -20°C

Contains: C14:0, C16:0, C16:1ω7, C18:1ω9, C18:1ω7, C18:2ω6, C18:4ω3, C20:1ω9, C20:2ω6, C20:5ω3, C22:1ω11, C22:1ω9, C22:5ω3, C22:6ω3

1081	PUFA-2 Qualitative mix	100 mg
-------------	----------------------------------	---------------

Source: natural, porcine **Appearance:** yellow oil **Solubility:** alcohols, hexane, chloroform **Storage:** -20°C

Contains: C14:0, C16:0, C16:1ω7, C18:0, C18:1ω9, C18:1ω7, C18:2ω6, C18:3ω6, C18:3ω3, C20:1ω9, C20:2ω6, C20:3ω6, C20:4ω6, C20:5ω3, C22:4ω6, C22:5ω3, C22:6ω3

1177	PUFA-3 Qualitative mix	100 mg
-------------	----------------------------------	---------------

Source: natural, menhaden oil **Appearance:** yellow oil **Solubility:** alcohols, hexane, chloroform **Storage:** -20°C

Contains: C14:0, C16:0, C16:1ω7, C16:2ω4, C16:3ω4, C16:4ω1, C18:0, C18:1ω9, C18:1ω7, C18:2ω6, C18:2ω4, C18:3ω4, C18:3ω3, C18:4ω3, C20:1ω9, C20:4ω6, C20:4ω3, C20:5ω3, C21:5ω3, C22:5ω3, C22:6ω3

Carbohydrate mixes

1124	Alditol acetate mix-1 Quantitative carbohydrate mix	50 mg/ml, 1 ml
-------------	---	-----------------------

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform **Storage:** -20°C

Contains: rhamnitol, fucitol, ribitol and arabinitol pentaacetates, 12.5 mg/ml each

1125 Alditol acetate mix-2 **50 mg/ml, 1 ml**
Quantitative carbohydrate mix

Source: synthetic **Appearance:** liquid **Solvent:** chloroform **Solubility:** chloroform
Storage: -20°C

Contains: mannitol, galactitol, glucitol and inositol hexaacetates, 12.5 mg/ml each

Other fatty acid methyl ester mixes

1722 2-Hydroxy methyl ester mix **10 mg/ml, 1 ml**
Source: synthetic **Appearance:** liquid **Solvent:** methylene chloride **Solubility:** methylene chloride **Storage:** -20°C

Quantitative mix contains: C14:0, 20%; C16:0, 20%; C18:0, 15%; C20:0, 15%; C22:0, 10%; C23:0, 10%; C24:0, 10%

1131 Cis-trans isomer standard **5 mg/ml, 5 ml**

Source: margarine **Appearance:** liquid **Solvent:** 5ml methylene chloride
Solubility: methylene chloride **Storage:** -20°C

Analysis of positional cis-trans fatty acid isomers is ever more important in light of the new food industry rules. These isomers can be resolved on Supelco SP-2560 or an equivalent capillary GC column. Use this specially formulated mix to ensure proper operation of your column for this tricky separation. Mix consists of cis-trans fatty acid isomers as methyl esters in methylene chloride.

This is a qualitative standard containing in order of elution: C16:0, C18:0, C18:1 trans isomers (4 peaks), C18:1 cis & trans isomers (2 peaks), C18:1 cis isomers (4 peaks), C18:2, C20:0, C20:1 and C18:3 (same peak), C22:0

AOCS animal and vegetable oil reference mixes (RM mixes)

By studying problems with the quantitative analysis of animal and vegetable oils and fats, the American Oil Chemists' Society has found certain mixtures to be useful as reference standards. The composition of each mixture (see Table I below) is similar to the fatty acid distribution of certain oils. All mixtures are in methyl ester form and ready for GC analysis

Table I. AOCS Oil Reference Mixes

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated in the Table.

Mix No. Catalog No.	RM-1 1084	RM-2 1085	RM-3 1086	Rapeseed 1083	RM-4 1087	RM-5 1088	RM-6 1089
C8:0 Caprylate						7.0	
C10:0 Caprate						5.0	
C12:0 Laurate						48.0	
C14:0 Myristate			1.0	1.0		15.0	2.0
C16:0 Palmitate	6.0	7.0	4.0	4.0	11.0	7.0	30.0
C16:1 Palmitoleate							3.0
C18:0 Stearate	3.0	5.0	3.0	3.0	3.0	3.0	14.0
C18:1 Oleate	35.0	18.0	45.0	60.0	80.0	12.0	41.0
C18:2 Linoleate	50.0	36.0	15.0	12.0	6.0	3.0	7.0
C18:3 Linolenate	3.0	34.0	3.0	5.0			3.0
C20:0 Arachidate	3.0		3.0	3.0			
C20:1 Eicosenoate				1.0			
C22:0 Behenate			3.0	3.0			
C22:1 Erucate			20.0	5.0			
C24:0 Lignocerate			3.0	3.0			

1083 Rapeseed oil reference mixture 25 mg/ml, 1 ml

AOCS rapeseed oil reference mix

Source: synthetic or plant **Appearance:** liquid **Solvent:** methylene chloride
Solubility: ethyl ether, methylene chloride **Storage:** -20°C

Suitable standard for low erucic acid oil

1084 RM-1 mix 50 mg

AOCS reference mix RM-1


Source: synthetic or plant **Appearance:** clear oil **Solubility:** chloroform, ethyl ether
Storage: -20°C

Suitable standard for corn, cottonseed, soybean, safflower, sesame, poppy seed, walnut kapok, and rice oils

1085	RM-2 mix AOCS reference mix RM-2	50 mg
	Source: synthetic or plant Appearance: clear oil Solubility: chloroform, ethanol, ethyl ether Storage: -20°C	
	Suitable standard for linseed, perilla, hempseed, and rubberseed oils	
1086	RM-3 mix AOCS reference mix RM-3	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: ethyl ether, methylene chloride Storage: -20°C	
	Suitable standards for peanut, rapeseed, and mustard seed oils	
1087	RM-4 mix AOCS reference mix RM-4	50 mg
	Source: synthetic or plant Appearance: clear oil Solubility: chloroform, ethyl ether Storage: -20°C	
	Suitable standard for olive, teased, and neatsfoot oils	
1088	RM-5 mix AOCS reference mix RM-5	50 mg
	Source: synthetic or plant Appearance: clear oil Solubility: chloroform Storage: -20°C	
	Suitable standard for coconut, palm kernel, babassu and ouri-ouri oils	
1089	RM-6 mix AOCS reference mix RM-6	50 mg
	Source: synthetic or plant Appearance: clear oil Solubility: ethyl ether, methylene chloride Storage: -20°C	
	Suitable standard for lard, beef tallow, mutton tallow, and palm oil	
1082	RM-7 kit AOCS reference mix RM-7 kit	50 mg ampules
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
	50 mg ampules of RM-1, RM-2, RM-3, RM-4, RM-5, RM-6, and 25 mg of Rapeseed oil reference mixture	

Custom mixes

Custom fatty acid methyl ester mixes can be prepared to your specification. Minimum quantity requirements apply to these orders.



Cat# 4210 spiked with 0.4 mg/ml C18:2t ester (methyl linoelaidate) and chromatographed on a Supelco SP 2330 fused silica column.

Peak number	FAME
1	C8:0
2	C10:0
3	C12:0
4	C13:0
5	C14:0
6	C14:1
7	C15:0
8	C16:0
9	C16:1
10	C17:0
11	C18:0
12	C18:1t-9
13	C18:1c-9
14	C18:2t,t-9,12
15	C18:2c,c-9,12
16	C20:0
17	C18:3
18	C20:1
19	C22:0
20	C22:1

Table II. Standards for GC analysis

GLC Standard mixes

GLC-10 through GLC-100 standards are **equal weight measures** of fatty acid methyl esters. They are quantitative standards, useful for determining relative **retention times and response factors**.

Each methyl ester mixture is carefully prepared by weight and the composition verified by gas chromatography. The weight percentage of each component is indicated below. All double bonds are cis.

Mixture Number Catalog Number	GLC-10 1095	GLC-30 1097	GLC-40 1098	GLC-50 1099	GLC-60 1100	GLC-70 1101	GLC-80 1102	GLC-90 1103	GLC-100 1104
C8:0 Caprylate		20.0				20.0			
C9:0 Nonanoate						20.0			
C10:0 Caprate		20.0				20.0			
C11:0 Undecanoate						20.0			
C12:0 Laurate		20.0				20.0			
C13:0 Tridecanoate							20.0	20.0	
C14:0 Myristate		20.0					20.0		
C15:0 Pentadecanoate							20.0	20.0	
C16:0 Palmitate	20.0	20.0	25.0				20.0		
C16:1 Palmitoleate-9 cis				20.0					
C17:0 Heptadecanoate							20.0	20.0	
C18:0 Stearate	20.0		25.0						20.0
C18:1 Oleate-9 cis	20.0			20.0					
C18:2 Linoleate-9,12 all cis	20.0								
C18:3 Linolenate-9,12,15 all cis	20.0								
C19:0 Nonadecanoate								20.0	20.0
C20:0 Arachidate			25.0		25.0				20.0
C20:1 Eicosenoate-11 cis				20.0	25.0				
C20:2 Eicosadienoate-11,14 all cis					25.0				
C20:3 Eicosatrienoate-11,14,17 all cis					25.0				
C21:0 Heneicosanoate								20.0	20.0
C22:0 Behenate			25.0						20.0
C22:1 Erucate-13				20.0					
C24:1 Nervonate				20.0					

1095 GLC-10 mix Quantitative GC mix **50 mg**

Source: synthetic or plant **Appearance:** clear oil **Solubility:** methylene chloride
Storage: -20°C

1097 GLC-30 mix 50 mg
Quantitative GC mix

Source: synthetic or plant **Appearance:** clear oil **Solubility:** methylene chloride
Storage: -20°C

1098	GLC-40 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
1099	GLC-50 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
1100	GLC-60 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
1101	GLC-70 mix Quantitative GC mix	50 mg
	Source: synthetic or plant Appearance: clear oil Solubility: methylene chloride Storage: -20°C	
1102	GLC-80 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
1103	GLC-90 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	
1104	GLC-100 mix Quantitative GC mix	50 mg/ml, 1 ml
	Source: synthetic or plant Appearance: liquid Solvent: methylene chloride Solubility: methylene chloride Storage: -20°C	

Water soluble fatty acid mixes

1106	WSFA-2 mix Water soluble fatty acid qualitative mix	5 ml
	Appearance: liquid Solvent: water Solubility: water Storage: Room Temp Contains: acetic, propionic, isobutyric, n-butyric, isovaleric and n-valeric acids	
1108	WSFA-4 mix Water soluble fatty acid qualitative mix	5 ml
	Appearance: liquid Solvent: water Solubility: water Storage: Room Temp Contains: acetic, propionic, isobutyric, n-butyric, 2-methyl butyric, isovaleric and n-valeric acids	

Microbiology standard mixes

1105	GLC-110 mix Bacterial lipid standard, qualitative mix	10 mg/ml, 1 ml
-------------	---	-----------------------

Source: various **Appearance:** liquid **Solvent:** chloroform **Solubility:** methylene chloride, choroform **Storage:** -20°C

Contains:

methyl 12-methyltridecanoate	(iso-C14:0)	methyl 14-methylpentadecanoate	(iso-C16:0)
methyl myristate	(C14:0)	methyl palmitate	(C16:0)
methyl 12-methyltetradecanoate	(anteiso-C15:0)	methyl 14-methylhexadecanoate	(anteiso-C17:0)
methyl pentadecanoate	(C15:0)		

1114	Bacterial acid methyl esters CP mix Qualitative mix	10 mg/ml, 1 ml
-------------	---	-----------------------

Source: various **Appearance:** liquid **Solvent:** methyl caproate **Solubility:** hexane, ethanol, methanol **Storage:** -20°C

A qualitative standard. Mixture consists of equal amounts of the compounds listed.

methyl undecanoate	C11:0	methyl cis-9-hexadecenoate	C16:1 ⁹
methyl 2-hydroxydecanoate	2-OH C10:0	methyl hexadecanoate	C16:0
methyl dodecanoate	C12:0	methyl 15-methylhexadecanoate	iso-C17:0
methyl tridecanoate	C13:0	methyl cis-9,10-methylenehexadecanoate	C17:0Δ ^{9,10}
methyl 2-hydroxydodecanoate	2-OH C12:0	methyl heptadecanoate	C17:0
methyl 3-hydroxydodecanoate	3-OH C12:0	methyl 2-hydroxyhexadecanoate	2-OH C16:0
methyl tetradecanoate	C14:0	methyl cis, cis-9,12-octadecadienoate	C18:2 ^{9,12}
methyl 13-methyltetradecanoate	iso-C15:0	methyl cis-9-octadecenoate	C18:1 ⁹
methyl 12-methyltetradecanoate	anteiso-C15:0	methyl trans-9-octadecenoate	C18:1 ⁹
methyl pentadecanoate	C15:0	methyl octadecanoate	C18:0
methyl 2-hydroxytetradecanoate	2-OH C14:0	methyl cis-9,10-methyleneoctadecanoate	C19:0Δ ^{9,10}
methyl 3-hydroxytetradecanoate	3-OH C14:0	methyl nonadecanoate	C19:0
methyl 14-methylpentadecanoate	iso-C16:0	methyl eicosanoate	C20:0

1075	Volatile acid mix Qualitative mix	100 ml
-------------	---	---------------

Appearance: liquid **Solvent:** water **Solubility:** water
Storage: 4-8°C

Contains: formic, acetic, propionic, isobutyric, n-butyric, isovaleric, n-valeric, isocaproic, n-caproic, and heptanoic acids

1077	Non-volatile acid mix Qualitative mix	100 ml
-------------	---	---------------

Appearance: liquid **Solvent:** water **Solubility:** water
Storage: 4-8°C

Contains: pyruvic, lactic, oxalacetic, oxalic, methyl malonic, malonic, fumaric and succinic acids.

Biochemical research standard mixes

These mixtures are prepared by precise gravimetric technique. All mixtures contain equal amounts of listed components. A data sheet is supplied with each mixture.

1127	Polar lipid mix TLC standards mix	25 mg/ml, 1 ml
-------------	---	-----------------------

Source: natural, egg, ovine **Appearance:** liquid **Solvent:** chloroform/methanol 2:1
Solubility: chloroform/methanol 2:1 **Storage:** -20°C

Contains: cholesterol, phosphatidylethanolamine, lecithin, and lyso-lecithin

1128	Sphingolipid mix TLC standards mix	25 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol 2:1 Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Contains: cerebrosides, sulfatides, and sphingomyelin	
1129	Non-polar lipid mix A TLC standards mix	25 mg/ml, 1 ml
	Source: natural, plant, ovine Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	Contains: cholesteryl palmitate, tripalmitin, palmitic acid, and cholesterol	
1130	Non-polar lipid mix B TLC standards mix	25 mg/ml, 1 ml
	Source: natural, plant, ovine Appearance: liquid Solvent: chloroform Solubility: chloroform Storage: -20°C	
	Contains: cholesteryl oleate, methyl oleate, triolein, oleic acid, and cholesterol	

Glycosphingolipid mixtures for TLC

These mixtures are qualitative standards prepared from our purified glycosphingolipids.

1505	Neutral glycosphingolipid qualmix, Glycosylceramides, qualitative mix	1 mg/ml, 1 ml
	Source: natural, bovine and porcine Appearance: liquid Solvent: chloroform/methanol 2:1 Solubility: chloroform/methanol 2:1 Storage: -20°C	
	Contains: cerebrosides, lactosylceramide, ceramide trihexoside, globoside	
1508	Monosialoganglioside mix GM ₃ , GM ₂ , GM ₁ qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: GM ₃ , GM ₂ , GM ₁	
1509	Disialoganglioside mix GD ₃ , GD _{1a} , GD _{1b} , qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: GD ₃ , GD _{1a} , GD _{1b}	
1510	Lactosylceramide and sialosyl derivatives mix LC, GM ₃ , GD ₃ qualitative mix	0.5 mg/ml, 1 ml
	Source: natural, bovine Appearance: liquid Solvent: chloroform/methanol/water 2:1:0.1 Solubility: chloroform/methanol/water 2:1:0.1 Storage: -20°C	
	Contains: LC, GM ₃ , GD ₃	

1511	Gangliotetraosylceramide and sialosyl derivatives mix asialo-GM ₁ , GM ₁ , GD _{1a} , GD _{1b} , GT _{1b} qualitative mix	0.5 mg/ml, 1 ml
------	---	-----------------

Source: natural, bovine **Appearance:** liquid **Solvent:** chloroform/methanol/water 2:1:0.1 **Solubility:** chloroform/methanol/water 2:1:0.1 **Storage:** -20°C

Contains: asialo-GM₁, GM₁, GD_{1a}, GD_{1b}, GT_{1b}


Biochemicals and reagents

Stable isotope labeled compounds

1914	N-Stearoyl-D₃₅-psychosine, perdeuterated Cerebrosides with N-C18:0-D ₃₅ fatty acid side chain C ₄₂ H ₄₆ D ₃₅ NO ₈	5 mg
	Source: semi-synthetic, bovine Mol. Wt.: 762 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform, hot ethanol, chloroform/methanol 2:1 Storage: -20°C	
1533	N-Palmitoyl-D₃-glucopsychosine, deuterated N-C16:0-D ₃ -Glucopsychosine; glucocerebroside with C16:0-D ₃ fatty acid side chain C ₄₀ H ₇₄ D ₃ NO ₈	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 703 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1534	N-Palmitoyl-D₃-lactosylceramide, deuterated N-C16:0-D ₃ -Lactosylceramide; lactosylceramide with C16:0-D ₃ fatty acid side chain C ₄₆ H ₈₄ D ₃ NO ₁₃	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 864 Purity: 98+% by TLC Appearance: white solid Solubility: chloroform/methanol/water 5:1:0.1 Storage: -20°C	
2200	N-1-¹³C-Palmitoyl-sphingosylphosphorylcholine D-erythro-Sphingomyelin with 1- ¹³ C-palmitic acid; SPM with ¹³ C labeled fatty acid C ₃₈ ¹² C ₃₈ ¹³ CH ₇₉ N ₂ O ₆ P	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 703 Purity: 98+% by TLC Appearance: waxy solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
2050	N-Octadecanoyl-D₃-monosialoganglioside GM₁ N-D ₃ -Stearoyl-GM ₁ C ₇₃ H ₁₂₈ N ₃ O ₃₁ D ₃	0.5 mg
	Source: semi-synthetic, bovine brain Mol. Wt.: 1548 Purity: 98+% by TLC Appearance: solid, Solubility: chloroform/methanol/water 2:1:0.1, forms micellar solution in water Storage: -20°C	
1536	N-Octadecanoyl-D₃-sulfatide N-C18:0-D ₃ -Sulfatide C ₄₂ H ₇₈ D ₃ NO ₁₁ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 833 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol/DI water 2:1:0.1 Storage: -20°C	

1537	N-Octadecanoyl-D₃-ceramide trihexoside C18:0-D ₃ -CTH; C18:0-D ₃ -Gb3; N-Octadecanoyl-D ₃ -globotriaosylceramide C ₅₄ H ₉₈ D ₃ NO ₁₈	0.5 mg
Source: semi-synthetic, porcine Mol. Wt.: 1055 Purity: 98+% by TLC Appearance: off-white solid Solubility: chloroform/methanol 2:1; DMSO Storage: -20°C		

Fluorescent compounds



1841	N-Hexanoyl-NBD-D-erythro-sphingosine	100 µg
1841-001	N-C6:0-NBD-ceramide; N-C6:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-sphingosine C ₃₀ H ₄₉ N ₅ O ₆ CAS#: 86701-10-2	1 mg

Source: synthetic **Mol. Wt.:** 575 **Melting Point (°C):** 85-88 **Purity:** 98+% by TLC
Appearance: orange film, vacuum dried **Solubility:** chloroform, ethanol, methanol
Storage: -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1618	N-Dodecanoyl-NBD-D-erythro-sphingosine	100 µg
1618-001	N-C12:0-NBD-ceramide; N-C12:0-NBD-D-erythro-sphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-sphingosine C ₃₆ H ₆₁ N ₅ O ₆	1 mg

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1857	N-Hexanoyl-NBD-L-threo-sphingosine	100 µg
1857-001	N-C6:0-NBD-ceramide; N-C6:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminocaproyl)-L-threo-sphingosine C ₃₀ H ₄₉ N ₅ O ₆	1 mg

Source: synthetic **Mol. Wt.:** 575 **Purity:** 98+% by TLC **Appearance:** red-orange solid
Solubility: chloroform, ethanol, methanol **Storage:** -20°C

Reference:
J. M. L. Hauser et al., J. Biol. Chem. **269**, 6803, 1994

1620	N-Dodecanoyl-NBD-L-threo-sphingosine	100 µg
1620-001	N-C12:0-NBD-ceramide; N-C12:0-NBD-L-threo-sphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-sphingosine C ₃₆ H ₆₁ N ₅ O ₆	1 mg

Source: synthetic **Mol. Wt.:** 660 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1624	N-Hexanoyl-NBD-L-threo-dihydroosphingosine	100 µg
1624-001	N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-L-threo-dihydroosphingosine, fluorescent; N-(NBD-aminocaproyl)-L-threo-dihydroosphingosine C ₃₀ H ₅₁ N ₅ O ₆	1 mg

Source: synthetic **Mol. Wt.:** 578 **Purity:** 98+% by TLC **Appearance:** orange solid
Solubility: chloroform/methanol 2:1, methanol **Storage:** -20°C

1623	N-Dodecanoyl-NBD-L-threo-dihydrosphingosine	100 µg
1623-001	N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-L-threo-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-L-threo-dihydrosphingosine <chem>C36H63N5O6</chem>	1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1626	N-Hexanoyl-NBD-D-erythro-dihydrosphingosine	100 µg
1626-001	N-C6:0-NBD-dihydroceramide; N-C6:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminocaproyl)-D-erythro-dihydrosphingosine <chem>C30H51N5O6</chem>	1 mg
	Source: synthetic Mol. Wt.: 578 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1625	N-Dodecanoyl-NBD-D-erythro-dihydrosphingosine	100 µg
1625-001	N-C12:0-NBD-dihydroceramide; N-C12:0-NBD-D-erythro-dihydrosphingosine, fluorescent; N-(NBD-aminolauroyl)-D-erythro-dihydrosphingosine <chem>C36H63N5O6</chem>	1 mg
	Source: synthetic Mol. Wt.: 662 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1628	N-Hexanoyl-NBD-phytosphingosine	100 µg
1628-001	N-C6:0-NBD-phytoceramide; N-C6:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminocaproyl)-phytosphingosine <chem>C30H51N5O7</chem>	1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 594 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1627	N-Dodecanoyl-NBD-phytosphingosine	100 µg
1627-001	N-C12:0-NBD-phytoceramide; N-C12:0-NBD-phytosphingosine, fluorescent; N-(NBD-aminolauroyl)-phytosphingosine <chem>C36H63N5O7</chem>	1 mg
	Source: semi-synthetic, bacteria Mol. Wt.: 678 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
1912	N-Hexanoyl-NBD-sphingosylphosphorylcholine	100 µg
1912-001	N-C6:0-NBD-sphingomyelin, fluorescent; N-C6:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminocaproyl)-sphingomyelin <chem>C35H61N6O9P</chem> CAS#: 94885-04-8	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 740 Purity: 98+% by TLC Appearance: red-brown solid Solubility: chloroform, ethanol, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	
1619	N-Dodecanoyl-NBD-sphingosylphosphorylcholine	100 ug
1619-001	N-C12:0-NBD-sphingomyelin, fluorescent; N-C12:0-NBD-sphingosylphosphorylcholine; fluorescent sphingomyelin; N-(NBD-aminolauroyl)-sphingomyelin <chem>C41H73N6O4P</chem>	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 825 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1, methanol Storage: -20°C	
	Mixture of D-erythro and L-threo isomers	

1621	N-Hexanoyl-NBD-galactosylceramide	100 µg
1621-001	N-C6:0-NBD-beta-D-galactosylsphingosine; N-C6:0-NBD-cerebrosides; N-C6:0-NBD-galactosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-galactosylsphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1622	N-Hexanoyl-NBD-glucosylceramide	100 µg
1622-001	N-C6:0-NBD-beta-D-glucosylsphingosine; N-C6:0-NBD-glucosylceramide, fluorescent; N-(NBD-aminocaproyl)-beta-D-glucosylsphingosine C ₃₆ H ₅₉ N ₅ O ₁₁	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 738 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 5:1, methanol Storage: -20°C	
1629	N-Hexanoyl-NBD-lactosylceramide	50 µg
1629-001	N-Hexanoyl-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-beta-D-lactosylsphingosine; N-C6:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminocaproyl)-beta-D-lactosylsphingosine C ₄₂ H ₆₉ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 900 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1630	N-Dodecanoyl-NBD-lactosylceramide	50 µg
1630-001	N-Dodecanoyl-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-beta-D-lactosylsphingosine; N-C12:0-NBD-lactosylceramide, fluorescent; fluorescent LC; N-(NBD-aminolauroyl)-beta-D-lactosylsphingosine C ₄₈ H ₈₁ N ₅ O ₁₆	1 mg
	Source: semi-synthetic, bovine buttermilk Mol. Wt.: 984 Purity: 98+% by TLC Appearance: orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	
1631	N-Dodecanoyl-NBD-ceramide trihexoside	100 µg
1631-001	N-C12:0-NBD-CTH; N-C12:0-NBD-globotriaosylceramide; N-(NBD-aminolauroyl)-ceramide trihexoside C ₅₄ H ₉₁ N ₅ O ₂₁	1 mg
	Source: semi-synthetic, porcine Mol. Wt.: 1145 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1: DMSO; hot methanol Storage: -20°C	
1632	N-Dodecanoyl-NBD-sulfatide	100 µg
1632-001	N-C12:0-NBD-sulfatide; N-Dodecanoyl-NBD-lyso-sulfatide; N-Dodecanoyl-NBD-sphingosyl-beta-D-galactoside-3-sulfate; N-(NBD-aminolauroyl)-sulfatide C ₄₂ H ₇₁ N ₅ O ₁₄ S	1 mg
	Source: semi-synthetic, bovine Mol. Wt.: 901 Purity: 98+% by TLC Appearance: red-orange solid Solubility: chloroform/methanol 2:1 Storage: -20°C	

Appendix

Table III. Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC.
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat. # 1044 Lecithin (egg)	Cat. # 1070 Lecithin (bovine)	Cat. # 1302 Lecithin (plant)	Cat. # 1045 Phosphatidyl- ethanolamine (egg)	Cat. # 1301 Phosphatidyl- ethanolamine (plant)	Cat. # 1046 lyso-Lecithin (egg)	Cat. # 1047 Phosphatidylserine (bovine)	Cat. # 1048 Phosphatidylinositol (plant)
Fatty Acids								
C14:0		trace						
C16:0	31	35	14	19	22	72	1	36
C16:1		1		1				
C18:0	16	14	4	26	3	24	42	
C18:1	31	33	11	23	7	3	27	7
C18:2	16		66	14	60			50
C18:3			6		8			7
C20:0							1	
C20:1							4	
C20:4				10			4	
C21:0								
C22:0							1	
C22:1							1	
C22:6							7	
C23:0								
C24:0								
C24:1								
C25:0								
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH								
C16:0 2-OH								
C18:0 2-OH								
C20:0 2-OH								
C22:0 2-OH								
C23:0 2-OH								
C24:0 2-OH								
C24:1 2-OH								
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	6	17	0	7	0	1	12	0
Total	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
 (actual composition may vary according to dietary history and growth condition of the source)

		Cat. # 1051 Sphingomyelin (bovine)	Cat. # 1328 Sphingomyelin (porcine RBC)	Cat. # 1053 Phosphatidic acid (semi-synthetic)	Cat. # 1057 Glucocerebroside (human)	Cat. # 1058 Monogalactosy- diglycerides (plant)	Cat. # 1059 Digalactosyldiglyceride (plant)	Cat. # 1061 Monosialoganglioside GM ₁	Cat. # 1062 Disialoganglioside GD _{1a}
Fatty Acids									
C14:0								trace	trace
C16:0	3	30	39	14	23	9	1	1	
C16:1									
C18:0	45	8	12	3	77	91	86	86	
C18:1		2	34				3	3	
C18:2			15						
C18:3									
C20:0	1	2		3			4	4	
C20:1									
C20:4									
C21:0									
C22:0	4	10		24			2	2	
C22:1									
C22:6									
C23:0		1		9			1	1	
C24:0	8	30		33			1	1	
C24:1	31	14		13			2	2	
C25:0									
C25:1									
C26:0		2							
C26:1									
C27:0									
C27:1									
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH									
C20:0 2-OH									
C22:0 2-OH									
C23:0 2-OH									
C24:0 2-OH									
C24:1 2-OH									
C25:0 2-OH									
C25:1 2-OH									
C26:0 2-OH									
C26:1 2-OH									
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	8	1	0	1	0	0	0	0	
Total	100	100	100	100	100	100	100	100	

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1063 Trisialoganglioside GT _{1b}	Cat: # 1064 Gangliotetraosylceramide	Cat: # 1065 Purified mixed gangliosides	Cat: # 1501 Disialoganglioside GD _{1b}	Cat: # 1050 Cerebrosides (bovine)	Cat: # 1066 Cerebrosides Kerasin (bovine)	Cat: # 1138 Cerebrosides Phrenosin (bovine)	Cat. # 1049 Sulfatides (bovine)
Fatty Acids								
C14:0	trace	trace	trace	trace				
C16:0	1	1	1	1	trace	trace		trace
C16:1								
C18:0	86	86	86	86	4	5		5
C18:1	3	3	3	3				trace
C18:2								
C18:3								
C20:0	4	4	4	4	1	1		1
C20:1								
C20:4								
C21:0								
C22:0	2	2	2	2	4	9		7
C22:1						trace		trace
C22:6								
C23:0	1	1	1	1	2	5		
C24:0	1	1	1	1	10	25		18
C24:1	2	2	2	2	15	43		29
C25:0					3	3		2
C25:1					1	3		2
C26:0					2	2		1
C26:1					1	4		3
C27:0					2			1
C27:1					2			
C14:0 2-OH								
C16:0 2-OH								
C18:0 2-OH					15		36	5
C20:0 2-OH					1		1	trace
C22:0 2-OH					6		8	3
C23:0 2-OH					5		6	
C24:0 2-OH					17		25	10
C24:1 2-OH					6		9	6
C25:0 2-OH					3		4	2
C25:1 2-OH							2	
C26:0 2-OH							2	
C26:1 2-OH							2	
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	0	0	0	0	0	0	5	5
Total	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1067 Ceramide trihexoside (porcine)	Cat # 1068 Globosides (porcine)	Cat: # 1118 Esterified steryl glucoside	Cat: # 1056 Ceramides (bovine)	Cat # 1322 Ceramides (non-hydroxy)	Cat # 1323 Ceramides (hydroxy)	Cat # 1535 Monosialoganglioside GM ₄	Cat: # 1332 Sphingomyelin, (egg, chicken)	Cat: # 1516 Tetrasialoganglioside GQ1 _b
Fatty Acids									
C14:0								trace	
C16:0	3	2	34	trace			4	72	5
C16:1									1
C18:0	2	1	8	4	11		2	8	80
C18:1	2		8					3	2
C18:2			36						3
C18:3			4						
C20:0	2	1	1	1	2		trace	2	4
C20:1							trace		
C20:4									
C21:0									
C22:0	17	15	4	4	10		3	5	2
C22:1							4		
C22:6									
C23:0	1	1	2	2	6		4	1	
C24:0	29	23	2	10	24		6	2	
C24:1	5	15		15	31		4	4	
C25:0				9	3				
C25:1				1	3				
C26:0		1		2	2				
C26:1		1		1	3				
C27:0				2					
C27:1				2					
C14:0 2-OH									
C16:0 2-OH									
C18:0 2-OH		trace		15		24	1		
C20:0 2-OH		trace		1		1	3		
C22:0 2-OH	3	3		6		8	25		
C23:0 2-OH	1			5		6	17		
C24:0 2-OH	19	11		17		35	18		
C24:1 2-OH	10	23				17	7		
C25:0 2-OH				3		4			
C25:1 2-OH									
C26:0 2-OH									
C26:1 2-OH						2			
C16 cis 9,10 methylene									
C18 cis 9,10 methylene									
Others	6	3	1	0	5	3	2	3	3
Total	100	100	100	100	100	100	100	100	100

Typical Fatty Acid Composition of Natural Lipids Made by Matreya LLC (continued)
 (actual composition may vary according to dietary history and growth condition of the source)

	Cat: # 1329 Sphingomyelin (buttermilk)	Cat: # 1500 Lactosyl ceramide (porcine)	Cat: # 1507 Lactosyl ceramide (buttermilk)	Cat: # 1502 Monosialoganglioside GM2	Cat: # 1503 Monosialoganglioside GM3 (buttermilk)	Cat: # 1504 Disialoganglioside GD3 (buttermilk)	Cat: # 1521 Glucocerebroside (buttermilk)	Cat: # 1522 Glucocerebrosides (plant)
Fatty Acids								
C14:0	1						trace	
C16:0	21	14	8	2	6	8	15	
C16:1								
C18:0	3	6	3	82	1	1	3	
C18:1		4	trace					
C18:2								
C18:3								
C20:0	1	1	2	7	1	1	2	
C20:1								
C20:4								
C21:0					1	2		
C22:0	23	9	28	4	23	24	31	
C22:1								
C22:6								
C23:0	29	1	33	trace	36	35	28	
C24:0	20	15	20	1	22	21	17	
C24:1	1	5	3	2	3	3		
C25:0								
C25:1								
C26:0								
C26:1								
C27:0								
C27:1								
C14:0 2-OH							trace	
C16:0 2-OH							79	
C18:0 2-OH		trace					trace	
C20:0 2-OH								
C22:0 2-OH		8					8	
C23:0 2-OH							1	
C24:0 2-OH		24					9	
C24:1 2-OH		13						
C25:0 2-OH								
C25:1 2-OH								
C26:0 2-OH								
C26:1 2-OH								
C16 cis 9,10 methylene								
C18 cis 9,10 methylene								
Others	1	0	3	2	7	5	4	3
Total	100	100	100	100	100	100	100	100